King's Research Portal

DOI:

[10.1016/S0140-6736\(17\)30823-1](https://doi.org/10.1016/S0140-6736(17)30823-1)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Prague, J. K., Roberts, R. E., Comninou, A. N., Clarke, S., Jayasena, C. N., Nash, Z., Doyle, C., Papadopoulou, D. A., Bloom, S. R., Mohideen, P., Panay, N., Hunter, M. S., Veldhuis, J. D., Webber, L. C., Huson, L., & Dhillon, W. S. (2017). Neurokinin 3 receptor antagonism as a novel treatment for menopausal hot flushes: A phase 2, randomised, double-blind, placebo-controlled trial. *The Lancet*, 389, 1809-1820. [https://doi.org/10.1016/S0140-6736\(17\)30823-1](https://doi.org/10.1016/S0140-6736(17)30823-1)

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Neurokinin 3 receptor antagonism as a novel treatment for menopausal hot flushes: a phase 2, randomised, double-blind, placebo-controlled trial

Julia K Prague, Rachel E Roberts, Alexander N Comninou, Sophie Clarke, Channa N Jayasena, Zachary Nash, Chedie Doyle, Deborah A Papadopoulou, Stephen R Bloom, Pharis Mohideen, Nicholas Panay, Myra S Hunter, Johannes D Veldhuis, Lorraine C Webber, Les Huson, Waljit S Dhillon

Summary

Background Hot flushes affect 70% of menopausal women and often severely impact physical, psychosocial, sexual, and overall wellbeing. Hormone replacement therapy is effective but is not without risk. Neurokinin B signalling is increased in menopausal women, and has been implicated as an important mediator of hot flushes.

Methods This phase 2, randomised, double-blind, placebo-controlled, single-centre, crossover trial assessed the effectiveness of an oral neurokinin 3 receptor antagonist (MLE4901) on menopausal hot flushes. Eligible participants were healthy women aged 40–62 years, having seven or more hot flushes in every 24 h of which some were reported as being severe or bothersome, who had not had a menstrual period for at least 12 months, and who had not been taking any medication shown to improve menopausal flushes in the preceding 8 weeks. Participants received 4 weeks of MLE4901 (40 mg, orally, twice daily) and placebo (orally, twice daily) in random order separated by a 2 week washout period. Randomisation was completed by a central computer, and participants were allocated to treatment number in numerical order. The primary outcome was the total number of hot flushes during the final week of both treatment periods. Analyses were by intention to treat and per protocol using generalised linear mixed models and standard crossover analysis. All analyses were prespecified in the study protocol. The trial is registered at ClinicalTrials.gov, number NCT02668185.

Findings 68 women were screened between Feb 3 and Oct 10, 2016, of which 37 were randomly assigned and included in an intention-to-treat analysis. 28 participants completed the trial and were included in a per-protocol analysis. MLE4901 significantly reduced the total weekly number of hot flushes by 45 percentage points (95% CI 22–67) compared with the placebo (intention-to-treat adjusted means: placebo 49·01 [95% CI 40·81–58·56] vs MLE4901 19·35 [15·99–23·42]; adjusted estimate of difference 29·66 [17·39–42·87], $p < 0·0001$). Treatment was well tolerated. Three participants developed a transaminase rise (alanine aminotransferase 4·5–5·9 times the upper limit of normal) with a normal bilirubin 28 days after starting MLE4901, which normalised within 90 days.

Interpretation Treatment with a neurokinin 3 receptor antagonist (MLE4901) could be practice changing as it safely and effectively relieves hot flush symptoms without the need for oestrogen exposure. Larger scale studies of longer duration are now indicated.

Funding UK Medical Research Council and National Institute for Health Research.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license.

Introduction

The menopause occurs when an insufficient number of ovarian follicles remain to sustain circulating oestrogen concentrations. Consequently, menstruation ceases, fertility is lost, and in most women a cluster of symptoms become prominent, which impact physical, psychosocial, sexual, and overall wellbeing. Symptoms can be long lasting (median 7·4 years),¹ and most women report that vasomotor symptoms (termed hot flushes or flashes or night sweats) are the most bothersome of all,² with at least 10% of women reporting them as intolerable.³ An effective treatment is hormone replacement therapy as this artificially restores circulating oestrogen. However, hormone replacement

therapy and in particular combined hormone replacement therapy is not without risk, and is contraindicated in many women due to the long-term safety concerns, including an increased risk of breast⁴ and ovarian⁵ cancers, thromboembolism,⁴ and stroke.⁴ Other treatments including selective serotonin reuptake inhibitors,⁶ gabapentin,⁶ tibolone,⁶ and cognitive behavioural therapy⁷ have been shown to have some efficacy but can cause side-effects. Herbal remedies, such as black cohosh and red clover might provide some relief but efficacy is variable between trials.⁸ It is estimated that a novel treatment for menopausal flushes could currently benefit 10 million women in the UK alone,⁹ and therefore a better understanding of the

Lancet 2017; 389: 1809–20

Published Online

April 3, 2017

[http://dx.doi.org/10.1016/S0140-6736\(17\)30823-1](http://dx.doi.org/10.1016/S0140-6736(17)30823-1)

S0140-6736(17)30823-1

See [Comment](#) page 1775

Department of Investigative Medicine, Imperial College London, London, UK

(J K Prague MBBS, R E Roberts MBBS, A N Comninou PhD, S Clarke MBBS, C N Jayasena PhD, Z Nash MBBS, C Doyle BSN, D A Papadopoulou BSc, Prof S R Bloom MD, Prof W S Dhillon PhD); Millendo Therapeutics, Inc, Ann Arbor, MI, USA (P Mohideen MD);

Department of Gynaecology, Queen Charlotte's & Chelsea Hospital and Chelsea and Westminster Hospital, London, UK (N Panay MRCOG); Institute of Reproductive and Developmental Biology, Imperial College London, London, UK (N Panay); Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

(Prof M S Hunter PhD); Mayo Clinic, Rochester, MI, USA (Prof J D Veldhuis MD); Scientific Partnering & Alliances, Innovative Medicines and Early Development Biotech Unit, AstraZeneca, Melbourne, UK (L C Webber MBBS); and Division of Experimental Medicine, Imperial College London, London, UK (L Huson PhD)

Correspondence to: Prof Waljit S Dhillon, Department of Investigative Medicine, 6th Floor, Commonwealth Building, Imperial College London, Hammersmith Campus, Du Cane Road, London, W12 0NN, UK
w.dhillon@imperial.ac.uk

Research in context

Evidence before this study

Hot flushes affect 70% of menopausal women and result in long-lasting symptoms, which severely impact physical, psychosocial, sexual, and overall wellbeing, in response to oestrogen withdrawal. Hormone replacement therapy is an effective treatment but is contraindicated in some women, and is not without risk in all due to long-term safety concerns, particularly an increased risk of breast cancer (though perhaps less so if it is oestrogen-only hormone replacement therapy), thromboembolism, and stroke. Alternative treatments—eg, some antidepressants and herbal remedies, have been shown to have some efficacy but not always greater than the effect of a placebo depending on the study and treatment class, and are not without side-effects too. For the past 20 years a growing body of evidence has accumulated that implicates neurokinin B (NKB) signalling (a hypothalamic neuropeptide) together with its receptor (NK3R) in the aetiology of menopausal hot flushes. We searched PubMed on Jan 6, 2017, using the keywords “neurokinin 3 receptor”, “hot flushes”, and “hot flashes” with no date or language restrictions. Of the four publications identified, one was a comprehensive review summarising all previous evidence in rodents, primates, and human post-mortem studies that shows that NKB neurons adapt in response to sex steroid deficiency, that this can be reversed by sex steroid replacement, and that NKB signalling is propagated via the hypothalamic median pre-optic nucleus (MnPO), which expresses NK3R, and receives input from, and projects to, the autonomic thermoregulatory pathway. This review included one of the other original articles identified. Of the other two, one was an original article that further supports the role of NKB signalling in the MnPO, and the other was a randomised,

double-blind, placebo-controlled, crossover trial that showed that peripheral infusion of NKB intravenously induces hot flushes in healthy premenopausal women that are typical of the location, duration, and observed physiological change of those described by postmenopausal women. An original article (not identified from the search) of a genome-wide association study found that genetic variation in *TAC3R*, which is the gene that encodes NK3R, might account for the variability in experience of hot flushes reported among women.

Added value of this study

We provide evidence from a phase 2, randomised, placebo-controlled, crossover trial that treatment with an oral, neurokinin 3 receptor antagonist taken twice daily significantly reduces the frequency of menopausal hot flushes. Residual flushes are also significantly less severe, less bothersome, and less interfering than when taking placebo. Treatment was well tolerated and safe. The only safety caution was a mild transient rise in transaminase concentrations in a small subgroup of participants.

Implications of all the available evidence

The finding that pharmacological blockade of NKB signalling with an oral NK3R antagonist can significantly improve hot flush symptoms independently of any hormonal effect fits entirely with the pre-existing data, and suggests great promise for such agents as a novel therapy. Larger scale studies of longer duration will determine whether such a therapeutic approach will change future clinical practice so that the lives of those women so deeply affected by hot flushes can be transformed without the need for increased oestrogen exposure.

aetiology of such flushes and an associated targeted therapeutic is required.

A hot flush is characterised by an intense feeling of heat, which often rises through the body, and intermittent activation of heat dissipation effectors, including peripheral cutaneous vasodilatation, sweating, and behavioural change, to reduce temperature.¹⁰ In rodents this is seen as a change in tail skin temperature secondary to vasodilatation of tail vessels, and behaviour change to increase heat loss such as grooming and hyperventilation.¹⁰ In human beings, a symptom diary and skin conductance monitor can reflect similar changes.^{11,12}

It has been known for many years that the hypothalamic-pituitary-gonadal axis regulates circulating sex steroid concentrations throughout life but it is only more recently that the role of the hypothalamic neurons containing colocalised kisspeptin, neurokinin B (NKB), and dynorphin receptors (so called KNDy neurons) has become clear.¹³ Furthermore, over the past 20 years, Rance and colleagues have contributed to a growing body of

evidence in rodents,^{14–17} primates,¹⁸ and post-mortem studies in human beings¹⁹ that the KNDy neurons, and in particular NKB and its receptor (NK3R), are implicated in the aetiology of the menopausal hot flush (summarised in a comprehensive review by Rance and colleagues¹⁰). Two recent publications further implicate NKB and NK3R in menopausal flushing: in a randomised, double-blind, placebo-controlled, crossover study, peripheral infusion of NKB intravenously to healthy premenopausal women induced hot flushes that were typical in location and duration to those described by postmenopausal women,²⁰ and Crandall and colleagues²¹ found that genetic variation in *TAC3R*, which is the gene that encodes NK3R, might account for the variability in experience of hot flushes reported among menopausal women. Such data led us to hypothesise that NKB–NK3R signalling was an important mediator of menopausal flushing, and therefore pharmacological blockade of NK3R with an oral antagonist could be a novel therapeutic target without the need for increased oestrogen exposure.

Methods

Study design and participants

This randomised, placebo-controlled, double-blind, two-way crossover trial was designed to assess the efficacy of an oral NK3R antagonist (MLE4901, previously termed AZD4901) in reducing the frequency of menopausal flushes in a proof-of-concept study. Eligible participants were healthy women aged 40–62 years, having seven or more hot flushes in every 24 h of which some were reported as being severe or bothersome, who had not had a menstrual period for at least 12 months, and who had not been taking any medication shown to improve menopausal flushes in the preceding 8 weeks. Full inclusion and exclusion criteria are detailed in the appendix.

Ethical approval was obtained from the West London Regional Ethics Committee (15/LO/1481), and approval was also gained from the Medicine and Healthcare products Regulatory Agency (EudraCT 2015-001553-32). All participants provided written informed consent before inclusion. The study was done in accordance with Good Clinical Practice guidelines. No changes were made to the protocol after study initiation and no interim analysis was done. An independent data monitoring committee monitored the safety of the study. Recruitment was stopped when sufficient numbers had enrolled to ensure the planned sample size was achieved, and the trial was stopped when all participants had completed the study protocol as no serious safety concerns occurred requiring early termination.

Randomisation and masking

Randomisation was provided by Almac Clinical Services using a balanced randomisation design (1:1, MLE4901 or placebo first) made using their in-house automated Medication Number List generator to provide a Medication Number List that linked subject number to a randomised treatment group. Subject number was assigned by JKP in numerical order immediately before

writing each drug prescription and taking it to the pharmacy. All authors, participants, and trial staff including the trial pharmacist and pharmacy technicians were masked until completion of the study. Unmasking only occurred once all participants had completed the study, and all their data had been entered into the electronic case record forms. To ensure safety in the case of a medical emergency, code break packs with individual scratch off panels were held by JKP, WSD, and the trial pharmacist but these were never required, and so remain unopened.

Procedures

For the first 2 weeks of the protocol (baseline period) data were collected to establish participants' steady state, to ensure familiarity with recording symptoms and completing the questionnaires appropriately, and to enable review of the inclusion criteria regarding number and severity of flushes at the end of week 2. Participants were then randomly assigned to either 4 weeks of MLE4901 (40 mg, orally, twice daily; group 1) or exact-match placebo (group 2) followed by a 2 week washout period (half-life of MLE4901 is 8.5 h). The dose of MLE4901 was chosen following a recent study in polycystic ovarian syndrome that showed 40 mg twice daily was biologically active and safe.²² The participants then received 4 weeks of whichever intervention they did not receive first followed by 2 weeks of monitoring to ensure they were monitored for the same length of time after dosing irrespective of treatment assignment order (figure 1).

Participants were ambulatory, and attended one of our clinical research facilities at Imperial College NHS Trust Hospitals. They met with one of the doctors in the study team (almost always the study coordinator [JKP]) every week for 14 weeks. At each weekly visit, adverse events (according to the Common Terminology Criteria for Adverse Events version 4.0 [CTCAE]), vital signs, and liver and kidney function were recorded, and twice-daily

See Online for appendix

Figure 1: Summary of protocol

Baseline period: participants underwent a 2 week period to gather baseline data on hot flush frequency, severity, bother, and interference (Hot Flash Related Daily Interference Scale). If the inclusion criteria regarding hot flush frequency and severity were met at the end of this period then they were assigned to the active phase of the study. Intervention 1 (double-blind): all participants randomly assigned to either 4 weeks of treatment with oral, twice daily 40 mg MLE4901 or exact-match placebo. Washout period: all participants underwent a 2 week washout period after intervention 1 (half-life of MLE4901 is 8.5 h). Intervention 2 (double-blind): all participants then switched to receive either 4 weeks of treatment with oral, twice daily exact-match placebo or oral, twice daily 40 mg MLE4901 depending on which intervention they received first. Monitoring period: a subsequent 2 week period to complete safety monitoring.

questionnaires recording total hot flush frequency, severity, bother, interference, associated menopausal symptoms, and treatment adherence were issued and collected. Additionally, a programmed Bahr Monitor (Simplex Scientific, Middleton, WI, USA) was issued to record skin conductance during the first 48 h of each week,¹² and serum was saved to later measure gonadotropins (see the appendix p 4 for the schedule of trial procedures).

Outcomes

The primary outcome was total number of hot flushes during the fourth week of treatment with MLE4901 or placebo to allow adequate time for an effect to be observed.²³ Comparison was also made between the total number of hot flushes during the fourth week of both treatment periods and the total number during the second week of the baseline period.²³ To ensure accurate records, participants recorded their flushes in real time using either a tally chart on a piece of paper ($n=34$) or an application on their smartphone such as Tally Counter (Pixel Research Labs, Minneapolis-Saint Paul, MN, USA; $n=3$), and then collated their total number of flushes twice daily on waking to record previous overnight symptoms and before bed to record daytime symptoms.

Secondary outcomes were hot flush severity, bother, and interference, associated menopausal symptom domain scores, gonadotropin concentrations, luteinising hormone pulsatility, and number of hot flushes detected by a skin conductance monitor. Hot flush severity and bother were reported twice daily (on waking to capture previous overnight symptoms and before bed to capture daytime symptoms) using a scale that ranged from nil to severe, and from not at all to a lot, respectively. Responses were then converted into a numerical scale for data analysis as per Joffe and colleagues²³ (severity: nil scored 1, mild scored 2, moderate scored 3, and severe scored 4; bother: not at all scored 1, a little scored 2, moderately scored 3, and a lot scored 4), and the morning and evening scores were then added to give a total daily score. Hot flush interference was recorded every evening using the Hot Flash Related Daily Interference Scale as per Carpenter,²⁴ and associated menopausal symptoms were recorded every morning using the Menopause-Specific Quality of Life (MENQOL) questionnaire (which is subdivided into the four symptomatic domains: vasomotor, psychosocial, physical, and sexual) as per Hilditch and colleagues.²⁵ Both of these scores were calculated as advised in the two original papers referenced but on a daily rather than a weekly or monthly basis, and then mean weekly total score was used for data analysis. To objectively measure hot flushes a skin conductance monitor (Bahr Monitor) was worn on the sternum for the first 48 h of each study week and the number of flushes was calculated by the software hot flush detection algorithm (SCM Conductance Software version 1.1.3.0, Simplex Scientific, Middleton, WI, USA).¹² The mean weekly value for each of the secondary outcomes was

compared during the fourth week of treatment with MLE4901 and placebo. Comparison was also made between the fourth week of both treatment periods and the second week of the baseline period. To assess compliance with treatment, participants also completed a modified medication adherence questionnaire twice daily, and returned their tablet bottles and any remaining tablets after both drug treatments for counting.

To measure any effect of MLE4901 on reproductive hormones, 3 mL of blood was taken from a peripheral venepuncture at each study visit to measure luteinising hormone, follicle-stimulating hormone, and oestradiol. Furthermore, to analyse luteinising hormone pulsatility a subgroup of participants based on their willingness to participate ($n=13$) attended our clinical research facility for three, 8 h studies. One was during the baseline period, and one during the second week of both treatment periods. During each study a 3 mL blood sample was taken every 10 min from a peripheral venous cannula that had been sited before the study start (time -30 min). All samples were left to clot for at least 30 min and then centrifuged so that the serum could be frozen for subsequent analysis. Analysis was completed in a batch when participants had finished the entire study protocol to eliminate any intra-assay variation. An automated chemiluminescent immunoassay method (Abbott Diagnostics, Maidenhead, UK) was used for analysis; for reference ranges, coefficients of variation, and analytical sensitivities see the appendix (p 5). Luteinising hormone pulsatility was determined using a blinded deconvolution method with 93% sensitivity and specificity by calculating the number and amplitude of luteinising hormone pulses, and how ordered they were.²⁶

Statistical analysis

The sample size was calculated to examine a reduction in hot flush frequency using published data from studies with similar designs.^{23,27,28} Allowing for a 25% reduction in hot flush frequency with placebo,^{11,29} we anticipated that MLE4901 would result in a 50% reduction from a baseline of seven hot flushes per 24 h since it had been previously determined that this was the magnitude of clinical improvement needed for an effective new therapeutic for menopausal flushing (ie, at least twice the effect achieved by placebo).^{23,27,28} We calculated an intraparticipant hot flush frequency SD of 2.02 hot flushes per day based on raw data from a previous study by Ayers and colleagues.⁷ 27 women were required to complete the protocol (90% power, alpha 0.05, two-sided), and allowing for a 10% dropout rate²⁹ we estimated that 30 women would enter the active phase of the study having met the inclusion criteria at week 2.

Data were analysed using generalised linear mixed models (as implemented in SAS PROC GLIMMIX). For the primary endpoint (weekly total hot flush count), a generalised linear mixed model with a Poisson error structure was used. For all secondary endpoints, except

	Group 1: allocated to MLE4901 and then placebo (n=16)	Group 2: allocated to placebo and then MLE4901 (n=12)
Age (years, range)	56 (50–62)	54 (49–61)
Ethnicity		
White British	8 (50%)	9 (75%)
White European	2 (13%)	0
White other	1 (6%)	0
White unspecified	0	1 (8%)
Asian, Pakistani	1 (6%)	0
Black Caribbean	3 (19%)	2 (17%)
Mixed white and black Caribbean	1 (6%)	0
Body-mass index (kg/m ²)	27.0 (4.2)	24.7 (3.7)
Blood pressure (mm Hg)		
Normotensive	15 (94%)	10 (83%)
Hypertensive	1 (6%)	2 (17%)
Body temperature at screening appointment (°C)	36.4 (0.4)	36.7 (0.4)
Duration of oligomenorrhoea (months)	135 (96)	96 (66)
Duration since last menstrual period (months)	119 (94)	77 (63)
Duration since hot flushes (months)	79 (47)	74 (65)
Menarche (age, years)	13 (1)	13 (2)
Regular menstrual cycles in adulthood		
Regular	13 (81%)	10 (83%)
Irregular	3 (19%)	2 (17%)
Gravida	2 (2)	3 (2)
Parity	1 (1)	2 (2)
Hysterectomy		
No	11 (69%)	12 (100%)
Hysterectomy	4 (25%)	0
Hysterectomy and bilateral salpingo-oophorectomy	1 (6%)	0
History of hormone replacement therapy use		
Yes	5 (31%)	1 (8%)
No	11 (69%)	11 (92%)
History of Mirena coil use		
Yes	2 (12%)	2 (17%)
No	14 (88%)	10 (83%)

(Table 1 continues in next column)

	Group 1: allocated to MLE4901 and then placebo (n=16)	Group 2: allocated to placebo and then MLE4901 (n=12)
(Continued from previous column)		
History of herbal remedy use of any type		
Yes	11 (69%)	6 (50%)
No	5 (31%)	6 (50%)
Smoking status		
Never smoked	13 (82%)	5 (42%)
Ex-smoker	1 (6%)	4 (33%)
Current smoker	2 (12%)	3 (25%)
Alcohol consumption		
≤14 units per week	16 (100%)	12 (100%)
>14 units per week	0	0
Luteinising hormone at screening appointment (IU/L)	31.7 (11.8)	33.2 (7.3)
Follicle-stimulating hormone at screening appointment (IU/L)	70.0 (20.5)	74.9 (23.1)
Progesterone at screening appointment (nmol/L)	0.9 (0.0)	0.9 (0.0)
Oestradiol at screening appointment (pmol/L)	69 (0)	69 (0)
Prolactin at screening appointment (mIU/L)	166 (47)	167 (69)
Androstenedione at screening appointment (nmol/L)	1.5 (0.8)	1.9 (1.0)
Dehydroepiandrosterone at screening appointment (μmol/L)	2.8 (2.1)	3.1 (2.0)
Sex hormone binding globulin at screening appointment (nmol/L)	61 (30)	56 (19)
Extracted testosterone at screening appointment (nmol/L)	0.7 (0.4)	0.8 (0.3)

Data are mean (SD) or n (%), unless stated otherwise.

Table 1: Demographics and baseline characteristics of the 28 women who completed the trial (per-protocol set)

the total number of flushes in luteinising hormone pulsatility analysis, which used a generalised linear mixed model with a Poisson error structure, a generalised linear mixed model with gamma error structure was used since exploratory analysis indicated that the usual assumption of normality was not valid for these endpoints. For all models used, a standard crossover analysis was implemented, with period, administration sequence, and treatment as fixed effects and subject as a

random effect. Tests for sequence (order) and period effects across all our models were done. All possible covariates were initially included in exploratory analyses but the only ones that were significant in each case were the baseline values of the endpoint being analysed, and so the final results reported are taken from models that included the baseline value as the only covariate. All of these models, although not assuming normally distributed errors, are entirely analogous to conventional analyses using ANCOVA and can be interpreted in the same way.

From each model, adjusted (least squares) means and differences between treatment means were estimated, together with associated 95% CIs, and a p value from a comparison of the mean values of the two treatments. For each endpoint, the percentage change from baseline

Figure 2: Trial profile

Intention-to-treat analysis included all participants who were randomly assigned and received study medication (placebo or MLE4901; $n=37$). Per-protocol analysis included all participants who appropriately completed both treatment periods ($n=28$). See the appendix (pp 1–3) for more details about inclusion and exclusion criteria, and the number of participants affected by each.

was also estimated using a generalised linear mixed model (with gamma error structure) with fold-change from baseline as the analysed response, with percentage change (and 95% CI) derived by calculation from the fold-change results. This ensures that the percentage change results take into account the crossover design.

All of these analyses were prespecified as per the study protocol. The primary and all secondary endpoints were analysed using both the per-protocol and intention-to-treat analysis sets, and all p values and adjusted means for all endpoints were very similar. Accordingly, results have been presented for the intention-to-treat and per-protocol sets for the primary endpoint and for the per-protocol set only for the secondary endpoints as originally planned (though intention-to-treat analyses for the secondary outcomes as well as the per-protocol analysis for the primary outcome are reported in the appendix). To facilitate the intention-to-treat analysis of the primary endpoint, weekly total flush counts were imputed where missing by setting the count totals to be equal on both active and placebo treatment groups. However, we also did a sensitivity analysis of the primary endpoint intention-to-treat analysis using multiple imputation techniques, and this gave similar results to those reported below, with a slightly larger estimate of treatment effect. The multiple imputation method was used for all intention-to-treat analyses of secondary endpoints. All significant p values reported remained significant after adjustment for multiple testing (using the false discovery rate) procedure. The study was registered at ClinicalTrials.gov, number NCT02668185.

Role of the funding source

This was an academic investigator initiated and led study, which was funded by the UK Medical Research Council (grant reference MR/M024954/1) and a National Institute for Health Research Professorship (to WSD, grant reference RP-2014-05-001). AstraZeneca agreed to provide their NK3R antagonist (AZD4901) and exact-matched placebo in a collaboration led by the principal investigator (WSD). The study was sponsored by the principal investigator's host institution (Imperial College London). One employee of AstraZeneca (LCW) contributed to the study design and continued to be involved in the trial management group while the study was ongoing. AstraZeneca held the background intellectual property regarding the compound. AstraZeneca subsequently exclusively licensed the compound to Millendo Therapeutics after the research protocol had been approved, and so one representative from their company (PM) also became involved in the trial management group while the study was ongoing. At this time the compound name was changed from AZD4901 to MLE4901. WSD and JKP (Imperial College London) had full access to all the data from the study and had final responsibility for the content of the report and decision to submit for publication. The data analysis was done by the trial statistician LH (Imperial College London).

Results

68 women were screened by JKP between Feb 3, 2016, and Oct 10, 2016. After discussion, eight did not meet the inclusion criteria, and 11 met an exclusion criterion, and so all 19 were withdrawn (see appendix pp 1–3).

	Group 1: allocated to MLE4901 and then placebo (n=16)	Group 2: allocated to placebo and then MLE4901 (n=12)
Total number of hot flushes per 24 h	13.09 (6.47)	12.56 (3.93)
Hot flush severity	5.74 (1.05)	6.36 (0.96)
Hot flush bother	5.86 (1.24)	6.14 (1.00)
Hot flush interference	48.68 (27.27)	27.38 (19.59)
MENQOL domains score		
Vasomotor	4.25 (1.93)	4.59 (1.57)
Psychosocial	3.50 (1.86)	2.36 (1.11)
Physical	3.24 (1.33)	2.96 (1.32)
Sexual	3.75 (2.54)	2.63 (1.83)
Number of flushes detected by sweat monitor per 24 h	24.86 (7.73)	27.33 (6.14)

Data are unadjusted mean (SD). MENQOL=Menopause-Specific Quality of Life questionnaire.

Table 2: Baseline outcome data presented by treatment assignment group (per-protocol set)

Four further participants withdrew before the study began due to time commitments. Therefore, 45 participants started the trial, of which seven were identified to have not met the inclusion criteria regarding frequency or severity of flushes at the end of week 2 (as per protocol) and so were not randomly assigned and withdrawn. One participant was randomly assigned but withdrew before receiving the allocated intervention due to new employment and so was not included in any analysis. Of the remaining 37 participants who were randomly assigned and received study medication (the intention-to-treat analysis set), 28 completed the protocol (the per-protocol analysis set; age range 49–62 years and had been experiencing hot flushes for 1–16 years). Extended demographic and baseline characteristics of the 28 participants who completed the trial are summarised in table 1, and participant flow is shown in figure 2. Baseline menopausal symptom scores were similar between treatment assignment groups (table 2). Baseline mean total number of hot flushes per 24 h was 13.09 (SD 6.47) in group 1 and 12.56 (3.93) in group 2 (table 2).

MLE4901 significantly reduced the total weekly number of hot flushes compared with placebo (intention-to-treat adjusted means: placebo 49.01 [95% CI 40.81–58.56] vs MLE4901 19.35 [15.99–23.42], $p<0.0001$; 45 percentage point decrease [22–67]; figure 3, table 3). The corresponding results for the per-protocol analysis were similar (placebo 59.27 [95% CI 51.52–68.17] vs MLE4901 16.85 [14.34–19.78], $p<0.0001$; 52 percentage point decrease [26–82]; appendix p 6).

Compared with placebo, MLE4901 also significantly reduced weekly hot flush severity (placebo

Figure 3: Primary endpoint ITT analysis

(A) Whole group ITT analysis (n=37) irrespective of treatment assignment order using adjusted means from crossover analysis with 95% CIs: percentage change in hot flush frequency (total number of hot flushes) during the final week of the 4 week treatment period with MLE4901 and placebo compared with hot flush frequency (total number of hot flushes) during the final week of the 2 week baseline period. Statistical analysis incorporated a total of seven daily counts for each of the study weeks analysed, and is based on a crossover model including treatment and period as fixed effects, subject as a random effect (within sequence), and baseline flush count as a covariate. All other possible demographic covariates were tested in the model but none were significant and therefore all were excluded from the final model. The model used is a generalised linear model with gamma error structure. Tests for sequence (order), and period, effect across all our models confirmed neither were significant. (B) Subgroup ITT analysis (n=37) by treatment assignment group using participants' unadjusted (raw) data with 95% CIs: percentage change in hot flush frequency (total number of hot flushes) during the final week of the 4 week treatment period with MLE4901 and placebo compared with hot flush frequency (total number of hot flushes) during the final week of the 2 week baseline period depending on whether the participant received MLE4901 or placebo as the first or second intervention. ITT=intention to treat.

5.70 [5.09–6.38] vs MLE4901 3.27 [2.92–3.66], $p<0.0001$; 41 percentage point decrease [32–49]), bother (placebo 5.56 [4.96–6.22] vs MLE4901 2.92 [2.61–3.27], $p<0.0001$; 45 percentage point decrease [36–53]), and interference (placebo 26.48 [20.02–35.03] vs MLE4901 7.94 [5.76–10.95], $p<0.0001$; 58 percentage point decrease [40–76]; table 4). The reported subjective effect of decreased flushes by MLE4901 was confirmed by objective measurement using the skin conductance

	Baseline	On treatment		Difference
		Placebo	MLE4901	
Unadjusted (raw) mean counts	84.54	62.84	25.24	..
Adjusted (least squares) means from crossover analysis (95% CI)*	..	49.01 (40.81 to 58.86)	19.35 (15.99 to 23.42)	..
Adjusted (least squares) estimate of difference (placebo–MLE4901) between treatment means (95% CI)	..	..	..	29.66 (17.39 to 42.87)
Adjusted (least squares) estimate of percentage change from baseline (95% CI)	..	–28% (–17 to –39)	–73% (–61 to –84)	..
Adjusted (least squares) estimate of percentage point difference (MLE4901–placebo) between treatment means (95% CI)	..	..	..	–45% (–22 to –67)

Total number of hot flushes during the final week of the 4 week treatment period with MLE4901 or placebo. Comparison was also made of the total number of hot flushes during the second week of the baseline period. Statistical analysis incorporated a total of seven daily counts for each of the study weeks analysed and is based on a crossover model including treatment and period as fixed effects, subject as a random effect (within sequence), and baseline flush count as a covariate. All other possible demographic covariates were tested in the model but none were significant and therefore all were excluded from the final model. The model used is a generalised linear model with Poisson error structure for the total flush counts, and with gamma error structure to estimate percentage change from baseline for the two treatments. *p<0.0001, comparison of treatment means.

Table 3: Primary endpoint (intention-to-treat analysis, n=37)

	Placebo (adjusted mean)	MLE4901 (adjusted mean)	Adjusted estimate of percentage point difference between treatment means	p value comparing adjusted treatment means
Hot flush severity	5.70 (5.09 to 6.38)	3.27 (2.92 to 3.66)	–41% (–32 to –49)	<0.0001
Hot flush bother	5.56 (4.96 to 6.22)	2.92 (2.61 to 3.27)	–45% (–36 to –53)	<0.0001
Hot flush interference	26.48 (20.02 to 35.03)	7.94 (5.76 to 10.95)	–58% (–40 to –76)	<0.0001
MENQOL domain score				
Vasomotor	3.98 (3.38 to 4.69)	2.05 (1.74 to 2.42)	–45% (–33 to –58)	<0.0001
Psychosocial	2.58 (2.30 to 2.90)	2.18 (1.94 to 2.45)	–15% (–5 to –25)	0.0083
Physical	2.93 (2.63 to 3.27)	2.42 (2.17 to 2.69)	–19% (–9 to –28)	0.0002
Sexual	2.15 (1.84 to 2.51)	1.98 (1.68 to 2.30)	–8% (5 to –23)	0.24
Number of flushes detected by sweat monitor per 24 h	26.91 (23.16 to 31.27)	16.22 (13.99 to 18.80)	–43% (–30 to –55)	<0.0001

All outcomes were compared during the final week of the 4 week treatment period with MLE4901 and exact-match placebo to allow adequate time for an effect to be observed. For all endpoints, a generalised linear model with gamma error structure was used. A standard crossover analysis was implemented, with period, administration sequence, and treatment as fixed effects and subject as a random effect. All possible baseline covariates were initially included in exploratory analyses but the only covariates that were significant, in each case, were the baseline values of the endpoint being analysed. All other demographic covariates were therefore excluded from the final model. From each model, adjusted (least squares) means and differences between means were estimated, together with associated 95% CIs. p values refer to the comparison of the mean values of the two treatments (placebo and MLE4901). MENQOL=Menopause-Specific Quality of Life questionnaire.

Table 4: Summary results of the secondary outcomes for the 28 women who completed the trial (per-protocol analysis set)

monitor and detection algorithm software (mean number of hot flushes per 24 h: placebo 26.91 [95% CI 23.16–31.27] vs MLE4901 16.22 [13.99–18.80], p<0.0001; 43 percentage point decrease [30–55]; table 4).

Mean weekly MENQOL domain scores showed that vasomotor symptoms improved when taking MLE4901 compared with placebo (placebo 3.98 [95% CI 3.38–4.69] vs MLE4901 2.05 [1.74–2.42], p<0.0001) as did the psychosocial domain score (placebo 2.58 [2.30–2.90] vs MLE4901 2.18 [1.94–2.45], p=0.0083), and physical domain score (placebo 2.93 [2.63–3.27] vs MLE4901 2.42 [2.17–2.69], p=0.0002; table 4). There was no significant difference between the sexual domain score when taking MLE4901 compared with when taking placebo (placebo 2.15 [95% CI 1.84–2.51] vs MLE4901 1.98 [1.68–2.30], p=0.24).

The magnitude of the placebo effect did vary between individuals but the variation in the treatment effect with MLE4901 was much smaller (tables 3 and 4).

Luteinising hormone pulsatility analysis showed that treatment with MLE4901 did not change the number of luteinising hormone pulses compared with placebo (placebo 6.30 [95% CI 4.92–8.05] vs MLE4901 5.48 [4.21–7.13], p=0.41) but did increase the amplitude of pulses (placebo 16.16 [11.03–23.67] vs MLE4901 26.66 [18.20–39.05], p=0.0243) and improved their orderliness (placebo 0.84 [0.75–0.95] vs MLE4901 0.59 [0.53–0.66], p=0.0006; figure 4). Oestradiol remained unchanged throughout.

Data for the per-protocol analysis of the primary outcome are shown in the appendix (p 6). Data for both analysis sets of secondary outcomes are shown in the

appendix (pp 7–25). Tests for sequence (order), and period, effect across all our models confirmed neither were significant (appendix p 26).

Compliance with treatment was high and both treatments were well tolerated. No serious adverse events occurred. Three participants developed a transient transaminase rise (alanine aminotransferase [ALT] greater than aspartate aminotransferase [AST]) with a normal bilirubin following treatment with MLE4901 (day 28), which remained asymptomatic, and returned to baseline in all cases. None exceeded 5·9 times the upper limit of normal (ULN) for ALT (4·5–5·9 times ULN) or 2·9 times the ULN for AST (1·7–2·9 times ULN; appendix pp 27–28). A small number of minor adverse events occurred in individual participants that were mostly unrelated to a drug effect—eg, grade 1 bruising to great toe after trauma (table 5).

Compliance with the protocol was also high; only one study visit (0·83%) was missed out of 120 in total (due to suspension of the train service required for the participant to attend her weekly visit), and only three participants had suboptimal sweat monitor data (two due to a localised skin irritation caused by the topical electrode plaster and one due to losing the monitor while wearing it). Two participants unintentionally missed more than three doses of the intervention, one of which was due to bereavement, and so were both withdrawn from the study as per protocol. The dropout rate was 24% (nine of 37), which was higher than anticipated, and was mostly due to participants choosing to book a holiday outside the UK for more than 7 days during the study period.

Discussion

In this randomised, double-blind, crossover study we have shown that an NK3R antagonist (MLE4901) significantly reduced the mean weekly number, severity, bother, and interference of hot flushes, which otherwise negatively impact a menopausal woman in her daily life. This result was achieved with a well tolerated, twice daily, oral preparation. Associated psychosocial and physical symptoms also improved with treatment (particularly those related to improved sleep such as fatigue and irritability) as otherwise disruptive night-time flushes were significantly reduced. The only safety caution was an asymptomatic rise in transaminase concentrations in a small subgroup of participants that was not associated with a rise in bilirubin. The results of ongoing and future clinical trials of MLE4901 will further explore this effect and characterise any appropriate follow-up studies required.

This study was designed as a 4 week crossover trial because it was a proof-of-concept study enabling the inclusion of fewer participants to achieve our sample calculation, a shorter trial duration,³⁰ and more importantly ensured each participant acted as their own control for treatment comparisons. Baseline data generated before randomisation allowed further treatment comparisons to

Figure 4: Luteinising hormone pulse analysis

JDV used a blinded deconvolution method with 93% sensitivity and specificity to analyse luteinising hormone pulsatility by calculating the number of luteinising hormone pulses (A), the mean amplitude of luteinising hormone pulses (B), and the orderliness of the pulses (approximate entropy; the lower the number the more ordered the pulses are, with zero denoting perfect orderliness; C). A generalised linear model was used for analysis with a Poisson error structure for number of pulses, and with a gamma structure for mean amplitude and orderliness of luteinising hormone pulses. A standard crossover analysis was implemented, with period, administration sequence, and treatment as fixed effects and subject as a random effect. Box plots: line, median; box, IQR; whiskers extend to the extremes of the data (minimum and maximum values).

	Baseline	Placebo	MLE4901
Respiratory, thoracic, and mediastinal disorders			
Upper respiratory infection (grade 1)	0	3	6
Nervous system disorders			
Headache (grade 1)	0	2	3
Nervous system disorder, other, migraine (grade 2)	0	2	3
Nervous system disorder, other, S1 shingles (grade 1)	0	1	0
Sinus pain (grade 1)	0	0	1
Dysesthesia (grade 1)			
Lip and nose	0	0	1
Jaw and mouth	0	0	1
Gastrointestinal disorders			
Dry mouth (grade 1)	0	0	2
Nausea (grade 1)	0	1	2
Vomiting (grade 1)	0	0	1
Diarrhoea (grade 1)	0	0	2
Constipation (grade 1)	0	1	0
Gastro-esophageal reflux disease (grade 1)	0	0	1
Investigations			
ALT increased (grade 1) with normal AST	1	1	0
AST increased (grade 1) with normal ALT	1	0	1
ALT increased (grade 1) with AST increased (grade 1)	0	1	3
ALT increased (grade 2) with AST increased (grade 1)	1	0	1
ALT increased (grade 3) with AST increased (grade 1)	0	0	2
Blood bilirubin increased (grade 1)	3	0	0
Alkaline phosphatase increased (grade 1)	1	0	1
Creatinine increased (grade 1)	1	1	0
Musculoskeletal and connective tissue disorders			
Chest wall pain (grade 1)	0	1	0
Bruising, great toe after trauma (grade 1)	0	1	0
Neck pain, after road traffic collision (grade 2)	0	1	0
Buttock pain, traumatic (grade 1)	0	0	1
Localised oedema, fingers (grade 1)	0	0	1
Myalgia			
Traumatic (grade 1)	0	0	1
Atraumatic (grade 1)	0	1	1
Skin and subcutaneous tissue disorders			
Hypertrichosis, chin (grade 1)	0	0	1
Skin and subcutaneous tissue disorders, other, weak nails (grade 1)	0	0	1
Pruritus			
Breast (grade 1)	0	0	1
Hands and feet, known eczema (grade 1)	0	0	1
Ear and labyrinth disorders			
Ear and labyrinth disorders, other, blocked ears (grade 1)	0	0	1

Number of events recorded during the study period in all participants who received at least one dose of study medication (placebo or MLE4901; n=37). All participants were asked about adverse events at each weekly visit by the study doctor. Any reported symptom was recorded and then coded according to the Common Terminology Criteria for Adverse Events version 4.0. Counting rules were used to determine group assignment for events within the crossover trial. Group assignment: baseline, from screening through to just before the first dose of the study medication (placebo or MLE4901); placebo, from first dose of placebo to either just before the first dose of MLE4901 (if placebo received first) or through to the end of follow-up (if placebo received second); MLE4901, from first dose of MLE4901 to either just before the first dose of placebo (if MLE4901 received first) or through to the end of follow-up (if MLE4901 received second). ALT=alanine aminotransferase. AST=aspartate aminotransferase.

Table 5: Adverse events

be made and baseline covariates to be used to reduce error variance. We allowed for a 25% reduction in hot flush frequency with placebo at 4 weeks based on previously published studies in menopausal flushing with a similar methodology to ours:^{11,29} our observed rate was a 28% reduction with placebo, which therefore fits closely with the existing literature.^{11,29} However, we acknowledge that there are risks of bias in crossover studies and so we avoided these by ensuring those recruited had chronic and stable flushes in the months preceding enrolment, randomisation, and blinding procedures, and the duration of the treatment and washout periods to eliminate a carryover effect. To ensure adverse events were appropriately assigned, counting rules were used to assess for treatment-emergent events. Intention-to-treat and per-protocol analyses were compared to ensure loss of participants or missing imputation methods did not distort our findings. All reported analyses conformed to the requirements of a crossover trial, and followed common practice for this study design. The objective reduction in hot flushes seen at 4 weeks using the skin conductance monitor with hot flush detection algorithm suggests that the duration of the treatment was sufficient to prove the concept. Furthermore, concordance was seen between subjective reporting and objective measurement of symptoms (45 percentage point decrease vs 43 percentage point decrease in hot flushes, respectively, with MLE4901 compared with placebo).

The findings from this ambulatory study should be generalisable as we placed no restriction on lifestyle, our cohort had a range of symptom duration, a quarter were not white, and some were smokers. Furthermore, no participants had received any treatment that could have improved their hot flushes for at least 8 weeks before study start, and therefore any treatment effect seen is likely to have been caused by the NK3R antagonist. Interestingly, a substantial proportion of our cohort had previously tried herbal remedies with far fewer having tried hormone replacement therapy either due to a contraindication or more commonly due to concerns regarding its use.

However, the trial is also not without limitations. The number of participants was small, the treatment duration was short, and the dropout rate was higher than anticipated. A larger and longer trial is needed to establish whether the treatment effect is long lasting in a greater number of individuals. The study was done in a single centre, and so this could be a potential source of bias. However, as a result all participants were seen and directed by the same trial doctor (JKP) during almost all of the study visits minimising variability in instructions given between participants and maximising compliance. Weekly questionnaire data regarding the primary and secondary outcomes were entered into the electronic clinical record forms by CD, RER, or ZN to reduce the likelihood of JKP observing the original data, or patterns of results that could influence opinion regarding treatment intervention when seeing the participants

before the end of the study. For the same reason, data were not collated and blood samples for hormone concentrations were not analysed until participants had finished the study. Alcohol intake has previously been found to exacerbate menopausal flushes but as alcohol dependence was in our exclusion criteria all participants consumed equal to, or less than, 14 units per week. Increased body-mass index has also previously been associated with increased flushing but was not found to be an independent covariate in our analysis; this might be due to the small study size. As many of the participants were widowed, separated, or not currently sexually active any improvement in the sexual MENQOL domain by treatment with MLE4901 could have been missed as these contributory factors remained unchanged throughout the study.

Over the past 20 years a growing body of evidence has implicated NKB–NK3R signalling in the aetiology of menopausal hot flushes. By studying brain specimens at post mortem, Rance and Young¹⁹ initially showed that in postmenopausal women, hypothalamic neurons are hypertrophied and have increased NKB gene expression and neuronal activity compared with premenopausal women. In keeping with this, the same was found to be true in ovariectomised monkeys but, moreover, this change could be reversed by treatment with sex steroid replacement, thus suggesting this was a dynamic change in response to circulating concentrations of oestradiol as occurs in the menopause.¹⁸ Subsequent work in rats highlighted the importance of the hypothalamic pre-optic nucleus (MnPO) in the propagation of the NKB-mediated signal that results in hot flushes. The MnPO is a neural area that receives input from, and projects to, the autonomic thermoregulatory pathway and expresses NK3R,^{14–17} and hence results in activation of heat dissipation effectors that characterise hot flushes. Furthermore, in a randomised, double-blind, placebo-controlled crossover study in healthy premenopausal women, peripheral infusion of NKB intravenously induced hot flushes that were typical in location, duration, and observed physiological change to those described by postmenopausal women.²⁰ Additionally, Crandall and colleagues²¹ recently found that genetic variation in *TACR3*, which is the gene that encodes NK3R, might account for the variability in experience of hot flushes reported among women. Therefore, the finding that pharmacological blockade of NKB signalling with an oral NK3R antagonist can significantly improve hot flush symptoms independent of any hormonal effect fits entirely with this pre-existing data. This finding suggests great promise for such agents as a novel therapeutic target to change future clinical practice so that the lives of those women so deeply affected by hot flushes could be transformed without the need for increased oestrogen exposure. Larger scale studies of longer duration are the required next step in assessing the feasibility and likelihood of this.

Contributors

ANC, CNJ, SRB, NP, MSH, LCW, LH, and WSD collated and submitted the UK Medical Research Council (MRC) Developmental Pathway Funding Scheme (DPFS) grant application. JKP, LCW, NP, MSH, LH, and WSD designed the final trial protocol. The trial management group comprised JKP, WSD, LCW, and PM. Almac Clinical Services provided the randomisation codes and tablet packs. JKP coordinated the trial, and was supported by RER, SC, CD, and DAP. Data entry was completed by RER, CD, and ZN. Blood samples for gonadotropins and oestradiol were analysed by SC. JDV did the blinded deconvolution analysis for luteinising hormone pulsatility. Data accuracy was checked by ZN, ANC, and WSD. JKP and RER did the analysis of the clinical data. JKP and WSD did the analysis of the safety data. JKP and WSD interpreted the data. LH did the statistical analysis. JKP wrote the first draft of the manuscript and all authors contributed to the further revisions and subsequent approval of the final version submitted for publication.

Declaration of interests

JKP is funded by the MRC DPFS. RER, SC, CD, and DAP are funded by the National Institute for Health Research (NIHR). CNJ is an inventor on a patent application 14762086.8-1453, which is registered to Imperial Innovations. PM is an employee of Millendo Therapeutics. NP has lectured for and acted in an advisory capacity for Abbott, Bayer, Besins, Consilient, Meda, MSD, Mylan, Novo Nordisk, Pfizer, and Shionogi. LCW is a full-time employee of AstraZeneca UK. AstraZeneca has now licensed the compound and associated patents to Millendo Therapeutics (WO2014170648A1, pending; US9475773 B2, granted). WSD is funded by the MRC DPFS and NIHR, is an inventor on a patent application 14762086.8-1453 (which is registered to Imperial Innovations), and is an investigator for a separate study of MLE4901 in polycystic ovarian syndrome, for which a consultancy fee was paid. Millendo Therapeutics provided some financial support for an administrative assistant for the study of MLE4901 in menopausal flushing. ANC, ZN, SRB, MSH, JDV, and LH declare no competing interests.

Acknowledgments

The study was funded by the MRC as part of their DPFS (grant reference MR/M024954/1) and an NIHR Professorship (to WSD; grant reference RP-2014-05-001). The Section of Investigative Medicine is funded by grants from the MRC and Biotechnology and Biological Sciences Research Council, and is supported by the NIHR Imperial Biomedical Research Centre Funding Scheme. The research study was supported by the NIHR/Wellcome Trust Clinical Research Facility at Imperial College Healthcare NHS Trust. JKP is funded by the MRC. RER is funded by an NIHR Academic Foundation Programme award. ANC is funded by the NHS and Biomedical Research Centre. SC is funded by NIHR. CD and DAP are supported by the NIHR. SRB is supported by an NIHR Senior Investigator Award and the MRC. WSD is funded by an NIHR Professorship. We also thank Vincenzo Libri (Director of the NIHR UCLH Clinical Research Facility and Head of the Leonard Wolfson Experimental Neurology Centre at UCL—Institute of Neurology), Tricia Tan (Imperial College London), and Niamh Martin (Imperial College London) for their time and expertise in monitoring the safety of the trial. The views expressed are those of the authors and not necessarily those of the above mentioned funders, the NHS, the NIHR, or the Department of Health.

References

- 1 Avis NE, Crawford SL, Greendale G, et al. Duration of menopausal vasomotor symptoms over the menopause transition. *JAMA Intern Med* 2015; **175**: 531–39.
- 2 Carpenter JS, Woods NF, Otte JL, et al. MsFLASH participants' priorities for alleviating menopausal symptoms. *Climacteric* 2015; **18**: 859–66.
- 3 Stearns V, Ullmer L, Lopez JF, Smith Y, Isaacs C, Hayes D. Hot flushes. *Lancet* 2002; **360**: 1851–61.
- 4 Marjoribanks J, Farquhar C, Roberts H, Lethaby A, Lee J. Long-term hormone therapy for perimenopausal and postmenopausal women. *Cochrane Database Syst Rev* 2017; **1**: CD004143.
- 5 Collaborative Group on Epidemiological Studies of Ovarian Cancer, Beral V, Gaitskell K, Hermon C, Moser K, Reeves G, Peto R. Menopausal hormone use and ovarian cancer risk: individual participant meta-analysis of 52 epidemiological studies. *Lancet* 2015; **385**: 1835–42.

- 6 Drewe J, Bucher KA, Zahner C. A systematic review of non-hormonal treatments of vasomotor symptoms in climacteric and cancer patients. *Springerplus* 2015; **4**: 65.
- 7 Ayers B, Smith M, Hellier J, Mann E, Hunter MS. Effectiveness of group and self-help cognitive behavior therapy in reducing problematic menopausal hot flushes and night sweats (MENOS 2): a randomized controlled trial. *Menopause* 2012; **19**: 749–59.
- 8 Franco OH, Chowdhury R, Troup J, et al. Use of plant-based therapies and menopausal symptoms: a systematic review and meta-analysis. *JAMA* 2016; **315**: 2554–63.
- 9 Office for National Statistics. UK census. 2011. <https://www.ons.gov.uk/census/2011census/2011censusdata> (accessed Jan 7, 2017).
- 10 Rance NE, Dacks PA, Mittelman-Smith MA, Romanovsky AA, Krajewski-Hall SJ. Modulation of body temperature and LH secretion by hypothalamic KNDy (kisspeptin, neurokinin B and dynorphin) neurons: a novel hypothesis on the mechanism of hot flushes. *Front Neuroendocrinol* 2013; **34**: 211–27.
- 11 Sloan JA, Loprinzi CL, Novotny PJ, Barton DL, Lavoie BI, Windschitl H. Methodologic lessons learned from hot flash studies. *J Clin Oncol* 2001; **19**: 4280–90.
- 12 Bahr DE, Webster JG, Grady D, et al. Miniature ambulatory skin conductance monitor and algorithm for investigating hot flash events. *Physiol Meas* 2014; **35**: 95–110.
- 13 Lehman MN, Coolen LM, Goodman RL. Minireview: kisspeptin/neurokinin B/dynorphin (KNDy) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinology* 2010; **151**: 3479–89.
- 14 Mittelman-Smith MA, Williams H, Krajewski-Hall SJ, McMullen NT, Rance NE. Role for kisspeptin/neurokinin B/dynorphin (KNDy) neurons in cutaneous vasodilatation and the estrogen modulation of body temperature. *Proc Natl Acad Sci USA* 2012; **109**: 19846–51.
- 15 Nakamura K, Morrison SF. A thermosensory pathway mediating heat-defense responses. *Proc Natl Acad Sci USA* 2010; **107**: 8848–53.
- 16 Dacks PA, Krajewski SJ, Rance NE. Activation of neurokinin 3 receptors in the median preoptic nucleus decreases core temperature in the rat. *Endocrinology* 2011; **152**: 4894–905.
- 17 Mittelman-Smith MA, Krajewski-Hall SJ, McMullen NT, Rance NE. Neurokinin 3 receptor-expressing neurons in the median preoptic nucleus modulate heat-dissipation effectors in the female rat. *Endocrinology* 2015; **156**: 2552–62.
- 18 Rometo AM, Krajewski SJ, Voytko ML, Rance NE. Hypertrophy and increased kisspeptin gene expression in the hypothalamic infundibular nucleus of postmenopausal women and ovariectomized monkeys. *J Clin Endocrinol Metab* 2007; **92**: 2744–50.
- 19 Rance NE, Young WS 3rd. Hypertrophy and increased gene expression of neurons containing neurokinin-B and substance-P messenger ribonucleic acids in the hypothalami of postmenopausal women. *Endocrinology* 1991; **128**: 2239–47.
- 20 Jayasena CN, Cominos AN, Stefanopoulou E, et al. Neurokinin B administration induces hot flushes in women. *Sci Rep* 2015; **5**: 8466.
- 21 Crandall CJ, Manson JAE, Hohnsbee C, et al. Association of genetic variation in the tachykinin receptor 3 locus with hot flashes and night sweats in the Women's Health Initiative Study. *Menopause* 2017; **24**: 252–61.
- 22 George JT, Kakkar R, Marshall J, et al. Neurokinin B receptor antagonism in women with polycystic ovary syndrome: a randomized, placebo-controlled trial. *J Clin Endocrinol Metab* 2016; **101**: 4313–21.
- 23 Joffe H, Guthrie KA, LaCroix AZ, et al. Low-dose estradiol and the serotonin-norepinephrine reuptake inhibitor venlafaxine for vasomotor symptoms: a randomized clinical trial. *JAMA Intern Med* 2014; **174**: 1058–66.
- 24 Carpenter JS. The Hot Flash Related Daily Interference Scale: a tool for assessing the impact of hot flashes on quality of life following breast cancer. *J Pain Symptom Manage* 2001; **22**: 979–89.
- 25 Hilditch JR, Lewis J, Peter A, et al. A menopause-specific quality of life questionnaire: development and psychometric properties. *Maturitas* 1996; **24**: 161–75.
- 26 Veldhuis JD, Keenan DM, Pincus SM. Motivations and methods for analyzing pulsatile hormone secretion. *Endocr Rev* 2008; **29**: 823–64.
- 27 Freeman EW, Guthrie KA, Caan B, et al. Efficacy of escitalopram for hot flashes in healthy menopausal women: a randomized controlled trial. *JAMA* 2011; **305**: 267–74.
- 28 Newton KM, Carpenter JS, Guthrie KA, et al. Methods for the design of vasomotor symptom trials: the menopausal strategies: finding lasting answers to symptoms and health network. *Menopause* 2014; **21**: 45–58.
- 29 Gordon PR, Kerwin JP, Boesen KG, Senf J. Sertraline to treat hot flashes: a randomized controlled, double-blind, crossover trial in a general population. *Menopause* 2006; **13**: 568–75.
- 30 Loprinzi CL, Diekmann B, Novotny PJ, Stearns V, Sloan JA. Newer antidepressants and gabapentin for hot flashes: a discussion of trial duration. *Menopause* 2009; **16**: 883–87.