



King's Research Portal

DOI:

<https://doi.org/10.1101/065540>
[10.1038/tpj.2017.21](https://doi.org/10.1038/tpj.2017.21)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Verbelen, M., Weale, M. E., & Lewis, C. M. (2017). Cost-effectiveness of pharmacogenetic-guided treatment: are we there yet? *The pharmacogenomics journal*, 17(5), 395-402. <https://doi.org/10.1101/065540>, <https://doi.org/10.1038/tpj.2017.21>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

REVIEW

Cost-effectiveness of pharmacogenetic-guided treatment: are we there yet?

M Verbelen¹, ME Weale² and CM Lewis^{1,2}

Pharmacogenetics (PGx) has the potential to personalize pharmaceutical treatments. Many relevant gene–drug associations have been discovered, but PGx-guided treatment needs to be cost-effective as well as clinically beneficial to be incorporated into standard health-care. We reviewed economic evaluations for PGx associations listed in the US Food and Drug Administration (FDA) Table of Pharmacogenomic Biomarkers in Drug Labeling. We determined the proportion of evaluations that found PGx-guided treatment to be cost-effective or dominant over the alternative strategies, and estimated the impact on this proportion of removing the cost of genetic testing. Of the 137 PGx associations in the FDA table, 44 economic evaluations, relating to 10 drugs, were identified. Of these evaluations, 57% drew conclusions in favour of PGx testing, of which 30% were cost-effective and 27% were dominant (cost-saving). If genetic information was freely available, 75% of economic evaluations would support PGx-guided treatment, of which 25% would be cost-effective and 50% would be dominant. Thus, PGx-guided treatment can be a cost-effective and even a cost-saving strategy. Having genetic information readily available in the clinical health record is a realistic future prospect, and would make more genetic tests economically worthwhile.

The Pharmacogenomics Journal advance online publication, 13 June 2017; doi:10.1038/tpj.2017.21

INTRODUCTION

Pharmacogenetics (PGx) studies the relationship between genetic variation and inter-individual variability in drug response in terms of efficacy and safety. Hence, PGx knowledge can be used to tailor pharmaceutical treatment to the genetic make-up of the patient. Several robust, well-replicated PGx associations exist, for example, the association of *HLA-B*5701* with abacavir hypersensitivity, *HLA-B*1502* with carbamazepine-induced Stevens–Johnson syndrome/toxic epidermal necrolysis, and *VKORC1* and *CYP2C9* with warfarin dosing.^{1–3} Accordingly, the US Food and Drug Administration (FDA) includes information about PGx associations in many drug labels in a wide range of therapeutic areas.⁴ These PGx drug labels cover tests that are commonly used, but also include weaker genetic associations that are reported without requiring adjustments to pharmaceutical treatment. Most drugs with mandatory genetic testing are used in oncology, but PGx tests in other therapeutic areas are already being offered by laboratories and some have become part of standard clinical practice.^{5,6}

Personalizing drug treatments through PGx testing could improve their efficacy and safety, as well as reduce costs.⁷ However, as health-care resources are finite, it is important that the cost-effectiveness of novel PGx-guided treatment strategies is assessed in addition to their clinical utility before they are widely applied. Economic evaluations, which compare costs and outcomes of at least two competing interventions, are a useful tool to inform decision making and prioritize health-care spending. In the context of PGx testing, a pharmaco-economic study might contrast PGx-guided treatment with standard treatment (ST) with the same drug, or with an alternative drug that does not require genetic testing, or with both alternatives. When the PGx strategy is

found to be more effective at an acceptable additional cost (cost-effective) or more effective at a lower cost (cost-saving or dominant), this provides a strong argument for the implementation of PGx testing.

Previously published literature reviews of PGx-guided treatment and personalized medicine reported that the majority of PGx strategies were cost-effective or even dominant, though they noted that there was large heterogeneity in methodology between studies.^{8–12} Concerns over the quality of the early economic evaluations of PGx-guided treatment have been raised, but the quality is generally considered to have improved over time.^{13–16}

Our review of pharmaco-economic studies of PGx-guided treatment provides an update on the literature in this rapidly evolving field (the most recent previous review covered studies up to early 2013 (ref. 10)). Furthermore, we include a more extensive range of economic evaluations, whereas recent literature reviews were limited to cost utility analyses (CUAs) only.^{10,11} We also assessed the impact of freely available genetic information on the cost-effectiveness of PGx-guided treatment. We adopted a narrow definition of PGx, limiting our scope to consideration of variation in germline DNA. In contrast to tests on tumour, viral or bacterial DNA, germline DNA has the advantage that genetic variants need to be typed only once, and results remain relevant throughout a patient's life.

MATERIALS AND METHODS

Data sources and search strategy

The FDA Table of Pharmacogenomic Biomarkers in Drug Labeling lists FDA-approved drugs that include PGx information on their

¹MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK and ²Division of Medical and Molecular Genetics, Faculty of Life Sciences and Medicine, King's College London, London, UK. Correspondence: Professor CM Lewis, MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, Denmark Hill, London SE5 8AF, UK. E-mail: cathryn.lewis@kcl.ac.uk

Received 21 September 2016; revised 15 February 2017; accepted 14 April 2017




Figure 1. Cost-effectiveness plane of pharmaco-economic studies. PGx, pharmacogenetics-guided treatment; ST, standard treatment.

drug label along with the biomarker gene (accessed on 18 September 2015).⁴ We used this table to identify drugs for which there is a genetic variant associated with the drug efficacy, safety or dosing. We excluded non-germline genetic biomarkers, for example, mutations in viral or tumour DNA.

We then searched for the selected drugs and biomarkers in the National Health Service Economic Evaluations Database (NHS EED), a UK Department of Health and National Institute for Health Research-funded registry of economic evaluations of health and social care interventions.^{17,18} This resource includes CUAs, cost-effectiveness analyses (CEAs), cost-benefit analyses (CBAs—see below for definitions of these terms) and commentaries by the Centre for Reviews and Dissemination of the University of York. Funding of the NHS EED ceased in March 2015 and the latest database update was December 2014.

For each drug included in our study, the NHS EED was searched for economic evaluations that contain (1) the drug name and (2) the specific gene from the FDA label or the search terms genetic, genotype, genotypic, pharmacogenetic or pharmacogenomic in any field. We only included studies that compared a PGx-guided treatment strategy with at least one alternative strategy.

We also searched PubMed to identify more recent papers (until September 2015) and any other studies missed by the NHS EED search. We searched for articles that included (1) the name of the drug and (2) the specific gene mentioned in the FDA label or the search terms genetic, genotype, genotypic, pharmacogenetic or pharmacogenomic in the title or abstract, and (3) Cost-Benefit Analysis as a Medical Subject Headings term. In addition, the reference lists of retrieved publications were used to identify additional studies missed in our database searches.

Overview of economic evaluation methodology

Measuring and comparing costs and health outcomes is essential in a pharmaco-economic study. Whereas costs are naturally expressed in monetary units, the effect of a healthcare intervention can be expressed in different ways. In CUAs, health outcomes are assessed as quality-adjusted life years (QALYs), which measure the expected number of post-treatment years of life accounting for the quality of life. QALYs allow comparisons of treatment strategies across therapeutic areas and populations, but are an abstract concept ('quality' is hard to define) and their validity has been questioned.¹⁹ CEAs evaluate the effect of an intervention in terms of a disease or treatment specific measure, for example the

number of adverse events avoided, the change in score on a depression rating scale or time taken to remission. CBAs quantify treatment outcome in purely monetary terms.

Furthermore, the perspective of a pharmaco-economic study determines which costs and benefits are taken into account. These can be limited to costs to the public health-care system or private insurers, for example, staff salaries, drugs and equipment costs, or may include broader costs such as productivity losses and informal care. Commonly used perspectives are the third-party payer and societal perspective, but some studies take a hospital or patient perspective.

The incremental cost-effectiveness ratio (ICER) summarizes the difference in costs and health outcomes between a PGx-guided strategy and ST:

$$ICER = \frac{(Cost_{PGx} - Cost_{ST})}{(Effect_{PGx} - Effect_{ST})}$$

If the PGx treatment reduces costs and achieves a better outcome than the ST, then the PGx strategy dominates the ST. Contrarily, if the PGx option costs more but is less effective than the ST, then the PGx treatment is dominated by the ST (Figure 1). When one treatment comes at a higher cost but is also more effective than the other, the ICER is compared to a willingness-to-pay threshold to determine cost-effectiveness. Generally, ICERs up to £20 000–£30 000 per QALY (or \$30 000–\$50 000 per QALY) are considered cost-effective.²⁰ As costs, health outcomes and willingness-to-pay thresholds differ between countries, or may differ according to the assumptions and perspectives adopted, economic studies evaluating the same PGx test may come to different conclusions.

Analyses

We extracted key parameters from the reviewed economic evaluations, including the unit of outcome, country, perspective, ICER if applicable and the conclusion regarding the cost-effectiveness of the PGx testing strategy (the interpretation of the result as described in the publication). A parameter of particular interest is the cost of the genetic test, as this can significantly affect the cost-effectiveness of the PGx testing strategy and may change over time. To allow comparison between studies, the price of the genetic test was corrected for inflation and converted to US dollars estimated at 2014 levels (2014 US\$).

A stepwise linear regression model was fitted to test whether publication year, geographic region (Asia, Oceania, United States and Canada or EU) or perspective (health care, society or other) had an influence on the price of genetic testing. A stepwise logistic regression model was also used to investigate whether publication year, geographic region, perspective, cost of genetic test, genetic variant (*HLA*, *TPMT* or other) or outcome (QALY or other) was associated with the PGx testing strategy being cost-effective. Statistical analysis was performed in R (version 3.1.2, R Foundation for Statistical Computing, Vienna, Austria).

We estimated the impact of freely available genetic information on the conclusions regarding the cost-effectiveness of PGx-informed strategies. The ICER under assumption of free genetic testing was calculated by adjusting the cost of the PGx-guided treatment for the cost of the test as reported in the reviewed studies

$$ICER_{free\ PGx} = \frac{[Cost_{PGx} - Cost_{genetic\ test} - Cost_{ST}]}{(Effect_{PGx} - Effect_{ST})}$$

When insufficient details were provided to estimate the $ICER_{free\ PGx}$, it was assumed that free genetic testing could not worsen the conclusion regarding PGx-guided treatment. For example, when a study found the PGx strategy to be cost-effective, we assumed that PGx-guided treatment with free genetic testing would also be at least as cost-effective.

RESULTS

Description of studies

The FDA Table of Pharmacogenomic Biomarkers in Drug Labeling listed 137 distinct drugs, of which 68 met our inclusion criteria (Figure 2). These drugs were from diverse clinical specialties, including cancer (11 drugs), infectious diseases (10 drugs), psychiatry (9 drugs) and neurology (8 drugs) (Table 1). Our literature search yielded economic evaluations for only 10 of these 68 drugs (14.7%; Table 2). All publications related to a single drug, except for one study investigating a PGx testing strategy for




Figure 2. Number of drugs and publications included in literature review.

carbamazepine and phenytoin treatment, which assumed both drugs to be interchangeable in terms of costs, efficacy and safety.²¹ To avoid duplication of studies in our review, this publication was counted as a carbamazepine study (there were no other publications on phenytoin).

We retrieved 44 economic evaluations that investigated the cost-effectiveness of a PGx-informed strategy (Table 2). Full details of the reviewed studies and extracted information are given in Supplementary Table 1. The earliest study included was published in 2000 and over 70% of studies were published in 2009 or later. Most publications were CUAs (30 studies, 68%) or CEAs (12 studies, 27%), with only two CBAs (5%). A health-care system perspective was adopted in 18 studies (41%), a societal perspective in 10 papers (23%), a third-party payer perspective in 5 studies (11%) and 11 papers (25%) did not state a clear perspective. In all, 20 studies (45%) were conducted in North America, 11 (25%) in Europe, 6 (14%) in Asia and 3 (7%) in Oceania; 4 studies (9%) did not specify a country. Warfarin had the most economic evaluations (12 studies), followed by azathioprine (9 studies); clozapine and mercaptopurine had only 1 economic evaluation each (Table 2).

Cost-effectiveness of PGx-informed treatment

We assessed the overall conclusions regarding cost-effectiveness of each PGx study. Over half of the 44 economic evaluations took a favourable view of the PGx-guided strategy: in 12 studies (27%) it was dominant (cost-saving) and in 13 studies (30%) it was cost-effective. Eleven publications (25%) found PGx testing not cost-effective and 8 studies (18%) did not reach a definitive conclusion (Figure 3a). The majority of economic evaluations concluded in favour of PGx testing for azathioprine (7 out of 9 studies), clopidogrel (4 out of 6 studies), abacavir (4 out of 5 studies), carbamazepine (3 out of 4 studies), irinotecan (3 out of 3 studies) and clozapine (1 study) (Figure 3b). Although warfarin had the highest number of economic studies, they reached diverging conclusions: 3 studies found PGx-guided dosing cost-effective, 4 studies were inconclusive and 5 studies concluded it was not cost-effective. No studies found unequivocally that PGx-guided

Table 1. Drugs from the FDA Table of Pharmacogenomic Biomarkers in Drug Labeling included in literature review

Therapeutic area	Count	Drugs
Oncology	11	Capecitabine, cisplatin, dabrafenib, fluorouracil, irinotecan , lapatinib, mercaptopurine , nilotinib, pazopanib, rasburicase, thioguanine
Infectious diseases	10	Abacavir , chloroquine, dapsone, mafenide, nalidixic acid, nitrofurantoin, primaquine, quinine sulphate, rifampin+isoniazid+pyrazinamide ^a , sulfamethoxazole+trimethoprim ^a
Psychiatry	9	Aripiprazole, atomoxetine, citalopram , clozapine , fluvoxamine, iloperidone, perphenazine, pimozide, thioridazine
Neurology	8	Carbamazepine , clobazam, dextromethorphan+quinidine ^a , divalproex, phenytoin , tetrabenazine, valproic acid, vortioxetine
Cardiology	5	Carvedilol, clopidogrel , isosorbide+hydralazine ^a , metoprolol, propafenone
Gastroenterology	5	Dexlansoprazole, esomeprazole, metoclopramide, PEG-3350+sodium sulphate+sodium chloride+potassium chloride+sodium ascorbate+ascorbic acid ^a , rabeprazole
Rheumatology	5	Azathioprine , carisoprodol, celecoxib, flurbiprofen, pegloticase
Endocrinology	4	Chlorpropamide, glimepiride, glipizide, glyburide
Haematology	3	Eltrombopag, methylene blue, warfarin
Analgesic	1	Tramadol
Anaesthesiology	1	Codeine
Dental	1	Cevimeline
Genitourinary	1	Tolterodine
Inborn errors of metabolism	1	Eliglustat
Pulmonary	1	Ivacaftor
Toxicology	1	Sodium nitrite
Transplantation	1	Mycophenolic acid

Abbreviation: FDA, Food and Drug Administration. Drugs in bold had economic evaluations available. ^aMultiple drugs on a single FDA label.

Table 2. Drugs for which a PGx-guided strategy was studied in economic evaluation(s)

Drug	Therapeutic area	Gene	Notes (based on PharmGKB.org ³⁷)	Number of reviewed publications
Abacavir	HIV	HLA-B	Abacavir is contraindicated for HLA-B*5701 carriers as they are at high risk of hypersensitivity reaction.	5 (refs 38–42)
Azathioprine	Rheumatology	TPMT	Carriers of one nonfunctional TPMT allele may require reduced azathioprine dose. Carriers of two nonfunctional TPMT alleles are at high risk of myelotoxicity and alternative treatment should be considered.	9 (refs 43–51)
Carbamazepine	Neurology	HLA-B, HLA-A	Carbamazepine is contraindicated for HLA-B*1502 carriers as they are at high risk Stevens–Johnson syndrome/toxic epidermal necrolysis. HLA-A*3101 has also been associated with hypersensitivity reactions.	4 (refs 21,52–54)
Citalopram	Psychiatry	CYP2C19, 5-HTTLPR ^a , HTR2A ^a	CYP2C19 poor metabolizers require reduced citalopram starting dose. Polymorphisms in 5-HTTLPR and HTR2A are associated with citalopram response. ^{25,56}	3 (refs 57–59)
Clopidogrel	Cardiology	CYP2C19	CYP2C19 poor metabolizers have reduced response to clopidogrel and alternative treatment should be considered.	6 (refs 25,26,60–63)
Clozapine	Psychiatry	CYP2D6, H2 ^a , 5-HTT ^a , 5-HT2A ^a	CYP2D6 poor metabolizers may require reduced clozapine dose. Six polymorphisms in H2, 5-HTT, 5-HT2A and 5-HT2C are associated with clozapine response. ⁶⁴	1 (ref. 65)
Irinotecan	Oncology	UGT1A1	Patients homozygous for the UGT1A1*28 allele are at higher risk of neutropenia and should receive a reduced starting dose of irinotecan.	3 (refs 66–68)
Mercaptopurine	Oncology	TPMT	Carriers of one nonfunctional TPMT allele may require reduced mercaptopurine dose. Carriers of two nonfunctional TPMT alleles are at high risk of myelotoxicity and alternative treatment should be considered.	1 (ref. 69)
Warfarin	Cardiology	CYP2C9, VKORC1	Genetic variation in VKORC1 and CYP2C9 explain 40% variance in warfarin dose. Genetic and clinical information can be used to determine starting dose.	12 (refs 22–24,30,70–77)

Abbreviation: PGx, pharmacogenetics. ^aGene not mentioned on FDA drug label but appears in economic evaluations.

citalopram (3 studies) or mercaptopurine (1 study) treatment was cost-effective.

We assessed the effect of study characteristics on the probability of concluding in favour of the PGx strategy. A logistic regression model detected that CUAs (studies using QALYs as outcome measure) were less likely than CEAs and CBAs to find the genetic testing strategy cost-effective (odds ratio=0.13, *P*-value < 0.05). However, there is no clear explanation for this and it may be a spurious result due to the relatively small sample size of 44 economic evaluations.

Effect of cost of genetic test on cost-effectiveness of PGx-informed treatment

The cost of genetic testing is an important parameter of economic evaluations of PGx interventions. After correcting for inflation and converting to 2014 US\$, the cost of genetic testing quoted by the reviewed studies ranged between US\$33 and US\$710 with a median value of US\$175. The price of genetic tests decreased slightly over time (not statistically significant) and this trend was more pronounced since 2009, the period when most economic evaluations were published (*P*-value < 0.05; Figure 4). Prices were on average higher in the United States and Canada than other regions of the world (mean United States and Canada: US\$363.65; mean other regions: US\$131.80; *P*-value < 0.05). We noted a wide variability in prices of tests for the same drug. For example, the lowest price quoted for warfarin PGx testing was US\$36 in a 2014 UK-based study,²² while US\$600 and US\$657 were used in a 2013 Canadian and 2009 US study, respectively.^{23,24} The prices for clopidogrel PGx testing also varied considerably: from US\$45 (2013 Australian study) to US\$ 543 (2013 US study).^{25,26}

Given the decreasing costs of genetic testing and its increasing availability, we looked ahead to a possible future where genotype information might be readily available, at negligible cost, for all patients as part of their electronic health record. Thirty-three economic evaluations (75%) would support PGx-guided treatment under this scenario, with 11 studies (25%) finding it cost-effective and 22 studies (50%) considering it dominant and cost-saving (Figure 3c). Five studies (11%) would still conclude that PGx testing was not cost-effective, while 3 studies (7%) would be inconclusive. A separate set of 3 studies had to be excluded, because the impact of free genetic testing could not be estimated. We note that the effect of freely available genetic information can be striking for some drugs. None of the published studies for citalopram and mercaptopurine found PGx-informed treatment to be cost-effective, but all studies switched in favour of PGx testing under the negligible test cost scenario (Figure 3d). For the 12 economic evaluations of warfarin, the number of cost-effective studies would increase from 3 to 7 with freely available genetic testing.

DISCUSSION

We have assessed published economic evaluations comparing the cost-effectiveness of PGx-guided treatment to ST for drugs listed in the FDA Table of Pharmacogenomic Biomarkers in Drug Labeling. The economic evaluations were drawn from the NHS EED database, which includes economic evaluations up to 31 December 2014. An alternative source of economic studies would be the Cost-Effectiveness Registry (CEA Registry) maintained by the Tufts Medical Centre. We opted to use the more comprehensive NHS EED as the CEA Registry is limited to CUAs (measuring health outcomes in QALYs), which would have reduced the number of evaluations available for assessment. Moreover, the CEA Registry was not updated beyond 2014 and it only provides advanced database searches for subscribers and contributors.¹⁸ A third resource, the Health Economic Evaluations Database curated by John Wiley & Sons, was discontinued in 2014.²⁷ As economic




Figure 3. Conclusions of reviewed economic evaluations regarding cost-effectiveness of PGx testing strategy (a) overall and (b) by drug, and estimated conclusions in scenario of no extra cost for genetic information (c) overall and (d) by drug. PGx, pharmacogenetics-guided treatment.

evaluations provide evidence for the introduction of PGx testing into clinical practice, we argue that an up-to-date, accessible database would be an important and valuable resource for both health-economic and PGx research.

Few of the FDA-listed drugs have been the subject of published economic evaluations assessing the economic aspects of PGx testing. This was previously also noted by Phillips *et al.*,¹¹ who found that only 13% of drugs on the FDA table and only 27% of available genetic tests were accompanied by economic studies. However, it is increasingly the case that clinical utility alone is not sufficient to recommend application of a PGx test in clinical practice, and a favourable economic assessment is therefore of increasing importance. We call for more pharmaco-economic studies in this field, which should be regularly updated to respond to the changing landscape of health-care and, in particular, genetic testing costs.

There are various limitations of our study that need to be taken into account. One is that the economic evaluations reviewed may not be representative of all PGx tests. For example, the economic aspects of PGx-guided treatment are of less relevance in cases where testing is clearly necessary, for example, because it

prevents life-threatening adverse events, and economic studies in such cases therefore tend to be lacking. Another possibility is that economic studies focus on PGx tests that are already applied in clinical practice and for which there is an apparent interest. Studies that find genetic testing to be not cost-effective may also be less likely to be published.

Notwithstanding the above issues, economic evaluations also have certain intrinsic limitations. One is that certain inputs into the model are difficult to quantify accurately. For example, parameters such as the response rate, the probability of adverse drug reactions and the cost of managing adverse drug reactions must sometimes be estimated from sparse information. Randomized clinical trials are the preferred source for these input data, but these are not always available. Ideally, the uncertainty in the input estimates should be accounted for in the economic modelling, and sensitivity analyses should be performed to verify how robust the result is to deviations in the inputs, but the level of uncertainty to apply can itself be a matter of subjective opinion, and vary from study to study.

Another intrinsic issue is that context and perspective may influence the conclusion of a study. For example, comparing

PGx-informed treatment. Increased costs may arise where the alternative drug for test-positive patients is more expensive, and this is exacerbated whether the test has a high proportion of false-positive results. For example, patients with heart disease or stroke who are *CYP2C19* poor metabolizers may be prescribed the more expensive ticagrelor in place of clopidogrel (which is metabolized into its active form by *CYP2C19*).²⁵ Thus, even if genetic information is freely accessible, economic evaluations of PGx testing are still relevant and necessary.

The economic evaluation studies reviewed here show that PGx has a positive impact on health-care quality and costs. Over half of reviewed studies concluded that the PGx-informed treatment strategy is more cost-effective than the alternatives considered under present-day economics. Only one in four economic evaluations found the genetic testing option unequivocally not cost-effective. This encouraging finding, with an even bigger projected benefit under low-cost genetic typing, suggests that PGx testing has the potential to be a cost-effective or even cost-saving intervention. It therefore seems likely that PGx testing will become a core clinical service, particularly as projects such as the 100 000 Genomes Project pushes genomics to become part of health-care infrastructure and as electronic health records become increasingly effective.³⁶

CONFLICT OF INTEREST

MV is funded by a studentship from the Medical Research Council and Eli Lilly and Company Ltd. MEW is a part-time employee of Genomics plc. CML declares no conflict of interest.

ACKNOWLEDGMENTS

This study was funded by an industrial CASE studentship to MV from the Medical Research Council with Eli Lilly and Company Ltd. This paper represents independent research part-funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

REFERENCES

- Mallal S, Phillips E, Carosi G, Molina J-M, Workman C, Tomažič J *et al*. HLA-B* 5701 screening for hypersensitivity to abacavir. *N Engl J Med* 2008; **358**: 568–579.
- Chung W-H, Hung S-I, Hong H-S, Hsieh M-S, Yang L-C, Ho H-C *et al*. Medical genetics: a marker for Stevens–Johnson syndrome. *Nature* 2004; **428**: 486–486.
- Wadelius M, Chen LY, Eriksson N, Bumpstead S, Ghori J, Wadelius C *et al*. Association of warfarin dose with genes involved in its action and metabolism. *Hum Genet* 2007; **121**: 23–34.
- U.S. Food and Drug Administration. Table of Pharmacogenomic Biomarkers in Drug Labeling. Available at <http://www.fda.gov/Drugs/ScienceResearch/ResearchAreas/Pharmacogenetics/ucm083378.htm> (accessed on 18 September 2015).
- Centers for Disease Control and Prevention. Genomic Tests and Family Health History by Levels of Evidence: Public Health Genomics. Available at <http://www.cdc.gov/genomics/gtesting/tier.htm> (accessed on 17 February 2016).
- Rubinstein WS, Maglott DR, Lee JM, Kattman BL, Malheiro AJ, Ovetsky M *et al*. The NIH genetic testing registry: a new, centralized database of genetic tests to enable access to comprehensive information and improve transparency. *Nucleic Acids Res* 2013; **41**: 925–935.
- Haycox A, Pirmohamed M, McLeod C, Houten R, Richards S. Through a glass darkly: economics and personalised medicine. *Pharmacoeconomics* 2014; **32**: 1055–1061.
- Phillips KA, Van Bebber SL. A systematic review of cost-effectiveness analyses of pharmacogenomic interventions. *Pharmacogenomics* 2004; **5**: 1139–1149.
- Vegter S, Boersma C, Rozenbaum M, Wilffert B, Navis G, Postma MJ. Pharmacoeconomic evaluations of pharmacogenetic and genomic screening programmes: a systematic review on content and adherence to guidelines. *Pharmacoeconomics* 2008; **26**: 569–587.
- Hatz MHM, Schremser K, Rogowski WH. Is individualized medicine more cost-effective? A systematic review. *Pharmacoeconomics* 2014; **32**: 443–455.
- Phillips KA, Ann Sakowski J, Trosman J, Douglas MP, Liang S-Y, Neumann P. The economic value of personalized medicine tests: what we know and what we need to know. *Genet Med* 2014; **16**: 251–257.
- Beaulieu M, de Denus S, Lachaine J. Systematic review of pharmacoeconomic studies of pharmacogenomic tests. *Pharmacogenomics* 2010; **11**: 1573–1590.
- Payne K, Shabaruddin FH. Cost-effectiveness analysis in pharmacogenomics. *Pharmacogenomics* 2010; **11**: 643–646.
- Shabaruddin FH, Fleeman ND, Payne K. Economic evaluations of personalized medicine: existing challenges and current developments. *Pharmacoecon Personalized Med* 2015; **8**: 115–126.
- Vegter S, Jansen E, Postma MJ, Boersma C. Economic evaluations of pharmacogenetic and genomic screening programs: update of the literature. *Drug Dev Res* 2010; **71**: 492–501.
- Wong WB, Carlson JJ, Thariani R, Veenstra DL. Cost effectiveness of pharmacogenomics: a critical and systematic review. *Pharmacoeconomics* 2010; **28**: 1001–1013.
- Nixon J, Stoykova B, Christie J, Glanville J, Kleijnen J, Drummond M. NHS Economic Evaluation Database for healthcare decision makers. *BMJ* 2000; **321**: 32.
- University of York Centre for Reviews and Dissemination. Welcome to the CRD Database. Available at <http://www.crd.york.ac.uk/CRDWeb/HomePage.asp> (accessed on 30 October 2015).
- Beresniak A, Medina-Lara A, Auray JP, De Wever A, Praet JC, Tarricone R *et al*. Validation of the underlying assumptions of the quality-adjusted life-years outcome: results from the ECHOUTCOME European project. *Pharmacoeconomics* 2015; **33**: 61–69.
- McCabe C, Claxton K, Culyer A. The NICE Cost-Effectiveness Threshold. *Pharmacoeconomics* 2008; **26**: 733–744.
- Dong D, Sung C, Finkelstein EA. Cost-effectiveness of HLA-B*1502 genotyping in adult patients with newly diagnosed epilepsy in Singapore. *Neurology* 2012; **79**: 1259–1267.
- Pink J, Pirmohamed M, Lane S, Hughes DA. Cost-effectiveness of pharmacogenetics-guided warfarin therapy vs. alternative anticoagulation in atrial fibrillation. *Clin Pharmacol Ther* 2014; **95**: 199–207.
- Nshimyumukiza L, Duplantie J, Gagnon M, Douville X, Fournier D, Lindsay C *et al*. Dabigatran versus warfarin under standard or pharmacogenetic-guided management for the prevention of stroke and systemic thromboembolism in patients with atrial fibrillation: a cost/utility analysis using an analytic decision model. *Thromb J* 2013; **11**: 14.
- Patrick AR, Avorn J, Choudhry NK. Cost-effectiveness of genotype-guided warfarin dosing for patients with atrial fibrillation. *Circ Cardiovasc Qual Outcomes* 2009; **2**: 429–436.
- Sorich MJ, Horowitz JD, Sorich W, Wiese MD, Pekarsky B, Karnon JD. Cost-effectiveness of using *CYP2C19* genotype to guide selection of clopidogrel or ticagrelor in Australia. *Pharmacogenomics* 2013; **14**: 2013–2021.
- Lala A, Berger JS, Sharma G, Hochman JS, Scott Braithwaite R, Ladapo JA. Genetic testing in patients with acute coronary syndrome undergoing percutaneous coronary intervention: a cost-effectiveness analysis. *J Thromb Haemost* 2013; **11**: 81–91.
- Wiley Online Library, John Wiley & Sons Ltd. HEED: Health Economic Evaluations Database. Available at: <http://onlinelibrary.wiley.com/book/10.1002/9780470510933> (accessed on 28 January 2016).
- Altar CA, Carhart J, Allen JD, Hall-Flavin D, Winner J, Dechairo B. Clinical utility of combinatorial pharmacogenomics-guided antidepressant therapy: evidence from three clinical studies. *Mol Neuropsychiatry* 2015; **1**: 145–155.
- Hornberger J, Li Q, Quinn B. Cost-effectiveness of combinatorial pharmacogenomic testing for treatment-resistant major depressive disorder patients. *Am J Manag Care* 2014; **21**: e357–e365.
- You JH. Pharmacogenetic-guided selection of warfarin versus novel oral anticoagulants for stroke prevention in patients with atrial fibrillation: a cost-effectiveness analysis. *Pharmacogenet Genomics* 2014; **24**: 6–14.
- Stergiopoulos K, Brown DL. Genotype-guided vs clinical dosing of warfarin and its analogues: meta-analysis of randomized clinical trials. *JAMA Intern Med* 2014; **174**: 1330–1338.
- U.S. Food and Drug Administration. Ziagen Prescribing Information. Available at http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/020977s030,020978s034lbl.pdf (accessed on 2 February 2016).
- 23andMe Inc. 23andMe: Welcome. Available at <https://www.23andme.com/en-gb> (accessed on 1 February 2016).
- Lu M, Lewis CM, Traylor M. Pharmacogenetic testing through the direct-to-consumer genetic testing company 23andMe. *bioRxiv* 2017; <https://doi.org/10.1101/098541> (accessed 6 Feb 2017).
- Wetterstrand KA. DNA Sequencing Costs: Data from the NHGRI Genome Sequencing Program (GSP). Available at www.genome.gov/sequencingcostsdata (accessed on 27 January 2016).
- Genomics England: Genomics England is delivering the 100,000 Genomes Project. Available at <http://www.genomicsengland.co.uk> (accessed on 31 March 2016).
- Whirl-Carrillo M, McDonagh EM, Hebert JM, Gong L, Sangkuhl K, Thorn CF *et al*. Pharmacogenomics knowledge for personalized medicine. *Clin Pharmacol Ther* 2012; **92**: 414–417.

- 38 Hughes DA, Vilar FJ, Ward CC, Alfrevic A, Park BK, Pirmohamed M. Cost-effectiveness analysis of HLA B*5701 genotyping in preventing abacavir hypersensitivity. *Pharmacogenetics* 2004; **14**: 335–342.
- 39 Schackman BR, Scott CA, Walensky RP, Losina E, Freedberg KA, Sax PE. The cost-effectiveness of HLA-B*5701 genetic screening to guide initial antiretroviral therapy for HIV. *AIDS* 2008; **22**: 2025–2033.
- 40 Kauf TL, Farkouh RA, Earnshaw SR, Watson ME, Maroudas P, Chambers MG. Economic efficiency of genetic screening to inform the use of abacavir sulfate in the treatment of HIV. *Pharmacoeconomics* 2010; **28**: 1025–1039.
- 41 Nieves Calatrava D, Calle-Martin Ode L, Iribarren-Loyarte JA, Rivero-Roman A, Garcia-Bujalance L, Perez-Escolano I et al. Cost-effectiveness analysis of HLA-B*5701 typing in the prevention of hypersensitivity to abacavir in HIV+ patients in Spain. *Enferm Infecc Microbiol Clin* 2010; **28**: 590–595.
- 42 Kapoor R, Martinez-Vega R, Dong D, Tan SY, Leo YS, Lee CC et al. Reducing hypersensitivity reactions with HLA-B*5701 genotyping before abacavir prescription: clinically useful but is it cost-effective in Singapore? *Pharmacogenomics* 2015; **25**: 60–72.
- 43 Tavadia SM, Mydlarski PR, Reis MD, Mittmann N, Pinkerton PH, Shear N et al. Screening for azathioprine toxicity: a pharmacoeconomic analysis based on a target case. *J Am Acad Dermatol* 2000; **42**: 628–632.
- 44 Marra CA, Esdaile JM, Anis AH. Practical pharmacogenetics: the cost effectiveness of screening for thiopurine s-methyltransferase polymorphisms in patients with rheumatological conditions treated with azathioprine. *J Rheumatol* 2002; **29**: 2507–2512.
- 45 Oh KT, Anis AH, Bae SC. Pharmacoeconomic analysis of thiopurine methyltransferase polymorphism screening by polymerase chain reaction for treatment with azathioprine in Korea. *Rheumatology (Oxford)* 2004; **43**: 156–163.
- 46 Winter J, Walker A, Shapiro D, Gaffney D, Spooner RJ, Mills PR. Cost-effectiveness of thiopurine methyltransferase genotype screening in patients about to commence azathioprine therapy for treatment of inflammatory bowel disease. *Aliment Pharmacol Ther* 2004; **20**: 593–599.
- 47 Dubinsky MC, Reyes E, Ofman J, Chiou C-F, Wade S, Sandborn WJ. A cost-effectiveness analysis of alternative disease management strategies in patients with Crohn's disease treated with azathioprine or 6-mercaptopurine. *Am J Gastroenterol* 2005; **100**: 2239–2247.
- 48 Priest VL, Begg EJ, Gardiner SJ, Frampton CM, Geary RB, Barclay ML et al. Pharmacoeconomic analyses of azathioprine, methotrexate and prospective pharmacogenetic testing for the management of inflammatory bowel disease. *Pharmacoeconomics* 2006; **24**: 767–781.
- 49 Van Den Akker-Van Marle ME, Gurwitz D, Detmar SB, Enzing CM, Hopkins MM, Gutierrez De Mesa E et al. Cost-effectiveness of pharmacogenomics in clinical practice: a case study of thiopurine methyltransferase genotyping in acute lymphoblastic leukemia in Europe. *Pharmacogenomics* 2006; **7**: 783–792.
- 50 Hagaman JT, Kinder BW, Eckman MH. Thiopurine S- methyltransferase [corrected] testing in idiopathic pulmonary fibrosis: a pharmacogenetic cost-effectiveness analysis. *Lung* 2010; **188**: 125–132.
- 51 Thompson AJ, Newman WG, Elliott RA, Roberts SA, Tricker K, Payne K. The cost-effectiveness of a pharmacogenetic test: a trial-based evaluation of TPMT genotyping for azathioprine. *Value Health* 2014; **17**: 22–33.
- 52 Rattanavipapong W, Koopitakkajorn T, Praditsitthikorn N, Mahasirimongkol S, Teerawattananon Y. Economic evaluation of HLA-B*15:02 screening for carbamazepine-induced severe adverse drug reactions in Thailand. *Epilepsia* 2013; **54**: 1628–1638.
- 53 Tiamkao S, Jitpimolmard J, Sawanyawisuth K, Jitpimolmard S. Cost minimization of HLA-B*1502 screening before prescribing carbamazepine in Thailand. *Int J Clin Pharm* 2013; **35**: 608–612.
- 54 Plumpton CO, Yip VL, Alfrevic A, Marson AG, Pirmohamed M, Hughes DA. Cost-effectiveness of screening for HLA-A*31:01 prior to initiation of carbamazepine in epilepsy. *Epilepsia* 2015; **56**: 556–563.
- 55 McMahon FJ, Buervenich S, Charney D, Lipsky R, Rush AJ, Wilson AF et al. Variation in the gene encoding the serotonin 2A receptor is associated with outcome of antidepressant treatment. *Am J Hum Genet* 2006; **78**: 804–814.
- 56 Serretti A, Kato M, De Ronchi D, Kinoshita T. Meta-analysis of serotonin transporter gene promoter polymorphism (5-HTTLPR) association with selective serotonin reuptake inhibitor efficacy in depressed patients. *Mol Psychiatry* 2007; **12**: 247–257.
- 57 Perlis RH, Patrick A, Smoller JW, Wang PS. When is Pharmacogenetic Testing for Antidepressant Response Ready for the Clinic[quest] A Cost-effectiveness Analysis Based on Data from the STAR[ast]D Study. *Neuropsychopharmacology* 2009; **34**: 2227–2236.
- 58 Serretti A, Olgiati P, Bajo E, Bigelli M, De Ronchi D. A model to incorporate genetic testing (5-HTTLPR) in pharmacological treatment of major depressive disorders. *World J Biol Psychiatry* 2011; **12**: 501–515.
- 59 Olgiati P, Bajo E, Bigelli M, De Ronchi D, Serretti A. Should pharmacogenetics be incorporated in major depression treatment? Economic evaluation in high- and middle-income European countries. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; **36**: 147–154.
- 60 Crespin DJ, Federspiel JJ, Biddle AK, Jonas DE, Rossi JS. Ticagrelor versus genotype-driven antiplatelet therapy for secondary prevention after acute coronary syndrome: a cost-effectiveness analysis. *Value Health* 2011; **14**: 483–491.
- 61 Panattoni L, Brown PM, Te Ao B, Webster M, Gladding P. The cost effectiveness of genetic testing for CYP2C19 variants to guide thienopyridine treatment in patients with acute coronary syndromes: a New Zealand evaluation. *Pharmacoeconomics* 2012; **30**: 1067–1084.
- 62 Reese ES, Daniel Mullins C, Beitelshes AL, Onukwugha E. Cost-effectiveness of cytochrome P450 2C19 genotype screening for selection of antiplatelet therapy with clopidogrel or prasugrel. *Pharmacotherapy* 2012; **32**: 323–332.
- 63 Kazi DS, Garber AM, Shah RU, Dudley RA, Mell MW, Rhee C et al. Cost-effectiveness of genotype-guided and dual antiplatelet therapies in acute coronary syndrome. *Ann Intern Med* 2014; **160**: 221–232.
- 64 Arranz MJ, Munro J, Birkett J, Bolonna A, Mancama D, Sodhi M et al. Pharmacogenetic prediction of clozapine response. *Lancet* 2000; **355**: 1615–1616.
- 65 Perlis RH, Ganz DA, Avorn J, Schneeweiss S, Glynn RJ, Smoller JW et al. Pharmacogenetic testing in the clinical management of schizophrenia: a decision-analytic model. *J Clin Psychopharmacol* 2005; **25**: 427–434.
- 66 Obradovic M, Mrhar A, Kos M. Cost-effectiveness of UGT1A1 genotyping in second-line, high-dose, once every 3 weeks irinotecan monotherapy treatment of colorectal cancer. *Pharmacogenomics* 2008; **9**: 539–549.
- 67 Gold HT, Hall MJ, Blinder V, Schackman BR. Cost effectiveness of pharmacogenetic testing for uridine diphosphate glucuronosyltransferase 1A1 before irinotecan administration for metastatic colorectal cancer. *Cancer* 2009; **115**: 3858–3867.
- 68 Pichereau S, Le Louarn A, Lecomte T, Blasco H, Le Guellec C, Bourgoin H. Cost-effectiveness of UGT1A1*28 genotyping in preventing severe neutropenia following FOLFIRI therapy in colorectal cancer. *J Pharm Pharm Sci* 2010; **13**: 615–625.
- 69 Donnan JR, Ungar WJ, Mathews M, Hancock-Howard RL, Rahman P. A cost effectiveness analysis of thiopurine methyltransferase testing for guiding 6-mercaptopurine dosing in children with acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2011; **57**: 231–239.
- 70 You JH, Chan FW, Wong RS, Cheng G. The potential clinical and economic outcomes of pharmacogenetics-oriented management of warfarin therapy—a decision analysis. *Thromb Haemost* 2004; **92**: 590–597.
- 71 Eckman MH, Rosand J, Greenberg SM, Gage BF. Cost-effectiveness of using pharmacogenetic information in warfarin dosing for patients with nonvalvular atrial fibrillation. *Ann Intern Med* 2009; **150**: 73–83.
- 72 Leey JA, McCabe S, Koch JA, Miles TP. Cost-effectiveness of genotype-guided warfarin therapy for anticoagulation in elderly patients with atrial fibrillation. *Am J Geriatr Pharmacother* 2009; **7**: 197–203.
- 73 You JH, Tsui KK, Wong RS, Cheng G. Potential clinical and economic outcomes of CYP2C9 and VKORC1 genotype-guided dosing in patients starting warfarin therapy. *Clin Pharmacol Ther* 2009; **86**: 540–547.
- 74 Meckley L, Gudgeon J, Anderson J, Williams M, Veenstra D. A policy model to evaluate the benefits, risks and costs of warfarin pharmacogenomic testing. *Pharmacoeconomics* 2010; **28**: 61–74.
- 75 You JHS, Tsui KKN, Wong RSM, Cheng G. Cost-effectiveness of dabigatran versus genotype-guided management of warfarin therapy for stroke prevention in patients with atrial fibrillation. *PLoS ONE* 2012; **7**: e39640.
- 76 Chong HY, Saokaew S, Dumrongpratt K, Permsuwan U, Wu DB, Sritara P et al. Cost-effectiveness analysis of pharmacogenetic-guided warfarin dosing in Thailand. *Thromb Res* 2014; **134**: 1278–1284.
- 77 Mitropoulou C, Fragoulakis V, Bozina N, Vozikis A, Supe S, Bozina T et al. Economic evaluation of pharmacogenomic-guided warfarin treatment for elderly Croatian atrial fibrillation patients with ischemic stroke. *Pharmacogenomics* 2015; **16**: 137–148.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017

Supplementary Information accompanies the paper on the The Pharmacogenomics Journal website (<http://www.nature.com/tpj>)