



King's Research Portal

DOI:

[10.1016/j.ijid.2017.04.020](https://doi.org/10.1016/j.ijid.2017.04.020)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Charani, E., Ahmad, R., Tarrant, C., Birgand, G., Leather, A., Mendelson, M., ... Holmes, A. (2017). Opportunities for system level improvement in antibiotic use across the surgical pathway. *INTERNATIONAL JOURNAL OF INFECTIOUS DISEASES*, 60, 29-34. <https://doi.org/10.1016/j.ijid.2017.04.020>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: www.elsevier.com/locate/ijid

Perspective

Opportunities for system level improvement in antibiotic use across the surgical pathway



E. Charani^{a,*}, R. Ahmad^a, C. Tarrant^b, G. Birgand^a, A. Leather^c, M. Mendelson^d,
S.R. Moonesinghe^e, N. Sevdalis^f, S. Singh^g, A. Holmes^a

^a NIHR Health Protection Research Unit in Antimicrobial Resistance and Healthcare Associated Infection, Imperial College London, Department of Medicine, London, UK

^b Department of Health Sciences, University of Leicester, Centre for Medicine, Leicester, UK

^c King's Centre for Global Health & Health Partnerships, Division of Health & Social Care Research, Faculty of Life Sciences & Medicine, King's College London, UK

^d Division of Infectious Diseases and HIV Medicine, Department of Medicine, University of Cape Town, Groote Schuur Hospital Observatory, Cape Town, South Africa

^e Centre for Anaesthesia Critical Care and Pain Medicine, University College London Hospitals, London, UK

^f Centre for Implementation Science, Institute of Psychiatry, King's College London, Denmark Hill, UK

^g School of Medicine, Amrita University, Tamilnadu, Kochi, India

ARTICLE INFO

Article history:

Received 24 January 2017

Received in revised form 19 April 2017

Accepted 25 April 2017

Corresponding Editor: Eskild Petersen, Aarhus, Denmark

Keywords:

Surgery

Infection

Antibiotic prescribing

Low and middle income countries

SUMMARY

Optimizing antibiotic prescribing across the surgical pathway (before, during, and after surgery) is a key aspect of tackling important drivers of antimicrobial resistance and simultaneously decreasing the burden of infection at the global level. In the UK alone, 10 million patients undergo surgery every year, which is equivalent to 60% of the annual hospital admissions having a surgical intervention. The overwhelming majority of surgical procedures require effectively limited delivery of antibiotic prophylaxis to prevent infections. Evidence from around the world indicates that antibiotics for surgical prophylaxis are administered ineffectively, or are extended for an inappropriate duration of time postoperatively. Ineffective antibiotic prophylaxis can contribute to the development of surgical site infections (SSIs), which represent a significant global burden of disease. The World Health Organization estimates SSI rates of up to 50% in postoperative surgical patients (depending on the type of surgery), with a particular problem in low- and middle-income countries, where SSIs are the most frequently reported healthcare-associated infections. Across European hospitals, SSIs alone comprise 19.6% of all healthcare-acquired infections. Much of the scientific research in infection management in surgery is related to infection prevention and control in the operating room, surgical prophylaxis, and the management of SSIs, with many studies focusing on infection within the 30-day postoperative period. However it is important to note that SSIs represent only one of the many types of infection that can occur postoperatively. This article provides an overview of the surgical pathway and considers infection management and antibiotic prescribing at each step of the pathway. The aim was to identify the implications for research and opportunities for system improvement.

© 2017 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Optimizing antibiotic prescribing across the surgical pathway (before, during, and after surgery) is a key aspect of tackling important drivers of antimicrobial resistance (AMR) and simultaneously decreasing the burden of infection at the global level. In the UK alone, 10 million patients undergo surgery every year (*The Royal College of Anaesthetists, 2015*), which is equivalent to 60% of the annual hospital admissions having a surgical intervention

* Corresponding author at: NIHR Health Protection Research Unit in Antimicrobial Resistance and Healthcare Associated Infection, Imperial College London, Department of Medicine, 8th Floor Commonwealth Building, Du Cane Road, London W12 0HS, UK. Tel: 02033132732.

E-mail address: e.charani@imperial.ac.uk (E. Charani).

(Anon, 2016). The overwhelming majority of surgical procedures require effectively limited delivery of antibiotic prophylaxis to prevent infections (Boucher et al., 2009; Bratzler et al., 2013). Evidence from around the world indicates that antibiotics for surgical prophylaxis are administered ineffectively, or are extended for an inappropriate duration of time postoperatively (Leeds et al., 2016; Tan et al., 2006). Ineffective antibiotic prophylaxis can contribute to the development of surgical site infections (SSIs), which represent a significant global burden of disease. The World Health Organization (WHO) estimates SSI rates of up to 50% in postoperative surgical patients (depending on the type of surgery), with a particular problem in low- and middle-income countries (LMICs), where SSIs are the most frequently reported healthcare-associated infections (HAIs) (Aveling et al., 2013; Aiken et al., 2013; World Health Organization, 2016). Across European hospitals, SSIs alone comprise 19.6% of all healthcare-acquired infections (Anon, 2013). Much of the scientific research in infection management in surgery is related to infection prevention and control in the operating room (Birgand et al., 2014; Cosgrove, 2015; Allo and Tedesco, 2005), surgical prophylaxis (Bratzler et al., 2013; Cusini et al., 2010; Davey et al., 2013), and the management of SSIs (Gaynes et al., 2001; Bergs et al., 2014), with many studies focusing on infection within the 30-day postoperative period (World Health Organization, 2016; Løwer et al., 2013). However it is important to note that SSIs represent only one of the many types of infection that can occur postoperatively.

This article provides an overview of the surgical pathway and considers infection management and antibiotic prescribing at each step of the pathway. The aim was to identify the implications for research and opportunities for system improvement.

Infection management in surgery

To date, research on surgical antibiotic prescribing has focused on the preoperative period and has largely been confined to prophylaxis, hospital inpatients, and SSIs. The WHO has recently published guidelines for the prevention of SSIs (World Health Organization, 2016). The guidelines provide detailed and evidence-based recommendations for the prevention of SSIs, including surgical antibiotic prophylaxis. The guidelines reinforce the recommendation against prolongation of surgical antibiotic prophylaxis beyond the surgical procedure as a means to prevent SSIs. In addition, they recommend worldwide surveillance of SSIs as a key component of any infection prevention and control programme. However, they do reflect that SSI surveillance is commonly poorly performed, with inconsistencies in practice, including in the type, duration, and quality of the surveillance (Table 1).

Beyond SSIs, there remains little research on the broader management of postoperative HAIs. In order to significantly mitigate against the drivers of AMR and simultaneously reduce infection rates, it is critical to optimize antibiotic use before, during, and after surgery, and to look at care settings beyond the hospital – where most patients originate from and return to. Many factors impact the risk of subsequent infection in a person who undergoes surgery; the patient's baseline health and wellbeing, patient's comorbidities, the healthcare professionals involved, and how these professionals work as a team can all influence infection-related surgical outcomes (Undre et al., 2006; Hull et al., 2011). The management of the patient in the immediate postoperative period is critical, as is their hospital length of stay. The risk of acquiring an HAI, e.g. hospital-acquired pneumonia (Garibaldi et al., 1981; Conde and Lawrence, 2008), device-related bacteraemia, or *Clostridium difficile*-associated diarrhoea, is high in the surgical patient (Krapohl et al., 2013) (Table 2). The use of invasive devices such as intravenous lines, urinary catheters, and mechanical

ventilation increases the risk of postoperative infection. In addition to this, the use of neuromuscular blocking agents during surgery to aid anaesthesia, together with reduced mobility in the immediate postoperative period, increases the risk of pneumonia (Bulka et al., 2016; Sachdev and Napolitano, 2012), which carries an estimated mortality of between 10% and 18%, even with appropriate antibiotic treatment (Croce, 2000).

Antibiotic prescribing in surgery—culture and context

The surgical pathway has many actors, steps, and actions specifically related to the management of infection and antibiotic use. Nurses, surgeons, anaesthetists, pharmacists, and allied healthcare professionals in the pre-assessment clinic, in the operating room, on the ward, and in the community during postoperative follow-up, all contribute to the care of the surgical patient. Within this complex pathway, the responsibility for the management of infection remains poorly defined at each step. In the operating room, it is unclear whether it is the anaesthetist or the operating surgeon who should assume responsibility for the timing, choice, and dose (or the need for a second dose) of the prophylactic antibiotic (Grocott and Pearse, 2012; Parker et al., 2000) (Table 1).

In the past few years, surgery has turned to aviation-inspired checklists to improve the reliability of the processes of care, and subsequently their outcomes. A number of checklists have been published and evaluated in some detail, with the best known one perhaps being the checklist developed by the WHO (Haynes et al., 2009). These checklists have identified antibiotic prophylaxis to be one of the objectives for safe surgery (World Health Organization, 2008). In the checklist, the requirement is that in the presence of the nurse, anaesthetist, and surgeon, the 'surgical team' confirm that antibiotic prophylaxis, where appropriate, is administered within 60 min of incision. However, the boundaries of responsibility for surgical antibiotic prophylaxis are not clear, specifically whose responsibility it is to decide what antibiotic prophylaxis is given and at what time (Tan et al., 2006). This is reported to be due in part to the culture and hierarchies that influence the behaviour of staff in the operating room, and also to the workflow and the environment of the operating room, which can act as obstacles to appropriate surgical antibiotic prophylaxis (Tan et al., 2006).

The lack of clarity around responsibility for antibiotic prescribing carries over into the postoperative period (Charani et al., 2017a). The responsibility for antibiotic prescribing in surgical teams is dispersed and the optimization of antibiotic therapy is often not prioritized. This leads to inappropriate antibiotic use with a prolonged duration (Leeds et al., 2016) (Table 1). This prevalent culture surrounding antibiotic decision-making in surgery needs to be understood and the expectation of surgical teams in relation to stewardship should be adjusted in view of this culture. Interventions in surgery should target the specific behaviour determinants and they should be developed in closer collaboration with surgical leaders.

Historically, most antibiotic stewardship programmes have been focused on medical specialties (Davey et al., 2013; Davey et al., 2017; Charani et al., 2017b). Within surgery, the 'low hanging fruit' is surgical antibiotic prophylaxis, with the majority of stewardship interventions targeting this single step (World Health Organization, 2016; Davey et al., 2013). Enhanced recovery after surgery (ERAS) programmes are now in place in many hospitals worldwide, and a recent systematic review has demonstrated that implementing such programmes can reduce HAIs (Grant et al., 2017). The ERAS protocols include recommendations for surgical antibiotic prophylaxis, but focus more on the entire perioperative plan for patients to promote rapid recovery and discharge postoperatively (Lassen et al., 2009). Engagement with and

Table 1

Gaps in the surgical pathway and opportunities for system change that will impact infection management and antibiotic prescribing.

	The surgical pathway		
	Perioperative period	Postoperative period	Follow-up care (primary/social/home) and surveillance
Gaps in practice that potentially impact infection management and antibiotic prescribing	Lack of clarity on whose responsibility it is to decide on the choice, dose, and timing of antibiotic prophylaxis	Gaps in the diagnosis and management of hospital-acquired infections in the postoperative patient	Follow-up care:
	Lack of clear understanding of the influence of culture and team dynamics (Tschan et al., 2015) on the implementation of the surgical checklists, e.g., WHO checklist Operating room design issues, e.g., thoroughfare, airflow disruption, or poor temperature control, or poorly designed surfaces LMIC-specific: lack of equipment to avoid infection, lack of access to antibiotics	Lack of leadership in antibiotic decision-making Lack of knowledge on the influence of culture and team dynamics on antibiotic prescribing decisions in surgery LMIC-specific: lack of access to antibiotics	Community follow-up of care in the postoperative period to ensure patient recovery Surveillance: Lack of consistency in the method of SSI surveillance, e.g.: Duration of surveillance Type of surveillance Quality of surveillance Feedback to surgical teams
Impact on clinical processes and patient outcomes	Inappropriate antibiotic prophylaxis resulting in increased risk of SSIs	Inappropriate management of hospital-acquired infections in the postoperative patient, including prolonged duration of antibiotic therapy	No system for linking SSI outcomes to antibiotic prescribing behaviours before, during, and after surgery
	Ineffective environmental precautions to prevent HCAs	Lack of postoperative critical care training leading to over-diagnosis of sepsis in the postoperative period	Lack of adherence to SSI programmes
Opportunities for change	Simple solutions such as: Clarity on roles and responsibilities for antibiotic prophylaxis in the operating room	Provision of education and training at post-graduate level Provision of national and local guidelines for antibiotic prescribing in surgery	Developing a co-ordinated package of follow-up care in the community Building on existing surveillance and developing new surveillance systems using a pragmatic approach, e.g., National Surgical Quality Improvement Programme (USA) or using mobile phone-based surveillance (LMIC)
	Monitoring operating room traffic and airflow to improve the operating room environment and prevent infections Organizational support and leadership in implementing the changes	Inclusion of and engagement with surgical teams in antibiotic stewardship interventions Developing better routes of access to antibiotics A greater understanding of the influence of culture and context on antibiotic prescribing behaviours Development of context-specific antimicrobial stewardship interventions	
Across the entire pathway	Lack of engagement with, and involvement of, the surgical teams in antibiotic prescribing interventions, together with a lack of knowledge on the optimization of antibiotic use in surgery (Charani et al., 2017a) Opportunity for change: Defining a role for antimicrobial stewardship within the surgical team, and developing targeted antibiotic prescribing interventions in surgery		

WHO, World Health Organization; LMIC, low- and middle-income countries; SSI, surgical site infection; HCAI, healthcare-associated infection.

inclusion of the surgical teams and surgical champions in the development of antimicrobial stewardship programmes is the first step in bringing about better outcomes. Organizations and national bodies can support this through legislation and policies that

promote and endorse better antibiotic prescribing across the whole surgical pathway.

Although surgical checklists are generally shown to have beneficial effects in reducing surgical infections (Treadwell et al.,

Table 2

The incidence of the most commonly occurring healthcare-associated infections in postoperative patients.

Type of hospital-acquired infection	Incidence range (as a percentage of postoperative patients) reported in the literature ^a
Hospital-acquired pneumonia	2.5% (Conde and Lawrence, 2008) to 17.5% (Garibaldi et al., 1981)
Surgical site infections	1% to 9.8% (England PH, 2016)
Urinary tract infections	2% (Chan et al., 2013) to 10% (Stéphan et al., 2006)
<i>Clostridium difficile</i> -associated diarrhoea	0.28% to 7.2% (Masgala et al., 2012; Flagg et al., 2014)

^a Depends on the type of surgery and patient population; figures are from the published literature.

2014) – though not always SSIs (Haugen et al., 2015) – overall, they have had a mixed reception (Aveling et al., 2013; Haynes et al., 2009; Russ et al., 2015). In India, for example, despite improvements in the use of the surgical site safety checklist, translation of its use in terms of patient safety outcomes has not been measured to understand the differences made in reducing SSIs, morbidity, and mortality (Patel et al., 2015). Historically, LMICs have sub-optimal investment in surgical services, and existing surgical systems are growing too slowly to meet the increasing demand (Ng-Kamstra et al., 2016). The success of interventions such as the WHO checklist targeting surgical safety are highly context-dependent and variable, and are influenced by economic, cultural, and social factors, including role identity and hierarchies within healthcare teams (Aveling et al., 2013). In particular, LMIC hierarchies have a significantly greater impact on the successful adoption of interventions in surgery (Aveling et al., 2013). Importantly, leadership, flexibility, and teamwork are required for the implementation of checklists to be effective in any setting (Walker et al., 2012; Gillespie and Marshall, 2015). This lesson applies to the successful implementation of any patient safety initiative, including interventions aiming to optimize antibiotic use (Undre et al., 2006; Dixon-Woods et al., 2013).

In particular for antibiotic prescribing, studies have described the influence of cultural determinants on prescribing outcomes (Charani et al., 2013). Antibiotic prescribing is a social act, influenced by the perceived need for clinical autonomy of individual prescribers and the existing hierarchies within teams (Charani et al., 2013). Changing antibiotic prescribing behaviours, with a view to optimizing patient-related infection outcomes, cannot be done in isolation simply by providing guidelines and policies. Existing guidelines and frameworks do not consider the variance in resources and infrastructure within LMICs (World Health Organization, 2016). In LMICs, inconsistencies in available surgical capability and resources is far more pronounced between rural and urban settings and between different LMIC countries, than it is in high-income countries where the minimum standard of care is provided in any setting. Research from different LMIC settings is lacking and is urgently needed to develop contextually sound and driven interventions that are sustainable.

The culture within specialties, organizations, and countries has the power to influence the outcome of interventions (Hofstede, 2011; Buzan, 2010; Al-Bannay et al., 2013). Culture refers to how individuals, as members of a team, learn and share knowledge in order to generate behaviours (Spradley, 1980). Culture is learned, and in order to influence practice, it needs to be studied. The culture of treatment of infection and antibiotic use in the surgical specialty has distinct features in contrast to acute medicine, because the dynamic of decision-making in the patient surgical pathway is different, as are the causal factors that influence patient-related outcomes. Understanding the contextual and cultural determinants of infection management and antibiotic prescribing in surgery is critical to the development of context-specific interventions that incorporate the need for flexibility, local leadership, and teamwork.

In an extensive review of the implementation of the Matching Michigan intervention for the prevention of blood stream infections by intensive care units in the UK (Dixon-Woods et al., 2013), the researchers reported that local culture was ‘highly consequential’ to the intervention outcome. Furthermore, the most successful units were reported to be those that had adapted the intervention to the local context and used local champions to drive and implement it. In surgery, one key step would be better engagement with the surgical leaders and the inclusion of surgeons in antimicrobial stewardship programmes. With the increasing threat of AMR, there is an urgent need to understand how to minimize the burden of infection and

optimize antibiotic use across the whole surgical pathway. Addressing the gaps in the surgical pathway will reduce the total antibiotic use and significantly mitigate drivers for AMR and the burden of infection (Table 1). Decision-making within multidisciplinary surgical teams and communication with professionals across organizational boundaries in LMICs remain unexplored. Resource limitations mean that fewer actors are involved in the surgical pathway, but there are also opportunities for innovative team composition. In the global context it is essential to redefine the antibiotic stewardship roles of the entire healthcare team across conventional organizational and professional boundaries. In addition, in LMICs the role of antibiotic stewardship in surgical care pathways is imminent, since the inappropriate use of antibiotic surgical prophylaxis has led to an increase in SSIs (Rana et al., 2013). The need for research on this topic in LMICs is particularly important to bring about efficiencies at the organizational level and to address the huge impact on out-of-pocket health expenditures and loss of earnings through prolonged postoperative recovery for the most vulnerable in society.

The way forward

Interventions in antibiotic use tend to consider and address only *one* point at a time on the patient pathway, and in the case of surgical patients, attention has been focused primarily on prophylactic antibiotic use. And yet multiple interventions have the potential to influence infection-related outcomes in the surgical patient. Social science perspectives to understand the structural, cultural contextual determinants of antibiotic use in surgery need exploring, with the potential to inform sustainable quality improvement and surgical safety initiatives. Identifying and mapping current actors and actions in the surgical specialty in different healthcare settings will inform interventions that are context-specific and relevant to the local patient population. They will also help ensure greater equity in access to safe surgery on a global scale.

Addressing antibiotic prescribing and AMR across sectors and cultures will enable surgical healthcare professionals to mobilize and drive organizational, national, and global change. Engaging with and involving surgeons and anaesthetists in the antibiotic stewardship agenda is essential, especially as it will enable the investigation of a critically unexplored domain in healthcare on a global scale, and optimize interventions within and beyond healthcare environments that have been entrenched in hierarchies and cultural norms. Addressing the gap of infection management in the surgical pathway will also impact on international policy and practice in antibiotic resistance and stewardship. Addressing the factors that influence AMR across the surgical pathway in resource-limited environments has the potential to profoundly impact the health outcomes of the millions of people who undergo surgery each year.

Funding

This work is funded by the National Institute of Health Research Imperial Biomedical Research Centre and the National Institute for Health Research Health Protection Research Unit (NIHR HPRU-2012-10047) in Healthcare Associated Infection and Antimicrobial Resistance at Imperial College London in partnership with Public Health England and the NIHR Imperial Patient Safety Translational Research Centre. NS research was supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care South London at King's College Hospital NHS Foundation Trust. NS is a member of King's Improvement Science, which is part of the NIHR CLAHRC South

London and comprises a specialist team of improvement scientists and senior researchers based at King's College London. Its work is funded by King's Health Partners (Guy's and St Thomas' NHS Foundation Trust, King's College Hospital NHS Foundation Trust, King's College London and South London and Maudsley NHS Foundation Trust), Guy's and St Thomas' Charity, the Maudsley Charity and the Health Foundation. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the UK Department of Health.

Conflict of interest

Charani, Tarrant, Ahmad, Birgand, Mendelson, Leather, Singh, Moonesinghe and Holmes have none to declare. Sevdalis is the Director of London Safety and Training Solutions Ltd, which provides team skills training and advice on a consultancy basis in hospitals and training programmes in the UK and internationally.

References

- Aiken AM, Wanyoro AK, Mwangi J, Juma F, Mugoya IK, Scott JAG. Changing use of surgical antibiotic prophylaxis in Thika Hospital, Kenya: A quality improvement intervention with an interrupted time series design. *PLoS One* 2013;8(11), doi: <http://dx.doi.org/10.1371/journal.pone.0078942>.
- Al-Bannay H, Jarus T, Jongbloed L, Yazigi M, Dean E. Culture as a variable in health research: perspectives and caveats. *Health Promot Int* 2013;1–9, doi: <http://dx.doi.org/10.1093/heapro/dat002>.
- Allo MD, Tedesco M. Operating room management: Operative suite considerations, infection control. *Surg Clin North Am* 2005;85(6):1291–7, doi: <http://dx.doi.org/10.1016/j.suc.2005.09.001>.
- A. Point Prevalence Survey of Healthcare-Associated Infections and Antimicrobial Use in European Hospitals 2011–2012. 2013.
- NHS. NHS Confed.. 2016 Available at: <http://www.nhsconfed.org/resources/key-statistics-on-the-nhs>. Accessed December 15, 2016.
- Aveling E-L, McCulloch P, Dixon-Woods M. A qualitative study comparing experiences of the surgical safety checklist in hospitals in high-income and low-income countries. *BMJ Open* 2013;3:e003039, doi: <http://dx.doi.org/10.1136/bmjopen-2013-003039>.
- Bergs J, Hellings J, Cleemput I, Zurel [340_TD\$DIFF]O, De Troyer V, Van Hiel M, et al. Systematic review and meta-analysis of the effect of the World Health Organization surgical safety checklist on postoperative complications. *Br J Surg* 2014;101(FEBRUARY 2014):150–8, doi: <http://dx.doi.org/10.1002/bjs.9381>.
- Birgand G, Azevedo C, Toupet G, Pissard-Gibollet [341_TD\$DIFF]R, Grandbastien B, Fleury E, et al. Attitudes, risk of infection and behaviours in the operating room (the ARIBO Project): a prospective, cross-sectional study. *BMJ Open* 2014;4(1):e004274, doi: <http://dx.doi.org/10.1136/bmjopen-2013-004274>.
- Boucher HW, Talbot GH, Bradley JS, Edwards [342_TD\$DIFF]E, Gilbert D, Rice LB, et al. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin Infect Dis* 2009;48(1):1–12, doi: [http://dx.doi.org/10.1016/S0090-3671\(10\)79814-1](http://dx.doi.org/10.1016/S0090-3671(10)79814-1).
- Bratzler DW, Dellinger EP, Olsen KM, Perl [343_TD\$DIFF]TM, Auwaerter PG, Bolon MK, et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Surg Infect (Larchmt)* 2013;14(1):73–156, doi: <http://dx.doi.org/10.1089/sur.2013.9999>.
- Bulka CM, Terekhov MA, Martin BJ, Dmochowski RR, Hayes RM, Ehrenfeld JM. Nondepolarizing Neuromuscular Blocking Agents, Reversal, and Risk of Postoperative Pneumonia. *Anesthesiology* 2016;4:1–8, doi: <http://dx.doi.org/10.1097/ALN.0000000000001279>.
- Buzan B. Culture and international society. *Int Aff* 2010;86(1):1–25, doi: <http://dx.doi.org/10.1111/j.1468-2346.2010.00866.x>.
- Chan JYK, Semenov YR, Gourin CG. Postoperative Urinary Tract Infection and (Short-Term) Outcomes and Costs in Head and Neck Cancer Surgery. *Otolaryngol Head Neck Surg* 2013;148:602–10, doi: <http://dx.doi.org/10.1177/0194599812474595>.
- Charani [344_TD\$DIFF]E, Castro-Sanchez E, Sevdalis N, Kyrtatsis [345_TD\$DIFF]Y, Drumright L, Shah N, et al. Understanding the determinants of antimicrobial prescribing within hospitals: the role of "prescribing etiquette". *Clin Infect Dis* 2013;57(2):188–96, doi: <http://dx.doi.org/10.1093/cid/cit212>.
- Charani E, Tarrant C, Moorthy K, Sevdalis N, Brennan L, Holmes AH. Understanding antibiotic decision making in surgery – a qualitative analysis. *Clin Microbiol Infect* 2017a;, doi: <http://dx.doi.org/10.1016/j.cmi.2017.03.013>.
- Charani E, Gharbi M, Moore LSP, Castro-Sánchez [347_TD\$DIFF]E, Lawson W, Gilchrist M, et al. Effect of adding a mobile health intervention to a multimodal antimicrobial stewardship programme across three teaching hospitals: an interrupted time series study. *J Antimicrob Chemother* 2017b;, doi: <http://dx.doi.org/10.1093/jac/dkx040>.
- Conde M, Lawrence V. Post-operative pulmonary infections. *BMJ Clin Evid* 2008;1–18.
- Cosgrove MS. Infection control in the operating room. *Crit Care Nurs Clin North Am* 2015;27(1):79–87, doi: <http://dx.doi.org/10.1016/j.cnc.2014.10.004>.
- Croce MA. Postoperative pneumonia. *Am Surg* 2000;66(2):133–7, doi: [http://dx.doi.org/10.1016/S0149-7944\(02\)00661-X](http://dx.doi.org/10.1016/S0149-7944(02)00661-X).
- Cusini A, Rampini SK, Bansal V, Ledergerber [348_TD\$DIFF]B, Kuster SP, Ruef C, et al. Different patterns of inappropriate antimicrobial use in surgical and medical units at a tertiary care hospital in Switzerland: A prevalence survey. *PLoS One* 2010;5(11):1–8, doi: <http://dx.doi.org/10.1371/journal.pone.0014011>.
- Davey P, Brown E, Charani E, Fenelon [349_TD\$DIFF]L, Im G, Holmes A, et al. Interventions to improve antibiotic prescribing practices for hospital inpatients (Review). *Cochrane database Syst Rev* 2013;4(5):CD003543, doi: <http://dx.doi.org/10.1002/14651858.CD003543.pub3>.
- Davey P, Brown [350_TD\$DIFF]E, Charani E, Fenelon L, Im G, Holmes A, et al. Interventions to improve antibiotic prescribing practices for hospital inpatients. *Cochrane Database Syst Rev* 2017;, doi: <http://dx.doi.org/10.1002/14651858.CD003543.pub4>.
- Dixon-Woods M, Leslie M, Tarrant C, Bion J. Explaining Matching Michigan: an ethnographic study of a patient safety program. *Implement Sci* 2013;8(1):70, doi: <http://dx.doi.org/10.1186/1748-5908-8-70>.
- England PH. Surveillance of Surgical Site Infections in NHS Hospitals in England. 2016 London.
- Flagg A, Koch CG, Schiltz N, Chandran Pillai [351_TD\$DIFF]A, Gordon SM, Petterson GB, et al. Analysis of Clostridium difficile infections after cardiac surgery: epidemiologic and economic implications from national data. *J Thorac Cardiovasc Surg* 2014;148(5):2404–9, doi: <http://dx.doi.org/10.1016/j.jtcvs.2014.04.017>.
- Garibaldi RA, Britt MR, Coleman ML, Reading JC, Pace NL. Risk factors for postoperative pneumonia. *Am J Med* 1981;70(3):677–80, doi: [http://dx.doi.org/10.1016/0002-9343\(81\)90595-7](http://dx.doi.org/10.1016/0002-9343(81)90595-7).
- Gaynes R, Culver D, Horan T, Edwards J, Richards C, Tolson J. Surgical Site Infection (SSI) rates in the United States, 1992–1998: The National Nosocomial Infections Surveillance System Basic SSI Risk Index. *Clin Infect Dis* 2001;33(Suppl 2):S69–77, doi: <http://dx.doi.org/10.1086/321860>.
- Gillespie BM, Marshall A. Implementation of safety checklists in surgery: a realist synthesis of evidence. *Implement Sci* 2015;10:137, doi: <http://dx.doi.org/10.1186/s13012-015-0319-9>.
- Grant M, Yang D, Wu C, Makary M, Wick E. Impact of Enhanced Recovery After Surgery and Fast Track Surgery Pathways on Healthcare-associated Infections: Results From a Systematic Review and Meta-analysis. *Ann Surg* 2017;265(1):68–79.
- Grocott MPW, Pearse RM. Perioperative medicine: The future of anaesthesia?. *Br J Anaesth* 2012;108(5):723–6, doi: <http://dx.doi.org/10.1093/bja/aes124>.
- Haugen AS, Sjøteland E, Almeland SK, Sevdalis [352_TD\$DIFF]N, Vonon B, Eide GE, et al. Effect of the World Health Organization Checklist on patient outcomes: a stepped wedge cluster randomized controlled trial. *Ann Surg* 2015;261(5):821–8, doi: <http://dx.doi.org/10.1097/SLA.0000000000000716>.
- Haynes A, Weiser TG, Berry WR, Lipsitz [353_TD\$DIFF]S, Breizat A-H, Dellinger E, et al. A Surgical Safety Checklist to Reduce Morbidity and Mortality in a Global Population. *N Engl J Med* 2009;360(5):491–9, doi: <http://dx.doi.org/10.1056/NEJMs0810119>.
- Hofstede G. Dimensionalizing Cultures: The Hofstede Model in Context. *Online Readings Psychol Cult* 2011;2(1):1–26, doi: <http://dx.doi.org/10.9707/2307-0919.1014>.
- Hull L, Arora S, Kassab E, Kneebone R, Sevdalis N. Observational teamwork assessment for surgery: content validation and tool refinement. *J Am Coll Surg* 2011;212(2), doi: <http://dx.doi.org/10.1016/j.jamcollsurg.2010.11.001> 234–243-5.
- Krapohl GL, Morris AM, Cai S, Englesbe [355_TD\$DIFF]MJ, Aronoff DM, Campbell DA, et al. Preoperative risk factors for postoperative Clostridium difficile infection in colectomy patients. *Am J Surg* 2013;205(3):343–8, doi: <http://dx.doi.org/10.1016/j.amjsurg.2012.10.028>.
- Løwer HL, Eriksen HM, Aavitsland P, Skjeldstad FE. Methodology of the Norwegian Surveillance System for Healthcare-Associated Infections: The value of a mandatory system, automated data collection, and active postdischarge surveillance. *Am J Infect Control* 2013;41:591–6, doi: <http://dx.doi.org/10.1016/j.ajic.2012.09.005>.
- Lassen K, Soop M, Nygren J. Consensus Review of Optimal Perioperative Care in Colorectal Surgery. *Am Med Assoc* 2009;144(10):961–9, doi: <http://dx.doi.org/10.1001/archsurg.2009.170>.
- Leeds IL, Fabrizio A, Cosgrove SE, Wick EC. Treating Wisely: The Surgeon's Role in Antibiotic Stewardship. *Ann Surg* 2016;1, doi: [http://dx.doi.org/10.1097/SLA.0000000000002034.XX\(Xx\)](http://dx.doi.org/10.1097/SLA.0000000000002034.XX(Xx)).
- Masgala A, Chronopoulos E, Nikolopoulos G, Sourlas [356_TD\$DIFF]J, Lallou S, Brilakis E, et al. Risk factors affecting the incidence of infection after orthopaedic surgery: The role of chemoprophylaxis. *Cent Eur J Public Health* 2012;20(4):252–6.
- Ng-Kamstra JS, Greenberg SLM, Abdullah F, Amado [358_TD\$DIFF]V, Anderson GA, Cossa M, et al. Global Surgery 2030: a roadmap for high income country actors. *BMJ Glob Heal* 2016;1(1):1–12, doi: <http://dx.doi.org/10.1136/bmjgh-2015-000011>.
- Parker BM, Tetzlaff JE, Litaker DL, Maurer WG. Redefining the preoperative evaluation process and the role of the anesthesiologist. *J Clin Anesth* 2000;12(5):350–6, doi: [http://dx.doi.org/10.1016/S0952-8180\(00\)00169-0](http://dx.doi.org/10.1016/S0952-8180(00)00169-0).

- Patel A, Sanghi V, Gupta V. Implementation of surgical safety checklist for all invasive procedures. *J Natl Accredited Board Hosp Healthc Provid* 2015;2(2):41–6, doi:<http://dx.doi.org/10.4103/2319-1880.174347>.
- Rana DA, Malhotra SD, Patel VJ. Inappropriate surgical chemoprophylaxis and surgical site infection rate at a tertiary care teaching hospital. *Brazilian J Infect Dis* 2013;17(1):48–53, doi:<http://dx.doi.org/10.1016/j.bjid.2012.09.003>.
- Russ S, Rout S, Caris J, Mansell [360_TD\$DIFF], Davies R, Mayer E, et al. Measuring variation in use of the WHO surgical safety checklist in the operating room: a multicenter prospective cross-sectional study. *J Am Coll Surg* 2015;220(1), doi:<http://dx.doi.org/10.1016/j.jamcollsurg.2014.09.021> 1–11.e4.
- Sachdev G, Napolitano LM. Postoperative Pulmonary Complications: Pneumonia and Acute Respiratory Failure. *Surg Clin North Am* 2012;92(2):321–44, doi:<http://dx.doi.org/10.1016/j.suc.2012.01.013>.
- Spradley JP. Ethnography for What?. *Participant Observation*. . p. 13–25.
- Stéphan F, Sax H, Wachsmuth M, Hoffmeyer P, Clergue F, Pittet D. Reduction of urinary tract infection and antibiotic use after surgery: a controlled, prospective, before-after intervention study. *Clin Infect Dis* 2006;42(11):1544–51, doi:<http://dx.doi.org/10.1086/503837>.
- Tan JA, Naik VN, Lingard L, Sussman [361_TD\$DIFF]JA, Mccaffrey CB, Leary DB, et al. Exploring obstacles to proper timing of prophylactic antibiotics for surgical site infections. *Qual Saf Health Care* 2006;15(1):32–8, doi:<http://dx.doi.org/10.1136/qshc.2004.012534>.
- The Royal College of Anaesthetists. *Perioperative Medicine the Pathway to Better Surgical Care* London. 2015.
- Treadwell JR, Lucas S, Tsou AY. Surgical checklists: a systematic review of impacts and implementation. *BMJ Qual Saf* 2014;23(4):299–318, doi:<http://dx.doi.org/10.1136/bmjqs-2012-001797>.
- Tschan F, Seelandt J, Keller S, Semmer [356_TD\$DIFF][362_TD\$DIFF]N, Kurmann A, Candinas D, et al. Impact of case-relevant and case-irrelevant communication within the surgical team on surgical-site infection. *Br J Surg* 2015;102:1718–25.
- Undre S, Sevdalis N, Healey AN, Darzi A, Vincent CA. Teamwork in the operating theatre: Cohesion or confusion?. *J Eval Clin Pract* 2006;12(2):182–9, doi:<http://dx.doi.org/10.1111/j.1365-2753.2006.00614.x>.
- Walker IA, Reshamwalla S, Wilson IH. Surgical safety checklists: Do they improve outcomes?. *Br J Anaesth* 2012;109(1):47–54, doi:<http://dx.doi.org/10.1093/bja/aes175>.
- World Health Organization. *WHO Surgical Safety Checklist Implementation Manual (First Edition)*. Spring 2008;1–28, doi:<http://dx.doi.org/10.1111/j.1749-4486.2009.02129.x>.
- World Health Organization. *Global guidelines for the prevention of surgical site infection*. 2016 Available at: <http://www.who.int/gpsc/global-guidelines-web.pdf>.