



## King's Research Portal

DOI:

[10.1016/j.scr.2016.01.006](https://doi.org/10.1016/j.scr.2016.01.006)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Jacquet, L., Hewitson, H., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Hobbs, C., Stephenson, E., & Ilic, D. (2016). Generation of KCL028 research grade human embryonic stem cell line carrying a mutation in the HTT gene. *Stem Cell Research*, 16(2), 278-281. <https://doi.org/10.1016/j.scr.2016.01.006>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.


### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Lab Resource: Stem Cell Line

## Generation of KCL028 research grade human embryonic stem cell line carrying a mutation in the *HTT* gene



Lauren Jacquet<sup>a</sup>, Heema Hewitson<sup>a</sup>, Victoria Wood<sup>a</sup>, Neli Kadeva<sup>a</sup>, Glenda Cornwell<sup>a</sup>, Stefano Codognotto<sup>a</sup>, Carl Hobbs<sup>b</sup>, Emma Stephenson<sup>a</sup>, Dusko Ilic<sup>a</sup>

<sup>a</sup> Stem Cell Laboratories, Division of Women's Health, Faculty of Life Sciences and Medicine, King's College London and Assisted Conception Unit, Guy's Hospital, London, United Kingdom

<sup>b</sup> Histology Laboratory, Wolfson Centre for Age-Related Diseases, Faculty of Life Sciences and Medicine, King's College London, London, United Kingdom

## ARTICLE INFO

## Article history:

Received 31 December 2015

Received in revised form 12 January 2016

Accepted 12 January 2016

Available online 14 January 2016

## ABSTRACT

The KCL028 human embryonic stem cell line was derived from an embryo donated for research that carried an autosomal dominant mutation affecting one allele of the *HTT* gene encoding huntingtin (43 trinucleotide repeats; 21 for the normal allele). The ICM was isolated using laser microsurgery and plated on  $\gamma$ -irradiated human foreskin fibroblasts. Both the derivation and cell line propagation were performed in an animal product-free environment. Pluripotent state and differentiation potential were confirmed by in vitro and in vivo assays.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Resource table

Name of stem cell line	KCL028
Institution	King's College London, London UK
Derivation team	Neli Kadeva, Victoria Wood, Glenda Cornwell, Stefano Codognotto, Emma Stephenson
Contact person and email	Dusko Ilic, email: <a href="mailto:dusko.ilic@kcl.ac.uk">dusko.ilic@kcl.ac.uk</a>
Date archived/stock date	May 25, 2011
Type of resource	Biological reagent: cell line
Sub-type	Human pluripotent stem cell line
Origin	Human embryo
Key marker expression	Pluripotent stem cell markers: NANOG, OCT4, TRA-1-60, TRA-1-81, alkaline phosphatase (AP) activity
Authentication	Identity and purity of line confirmed
Link to related literature (direct URL links and full references)	1) Ilic, D., Stephenson, E., Wood, V., Jacquet, L., Stevenson, D., Petrova, A., Kadeva, N., Codognotto, S., Patel, H., Semple, M., Cornwell, G., Ogilvie, C., Braude, P., 2012. Derivation and feeder-free propagation of human embryonic stem cells under xeno-free conditions. <i>Cytotherapy</i> . 14 (1), 122–128. doi: 10.3109/14653249.2011.623692 <a href="http://www.ncbi.nlm.nih.gov/pubmed/22029654">http://www.ncbi.nlm.nih.gov/pubmed/22029654</a> 2) Stephenson, E., Jacquet, L., Miere, C., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Dajani, Y., Braude, P., Ilic, D., 2012. Derivation and propagation of human embryonic stem cell lines from frozen embryos in an animal product-free environment. <i>Nat. Protoc.</i> 7 (7), 1366–1381. doi: 10.1038/nprot.2012.080 <a href="http://www.ncbi.nlm.nih.gov/pubmed/22722371">http://www.ncbi.nlm.nih.gov/pubmed/22722371</a>

(continued)

Name of stem cell line	KCL028
Information in public databases	3) Jacquet, L., Neueder, A., Földes, G., Karagiannis, P., Hobbs, C., Jolinon, N., Mioulane, M., Sakai, T., Harding, S.E., Ilic, D., 2015. Three Huntington's disease specific mutation-carrying human embryonic stem cell lines have stable number of CAG repeats upon in vitro differentiation into cardiomyocytes. <i>PLoS One</i> . 10(5), e0126860. <a href="http://doi.org/10.1371/journal.pone.0126860">http://doi.org/10.1371/journal.pone.0126860</a> <a href="http://www.ncbi.nlm.nih.gov/pubmed/25993131">http://www.ncbi.nlm.nih.gov/pubmed/25993131</a> KCL028 is a National Institutes of Health (NIH) registered hESC line NIH registration number: 0224 NIH approval number: NIHhESC-13-0224 <a href="http://grants.nih.gov/stem_cells/registry/current.htm?id=664">http://grants.nih.gov/stem_cells/registry/current.htm?id=664</a>
Ethics	The hESC line KCL028 is derived under license from the UK Human Fertilisation and Embryology Authority (research license numbers: R0075 and R0133) and also has local ethical approval (UK National Health Service Research Ethics Committee Reference: 06/Q0702/90). Informed consent was obtained from all subjects and the experiments conformed to the principles set out in the WMA Declaration of Helsinki and the NIH Belmont Report. No financial inducements are offered for donation.

## Resource details

Consent signed	Jan 27, 2011
Embryo used	May 04, 2011
UK Stem Cell Bank deposit approval	Dec 01, 2011 Reference: SCSC11-47

(continued)


Sex	Male 46, XY
Grade	Research
Disease status (Fig. 1)	Mutation affecting one allele of the <i>HTT</i> gene encoding huntingtin (~43 CAG repeats; 21 for the normal allele) associated with Huntington's disease (Ilic et al., 2015)
Karyotype (aCGH)	No imbalance detected.
DNA fingerprint	Allele sizes (in bp) of 17 microsatellite markers specific for chromosomes 13, 18 and 21 (Jacquet et al., 2015)
HLA typing	HLA-1: 02,03; –B: 07,35; –C: 04,07; Bw: 6; DRB1: 01; DQB1: 05
Viability testing	Pass
Pluripotent markers (immunostaining) (Fig. 2)	NANOG, OCT4, TRA-1-60, TRA-1-81, AP activity (Jacquet et al., 2015)
Three germ layers differentiation in vitro (immunostaining) (Fig. 3)	Endoderm: AFP ( $\alpha$ -fetoprotein); Ectoderm: TUBB3 (tubulin, $\beta$ 3 class III); Mesoderm: ACTA2 (actin, $\alpha$ 2, smooth muscle) (Jacquet et al., 2015)
Three germ layer differentiation in vivo (teratomas) (Fig. 4)	Endoderm: AFP, GATA4. Ectoderm: TUBB3, GFAP (glial fibrillary acidic protein). Mesoderm: DES (desmin), Alcian Blue and periodic acid–Schiff (PAS)-stained cartilage (Jacquet et al., 2015)
Targeted differentiation	Cardiomyocytes: TNNT2 (cardiac troponin T) immunostaining
Sibling lines available	KCL027

We generated KCL028 research grade hESC line following protocols, established previously (Ilic et al., 2012; Stephenson et al., 2012). The expression of the pluripotency markers was tested after freeze/thaw cycle (Ilic et al., 2012). Differentiation potential into three germ layers was verified in vitro (Jacquet et al., 2015) and in vivo (Jacquet et al., 2015).


## Materials and methods

### Consenting process

We distribute Patient Information Sheet (PIS) and consent form to the in vitro fertilization (IVF) patients if they opted to donate to research



**Fig. 1.** Genetic pedigree tree. The couple undergoing IVF had 16 embryos in this particular cycle. Three embryos were normal, whereas those that carried the mutation in *HTT* were donated for research. We derived hESC lines from two of them.



**Fig. 2.** Expression of pluripotency markers. Pluripotency is confirmed by immunostaining (Oct. 4, Nanog, TRA-1-60, TRA-1-81) and alkaline phosphatase (AP) activity assay. Actin stress fibers, visualized with rhodamine-phalloidin (red), are present in both feeders and hES cell colonies, whereas AP activity (green) is detected only in hES cells. Scale bar, 50  $\mu$ m.

embryos that were stored for 5 or 10 years. They mail signed consent back to us and that might be months after the PIS and consent were mailed to them. If in the meantime new versions of PIS/consent are implemented, we do not send these to the patients or ask them to re-sign; the whole process is done with the version that was given them initially. The PIS/consent documents (PGD-V.8) were created on Jul. 01, 2010. HFEA Code of Practice that was in effect at the time of document creation: Edition 8 – R.2 (<http://www.hfea.gov.uk/2999.html>). The donor couple signed the consent on Jan. 12, 2011. HFEA Code of Practice that was in effect at the time of donor signature: Edition 8 – R.2. HFEA Code of Practice Edition 8 – R.2 was in effect: Apr. 07, 2010–Apr. 06, 2011.

### Embryo culture and micromanipulation


Embryo culture and laser-assisted dissection of inner cell mass (ICM) were carried out as previously described in details (Ilic et al., 2012; Stephenson et al., 2012). The cellular area containing the ICM was then washed and transferred to plates containing mitotically inactivated human neonatal foreskin fibroblasts (HFF).

### Cell culture

ICM plated on mitotically inactivated HFF were cultured as described (Ilic et al., 2012; Stephenson et al., 2012). TE cells were removed mechanically from outgrowth (Ilic et al., 2007; Ilic et al., 2010). hESC colonies were expanded and cryopreserved at the third passage.

### Viability test

Straws with the earliest frozen passage (p. 2–3) are thawed and new colonies are counted three days later. These colonies are then expanded up to passage 8, at which point cells were part frozen and part subjected to standard battery of tests (pluripotency markers, in vitro and in vivo differentiation capability, genetics, sterility, mycoplasma).



**Fig. 3.** Differentiation of three germ layers in vitro is confirmed by detection of markers: smooth muscle actin (ACTA2, red) for mesoderm,  $\beta$ -III tubulin (TUBB3, red) for ectoderm and  $\alpha$ -fetoprotein (AFP, red) for endoderm. Nuclei are visualized with Hoechst 33342 (blue). Scale bar, 100  $\mu$ m.

#### Pluripotency markers

Pluripotency was assessed using two different techniques: enzymatic activity assay [alkaline phosphatase (AP) assay] and immunostaining as described (Ilic et al., 2012; Stephenson et al., 2012; Petrova et al., 2014).

#### Differentiation

Spontaneous differentiation into three germ layers was assessed in vitro as described (Ilic et al., 2012; Stephenson et al., 2012; Petrova et al., 2014) and in vivo (Jacquet et al., 2015). Targeted differentiation in cardiomyocytes followed the protocols described earlier (Jacquet et al., 2015; Laflamme et al., 2007).

#### Genotyping

DNA was extracted from hESC cultures using a Chemagen DNA extraction robot according to the manufacturer's instructions. Amplification of polymorphic microsatellite markers was carried out as described (Ilic et al., 2012). Allele sizes were recorded to give a unique fingerprint of each cell line.

#### Array comparative genomic hybridization (aCGH)

aCGH was performed as described in details (Ilic et al., 2012).

#### HLA typing

HLA-A, -B and -DRB1 typing was performed with a PCR sequence-specific oligonucleotide probe (SSOP; Luminex, Austin, TX, USA) hybridization protocol at the certified Clinical Transplantation Laboratory, Guy's

and St Thomas' NHS Foundation Trust and Serco Plc. (GSTS) Pathology (Guy's Hospital, London, UK) as described (Jacquet et al., 2013).

#### Author disclosure statement


There are no competing financial interests in this study.

#### Acknowledgments

This work was supported by the UK Medical Research Council grants G0701172 and G0801061. We thank Dr. Yacoub Khalaf, Director of the Assisted Conception Unit of Guy's and St Thomas' NHS Foundation Trust and his staff for supporting the research program. We are especially indebted to Prof. Peter Braude and to the patients who donated embryos.

#### References

- Ilic, D., Genbacev, O., Krtolica, A., 2007. Derivation of hESC from intact blastocysts. *Curr. Protoc. Stem Cell Biol.* (Chapter 1: Unit 1 A.2).
- Ilic, D., Caceres, E., Lu, S., Julian, P., Foulk, R., Krtolica, A., 2010. Effect of karyotype on successful human embryonic stem cell derivation. *Stem Cells Dev.* 19 (1), 39–46.
- Ilic, D., Stephenson, E., Wood, V., Jacquet, L., Stephenson, D., Petrova, A., Kadeva, N., Codognotto, S., Patel, H., Semple, M., Cornwell, G., Ogilvie, C., Braude, P., 2012. Derivation and feeder-free propagation of human embryonic stem cells under xeno-free conditions. *Cytotherapy* 14 (1), 122–128.
- Jacquet, L., Stephenson, E., Collins, R., Patel, H., Trussler, J., Al-Bedaery, R., Renwick, P., Ogilvie, C., Vaughan, R., Ilic, D., 2013. Strategy for the creation of clinical grade hESC line banks that HLA-match a target population. *EMBO Mol. Med.* 5 (1), 10–17.
- Jacquet, L., Neueder, A., Földes, G., Karagiannis, P., Hobbs, C., Jolinon, N., Mioulane, M., Sakai, T., Harding, S.E., Ilic, D., 2015. Three Huntington's disease specific mutation-carrying human embryonic stem cell lines have stable number of CAG repeats upon in vitro differentiation into cardiomyocytes. *PLoS ONE* 10 (5), e0126860.
- Laflamme, M.A., Chen, K.Y., Naumova, A.V., Muskheli, V., Fugate, J.A., Dupras, S.K., Reinecke, H., Xu, C., Hassaniipour, M., Police, S., O'sullivan, C., Collins, L., Chen, Y., Minami, E., Gill, E.A., Ueno, S., Yuan, C., Gold, J., Murry, C.E., 2007. Cardiomyocytes derived from human embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts. *Nat. Biotechnol.* 25 (9), 1015–1024.



**Fig. 4.** Differentiation of three germ layers in vivo. Teratomas were encapsulated and did not invade surrounding tissue. Sections are counterstained with hematoxylin and eosin and specific stains are brown (immunohistochemistry). Germ layer marker: DES for mesoderm, TUBB3 for ectoderm, and GATA4 for endoderm. Scale bars are 100  $\mu$ m.

- Petrova, A., Celli, A., Jacquet, L., Dafou, D., Crumrine, D., Hupe, M., Arno, M., Hobbs, C., Cvorovic, A., Karagiannis, P., Devito, L., Sun, R., Adame, L.C., Vaughan, R., McGrath, J.A., Mauro, T.M., Ilic, D., 2014. 3D in vitro model of a functional epidermal permeability barrier from human embryonic stem cells and induced pluripotent stem cells. *Stem Cell Rep.* 2 (5), 675–689.
- Stephenson, E., Jacquet, L., Miere, C., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Dajani, Y., Braude, P., Ilic, D., 2012. Derivation and propagation of human embryonic stem cell lines from frozen embryos in an animal product-free environment. *Nat. Protoc.* 7 (7), 1366–1381.