King's Research Portal

DOI:

[10.12968/bjom.2018.26.1.31](https://doi.org/10.12968/bjom.2018.26.1.31)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Carlisle, N., Carter, J., Radford, S., & Shennan, A. (2018). Women's experiences of tests and procedures carried out at a preterm birth surveillance clinic. *British Journal of Midwifery*, 26(1), 31-34.
<https://doi.org/10.12968/bjom.2018.26.1.31>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

British Journal of Midwifery

Women's experiences of tests and procedures carried out at a preterm birth surveillance clinic --Manuscript Draft--

Manuscript Number:	
Full Title:	Women's experiences of tests and procedures carried out at a preterm birth surveillance clinic
Short Title:	Women's experiences of tests and procedures carried out at a PTB clinic
Article Type:	Original research
Keywords:	Preterm birth Late miscarriage Speculum examination Transvaginal scan
Corresponding Author:	Naomi Carlisle, MRes St Thomas Hospital, London London, UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	St Thomas Hospital, London
Corresponding Author's Secondary Institution:	
First Author:	Naomi Carlisle, MRes
First Author Secondary Information:	
Order of Authors:	Naomi Carlisle, MRes Samara Radford Jenny Carter Andrew Shennan
Order of Authors Secondary Information:	
Abstract:	This study sought to explore the experiences and views of pregnant women at high-risk of preterm birth who were undergoing tests requiring a speculum examination and procedures at a preterm birth surveillance clinic. There was a total of 102 participants who completed the questionnaire. Overall 97% (n=99) of women found the speculum examination acceptable and 88% (n=90) stated they would be happy to have the examination again if required. This was comparable to 95% (n=97) of women who stated they found the transvaginal ultrasound examination acceptable. Our findings suggest that women at risk of preterm birth find speculum examinations and transvaginal ultrasound scans for cervical length measurement acceptable.
Suggested Reviewers:	
Additional Information:	
Question	Response
Please enter the word count of your manuscript	1479

Women's experiences of tests and procedures carried out at a preterm birth surveillance clinic

Introduction

Preterm birth remains a major cause of neonatal morbidity and mortality, and in the United Kingdom (UK), 7.8% of babies born were born before 37 weeks gestation in the year 2010 (WHO, 2012). Some UK hospitals offer a specialist preterm birth service for women deemed to be at risk of a preterm birth or late miscarriage including those with a history of preterm birth, late miscarriage or cervical surgery. However practice in specialist preterm birth clinics varies in both surveillance methods and clinical management (Sharp & Alfirevic, 2014).

Surveillance methods are used to identify those women who may benefit from intervention, such as cerclage or progesterone treatment and closer monitoring (Min *et al.*, 2016). Methods include transvaginal ultrasound measurement of cervical length, and collection of vaginal fluid for evaluation using predictive biomarker tests such as Fetal Fibronectin, PartoSure™ or Actim® Partus.

Most biomarker tests require the use of a speculum, however one test, PartoSure™, does not and the reduced need for a speculum examination with a speculum may be more acceptable to some women. While there is more evidence to support the utility of quantitative Fetal Fibronectin (qfFn) in predicting preterm birth compared to Partosure™, Fetal Fibronectin does require the use of a speculum (Abbott *et al.*, 2013; Abbott *et al.*, 2015; Kurtzman *et al.*, 2009).

Although previous studies exploring transvaginal ultrasound scans in pregnancy have found them to be acceptable to women (Atalabi *et al.*, 2012; Bello & Odeku, 2015; Clement *et al.*, 2003), we found no studies focusing on women's experiences of speculum examinations in a pregnancy at high-risk of preterm delivery.

This study sought to explore the experiences and views of pregnant women at high-risk of preterm birth who were undergoing tests requiring a speculum examination and procedures at a preterm birth surveillance clinic.

Methods

Participants were a sub-set of women enrolled in the ***** study (REC Ref.*****). Women were enrolled through convenience sampling as they attended an appointment at an antenatal preterm surveillance clinic at a large UK inner-city teaching hospital between February and August 2011. All women presenting to the clinic were provided with the study questionnaire and were asked to return their completed paperwork anonymously after their appointment to the clinic receptionist. This ensured that women did not feel pressured by the clinical research staff working in the clinic.

Data collected included demographic information, parity and gestational age. Eight questions were asked about their experiences of the speculum examination and transvaginal ultrasound scan for cervical length measurement. These eight questions could be answered “yes”, “no” or “I don’t know”. The data were then collected and analysed on an Excel spreadsheet.

Results

There was a total of 102 participants who completed the questionnaire. Overall 97% (n=99) of women found the speculum examination acceptable and 88% (n=90) stated they would be happy to have the examination again if required. This was comparable to 95% (n=97) of women who stated they found the transvaginal ultrasound examination acceptable.

Participant characteristics

The age of the participants at time of completing the questionnaire ranged from 17 years to 44 years, with a mean and median age of 32 years.

The demographics of the participants can be seen in Table 1.

Table 1 Demographics of participants

	Number of participants n=	Percentage of participants %
Age Range		
> 20	2	2
20-24	7	7
25-29	14	14
30-34	45	44
35-39	25	26
40-44	4	4
No response	5	5
Ethnicity		
White British	29	28
White Other	16	16
Black African	27	26
Black Caribbean	8	8
Asian	6	6
Mixed race	3	3
Other	12	12
No response	1	1
Parity		
Nulliparous	53	52

Multiparous

49

48

The gestational age (GA) of the participants at the time of completing the questionnaire ranged from 18 to 34 weeks, with a mean and median GA of 22 weeks. 5 participants did not disclose their gestation.

Experience of having speculum examinations and procedures

Results for the remainder of the survey are presented in Table 2 and Figure 1.

Table 2 Patient responses to the questionnaire

	Yes		No		I don't know		No response	
	n	%	n	%	n	%	n	%
1. Did you find the speculum examination acceptable?	99	97	1	1	2	2	0	0
2.If "no", would you accept the test if it did not involve a speculum?	1	1	-	-	-	-	-	-
3. Did you understand why the test was done?	100	98	1	1	0	0	1	1

**4. Did you understand
what the test result
meant?**

89   86   0   0   5   5   8   8

**5. Would you be happy
to have the test again in
this pregnancy?**

90   88   4   4   5   5   3   3

**6. Would you be
prepared to take the
swab yourself without
the speculum?**

48   47   35   34   15   15   4   4

**7. Do you find the
cervical length scan
acceptable?**

97   95   2   2   2   2   1   1

**8. Given the choice,
would you come back
to this clinic?**

97   95   2   2   2   2   1   1

Total n=102

Figure 1 Stacked column showing the participants' experiences of tests and procedures at the preterm birth surveillance clinic

Discussion

This is the first study exploring women's experience of speculum examination for preterm surveillance. Studies exploring women's experiences of a vaginal speculum examination suggest that they are associated with anxiety, embarrassment and discomfort (Hoyo *et al.*, 2005; Yanikkerem *et al.*, 2009). Anxiety in pregnancy is closely related to the woman's perception of her pregnancy being high-risk (Bayrampour *et al.*, 2012). Women attending preterm birth surveillance clinics are aware

that they are at high-risk, and therefore likely to be anxious and apprehensive even before any procedures or tests are carried out.

The majority of participants (99/102, 97%) found the speculum examination acceptable, with only one participant (1%) finding the speculum unacceptable. This one participant responded that they would have accepted the Fetal Fibronectin test if it had not involve a speculum. These results demonstrate that the vast majority of this pregnant cohort found the speculum examination acceptable.

This study found that 88% of participants who responded would be happy to have the test again in their current pregnancy, while 4% would not be happy to. These results are encouraging, as they suggest that this would not prevent the majority of women adhering to premature birth surveillance due to lack of acceptability with speculum use. Previous studies suggest that discomfort associated with speculum examinations may have led to some women not attending appointments for cervical cancer screening (Hoyo *et al.*, 2005).

Of participants who responded, 47% of women would be prepared to take the swab themselves without a speculum, while 34% of women would not be prepared to. As previous studies have found women's experiences of speculums as anxiety provoking, embarrassing and uncomfortable this was surprising. For example, a pilot study on speculum 'self-insertion' found that the majority of women were satisfied and would choose to self-insert a speculum again as it made the process less threatening (Wright *et al.*, 2005). These differences may be explained by the fact the women in this study were pregnant and in a 'high-risk' cohort. They may have fears of harming the pregnancy or their baby. As their gestation increases, women may also find it physically difficult to take their own vaginal swabs.

95% of women who responded found the transvaginal scan to measure their cervical length acceptable, while 2% of women did not. These results are similar to those in previous studies which

found that transvaginal ultrasound scans are acceptable to pregnant women (Atalabi *et al.*, 2012; Bello & Odeku, 2015; Clement *et al.*, 2003).

Limitations of this study include that no formal power calculation or advanced statistical analysis was performed on the data. Also as this was a convenience sample, there may be a possibility of selection bias. The women in this study were from an ethnically diverse background with a range of maternal ages and gestational ages. Caution must therefore be taken when applying these results to a different demographic of pregnant women.

Conclusions

Our findings suggest that women at risk of preterm birth find speculum examinations and transvaginal ultrasound scans for cervical length measurement acceptable. Any intimate examination should be approached with care, ensuring the patient is treated with dignity and respect and consent. While it is acknowledged that intimate medical examinations and procedures are not pleasant experiences, this study illustrates that the majority of pregnant women in a high-risk pregnancy are not opposed to them being performed.

More research is needed, however, to confirm this in a larger cohort using a formal power calculation and advanced statistical analysis. Further research could also focus on women's experience of speculum use in low-risk pregnancies (for example, for other swab tests), and if pregnant women would consider self-insertion of speculums.

References

- Abbott, D.S., Radford, S.K., Seed, P.T., Tribe, R.M. and Shennan, A.H. (2013) 'Evaluation of a quantitative fetal fibronectin test for spontaneous preterm birth in symptomatic women', *Am J Obstet Gynecol*, 208(122).e1–6.
- Abbott, D.S., Hezelgrave, N.L., Seed, P.T., Norman, J.E., David, A.L., Bennett, P.R., Girling, J.C., Chandirimani, M., Stock, S.J., Carter, J., Cate, R., Kurtzman, J., Tribe, R.M. and Shennan, A.H. (2015) 'Quantitative fetal fibronectin to predict preterm birth in asymptomatic women at high risk', *Obstetrics and Gynecology*, 125(5), 1168-1176.
- Atalabi, O. M., Morhason-Bello, I. O., Adekanmi, A. J., Marinho, A. O., Adedokun, B. O., Kalejaiye, A. O., Sogo, K. and Gbadamosi, S. A. (2012). 'Transvaginal ultrasonography: a survey of the acceptability and its predictors among a native African women population', *International Journal of Women's Health*, 4, 1–6.
- Bayrampour, H., Heaman, M., Duncan, K.A. & Tough, S. (2012). Predictors of Perception of Pregnancy Risk among Nulliparous Women. *Journal of Obstetric, Gynecologic & Neonatal Nursing*, 42(4): 416-427.
- Bello, F.A. and Odeku, A.O. (2015). 'Transvaginal sonography is feasible and universally acceptable to women in Ibadan, Nigeria: Experience from the 1st year of a novel service', *Annals of African Medicine*, 14(1), 52-56.
- Clement, S., Candy, B., Heath, V., To, M. and Nicolaides, K. H. (2003), 'Transvaginal ultrasound in pregnancy: its acceptability to women and maternal psychological morbidity'. *Ultrasound Obstet Gynecol*, 22: 508–514.
- Hoyo, C., Yarnall, K. S. H., Skinner, C. S., Moorman, P. G., Sellers, D. and Reid, L. (2005). 'Pain predicts non-adherence to Pap smear screening among middle-aged African American women'. *Preventive Medicine*, 41(2), 439–445.

Kurtzman, J., Chandiramani, M., Briley, A., Poston, L., Das, A. and Shennan, A. (2009) 'Quantitative fetal fibronectin screening in asymptomatic high-risk patients and the spectrum of risk for recurrent preterm delivery', *Am J Obstet Gynecol*, 200(263).e1–6.

Min, J., Watson, H. A., Hezelgrave, N. L., Seed, P. T. and Shennan, A. H. (2016). Ability of a preterm surveillance clinic to triage risk of preterm birth: a prospective cohort study. *Ultrasound Obstet Gynecol*, 48: 38–42.

Sharp, A. N., and Z. Alfrevic. (2014). "Provision and practice of specialist preterm labour clinics: a UK survey of practice." *BJOG: An International Journal of Obstetrics & Gynaecology*, 121(4): 417-421.

WHO (World Health Organization), 2012. *Born Too Soon: The Global Action Report on Preterm Birth*. Available at: http://www.who.int/pmnch/media/news/2012/preterm_birth_report/en/index5.html (Accessed 17/06/2017).

Yanikkerem, E., Özdemir, M., Bingol, H., Tatar, A. and Karadeniz, G. (2009). 'Women's attitudes and expectations regarding gynaecological examination'. *Midwifery*, 25(5), 500–508.