King's Research Portal

DOI:

[10.1097/RLU.0000000000001974](https://doi.org/10.1097/RLU.0000000000001974)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kim, B. J., Bradley, K. M., & Subesinghe, M. (2018). 18F-FDG PET/CT Detected Delayed Endoleak in an Aortoiliac Endovascular Aneurysm Repair. *Clinical Nuclear Medicine*, 43(3).
<https://doi.org/10.1097/RLU.0000000000001974>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Clinical Nuclear Medicine

F-18-FDG PET/CT detected delayed endoleak in an aorto-iliac endovascular aneurysm repair (EVAR) --Manuscript Draft--

Manuscript Number:	CNM-D-17-00916R1
Full Title:	F-18-FDG PET/CT detected delayed endoleak in an aorto-iliac endovascular aneurysm repair (EVAR)
Article Type:	Interesting Image
Keywords:	F-18-FDG PET/CT; EVAR; angiogram; endoleak; abdominal aortic aneurysm; diagnosis
Corresponding Author:	Byung Jin Kim, BA University of Oxford Oxford, Oxfordshire UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Oxford
Corresponding Author's Secondary Institution:	
First Author:	Byung Jin Kim, BA
First Author Secondary Information:	
Order of Authors:	Byung Jin Kim, BA
	Kevin M Bradley, FRCP, FRCR
	Manil Subesinghe, MRCP, FRCR
Order of Authors Secondary Information:	
Abstract:	<p>We present a case of a type Ia endoleak detected by F-18-FDG PET/CT 10 years following an abdominal aorto-iliac EVAR in an 83-year-old man. The F-18-FDG PET/CT scan was performed to evaluate a solitary pulmonary nodule but in addition demonstrated unexpected blood pool activity outside of the stent graft and within the abdominal aortic aneurysm (AAA) sac, indicating an endoleak; no FDG activity should be present within the aneurysm sac following stent graft placement. A subsequent CT angiogram confirmed the endoleak. This case illustrates the value of PET/CT in unexpectedly demonstrating this serious, late complication of EVAR.</p>

F-18-FDG PET/CT detected delayed endoleak in an aorto-iliac endovascular aneurysm repair (EVAR).

Authors

Byung Jin Kim, BA (Hons), Medical Sciences Division, John Radcliffe Hospital, OX3 9DU

Kevin M Bradley, FRCP, FRCR, Department of Radiology, Churchill Hospital, Oxford, OX3 7LE

Manil Subesinghe, MRCP, FRCR, Department of Radiology, Churchill Hospital, Oxford, OX3 7LE

Correspondence

Byung Jin Kim, BA (Hons), Medical Sciences Division, John Radcliffe Hospital, OX3 9DU

Email: byungjin.kim@ccc.ox.ac.uk

Short title

PET/CT detected delayed endoleak

Conflicts of Interest

None

F-18-FDG PET/CT detected delayed
endoleak in an aorto-iliac
endovascular aneurysm repair (EVAR).

Abstract

We present a case of a type Ia endoleak detected by F-18-FDG PET/CT 10 years following an abdominal aorto-iliac EVAR in an 83-year-old man. The F-18-FDG PET/CT scan was performed to evaluate a solitary pulmonary nodule but in addition demonstrated unexpected blood pool activity outside of the stent graft and within the abdominal aortic aneurysm (AAA) sac, indicating an endoleak; no FDG activity should be present within the aneurysm sac following stent graft placement. A subsequent CT angiogram confirmed the endoleak. This case illustrates the value of PET/CT in unexpectedly demonstrating this serious, late complication of EVAR.

Key words

F-18-FDG PET/CT, EVAR, angiogram, endoleak, abdominal aortic aneurysm, diagnosis

Figure Legend

An 83-year-old man underwent a whole-body F-18-Fludeoxyglucose (FDG) PET/CT scan to characterise/stage an enlarging 1.4cm pulmonary nodule. As well as demonstrating the pulmonary nodule, the PET/CT scan revealed blood pool activity extending for approximately 7 cm within the left side of the original large AAA sac (**A coronal, B axial; standard routine non-contrast PET images acquired at 90 minutes; SUV 0-6 scale; arrows to abnormal, extraluminal blood pool activity; C corresponding unenhanced axial CT image from PET/CT**), which had undergone an abdominal aorto-iliac EVAR ten years previously. The left iliac limb was known to be chronically occluded, and had been successfully treated by a femoral-femoral crossover graft. The abnormal activity was localised to within the aneurysm sac and outside of the EVAR stent graft. A CT angiogram was performed (**D and E**) and confirmed there was an active, type Ia endoleak.¹ The patient underwent open aortic neck banding but subsequently deceased due to pneumonia.

Endoleak is a unique complication of EVAR, caused by non-resolving pressure on the aneurysmal sac and it remains a significant problem in approximately 25% of patients, who have undergone EVAR to treat AAA² with most endoleaks occurring in the first 30 days post-EVAR^{3, 4}. A type 1 endoleak results from blood entering the aneurysm sac from an attachment site with consensus suggesting that this occurs in approximately 7.5% of procedures.⁵ Type 1a is from the proximal attachment site. Endoleak predisposes to post-operative morbidities, including aneurysmal sac expansion and subsequent rupture, making identification and treatment important.^{6, 7} Various methods of investigation exist to monitor for endoleaks, including magnetic resonance angiography and duplex ultrasound with or without contrast. However, the gold standard to detection remains multiphase CT angiography.⁷

Increased FDG uptake involving the aortic wall, resulting from inflammation, has been suggested to be a predictor for an endoleak and AAA rupture risk.⁸ Recently, early, dynamic PET/CT and PET angiography for endoleak detection after EVAR has been proposed⁹ although is only likely to find utility

if intravenous contrast is contraindicated. Given the increasing use of endovascular techniques for the repair of AAA, with their associated endoleaks, the recognition of delayed endoleaks, outside of routine CT surveillance programmes, on PET/CT scans performed for other indications is of critical importance.

References

1. Bashir MR, Ferral H, Jacobs C, et al. Endoleaks after endovascular abdominal aortic aneurysm repair: management strategies according to CT findings. *AJR Am J Roentgenol*. 2009;192:W178-186.
2. White SB, Stavropoulos SW. Management of Endoleaks following Endovascular Aneurysm Repair. *Semin Intervent Radiol*. 2009;26:33-38.
3. Shah A, Stavropoulos SW. Imaging Surveillance following Endovascular Aneurysm Repair. *Semin Intervent Radiol*. 2009;26:10-16.
4. Liaw JV, Clark M, Gibbs R, et al. Update: Complications and management of infrarenal EVAR. *Eur J Radiol*. 2009;71:541-551.
5. Veith FJ, Baum RA, Ohki T, et al. Nature and significance of endoleaks and endotension: summary of opinions expressed at an international conference. *J Vasc Surg*. 2002;35:1029-1035.
6. Tornqvist P, Resch T. Endoleaks after EVAR and TEVAR: indications for treatment and techniques. *J Cardiovasc Surg (Torino)*. 2014;55:105-114.
7. Avgerinos ED, Chaer RA, Makaroun MS. Type II endoleaks. *J Vasc Surg*. 2014;60:1386-1391.
8. Paraskevas KI, Tzovaras AA, Stathopoulos V, et al. Increased fluorodeoxyglucose uptake following endovascular abdominal aortic aneurysm repair: a predictor of endoleak? *Open Cardiovasc Med J*. 2010;4:117-119.
9. Drescher R, Guhne F, Freesmeyer M. Early-Dynamic Positron Emission Tomography (PET)/Computed Tomography and PET Angiography for Endoleak Detection After Endovascular Aneurysm Repair. *J Endovasc Ther*. 2017;24:421-424.