King's Research Portal

DOI:

[10.1016/j.puhe.2017.12.002](https://doi.org/10.1016/j.puhe.2017.12.002)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Vancampfort, D., Lara, E., Stubbs, B., Swinnen, N., Probst, M., & Koyanagi, A. (2018). Physical activity correlates in people with mild cognitive impairment: findings from six low- and middle-income countries. *Public Health*, 156, 15-25. <https://doi.org/10.1016/j.puhe.2017.12.002>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Abstract

Objectives: Despite promising research showing that physical activity (PA) might improve cognitive functioning in people with mild cognitive impairment (MCI), people with MCI are less physically active compared to the general population. Therefore, the aim of this study was to assess PA correlates among community-dwelling older people with MCI in six low- and middle-income countries.

Design: Cross-sectional data from the World Health Organization's Study on Global Ageing and Adult Health were analysed.

Methods: PA level was assessed by the Global Physical Activity Questionnaire. 4,854 participants with MCI (mean age 64.4 years; 55.1% females) were grouped into those who do and do not (low PA) meet the 150 minutes of moderate-to-vigorous PA per week recommendation. Associations between PA and the correlates were examined using multivariable logistic regressions.

Results: The prevalence of low PA was 27.4% (95%CI=25.0-30.0). In the multivariable analysis, older age and unemployment were the only sociodemographic correlates of low PA. The significant positive correlates of low PA in other domains included depression, being underweight, obesity, asthma, chronic lung disease, hearing problems, visual impairment, slow gait, weak grip strength, poor self-rated health, and lower levels of social cohesion.

Conclusions: The current data illustrate that a number of sociodemographic and health factors are associated with PA levels among older people with MCI. The promotion of social cohesion may increase the efficacy of public health initiatives while from a health care perspective, somatic co-morbidities, muscle strength and slow gait need to be considered when activating those at risk for dementia.

Keywords: physical activity; exercise; cognition; exercise

Introduction

With an advancing aging population and associated worldwide rise in dementia prevalence, the associated costs and disease burden have exerted significant pressure on global economic and social systems¹. Currently, it is estimated that approximately 50 million people worldwide are diagnosed with dementia, this figure is expected to increase to 132 million by 2050². There is increasing recognition that due to an upward trend in physical inactivity and non-communicable diseases in low- and middle-income countries (LMICs)³, both risk factors for developing dementia^{4,5}, this part of the world is likely to experience the largest burden of dementia. Currently 58% of the world population having dementia lives in LMICs, but this percentage is estimated to increase to 68% in 2050⁶.

There are at the moment no proven treatments available to cure dementia or to alter its clinical course^{7,8}. Therefore, primary prevention and public health initiatives in the precursory stages of dementia is essential. Mild cognitive impairment (MCI) is considered to be a preclinical state of dementia⁹. Identifying modifiable risk factors for dementia in people with MCI is important in order to establish public and clinical health initiatives that might prevent or delay the onset of dementia. It has been demonstrated that low physical activity (PA) is such a risk factor with population attributable risk for dementia being 21.0% (95%CI=5.8-36.6), 20.3% (95%CI=5.6-35.6), and 21.8% (95%CI=6.1-37.7) in the USA, Europe and UK, respectively¹⁰. It is estimated that regular PA could prevent almost 300,000 cases of dementia per year, worldwide, if everyone were to comply with the public health recommendation of 150 minutes of moderate-to-vigorous PA per week¹¹. There are likely cellular (e.g. neurogenesis, synaptogenesis, neuro-angiogenesis) and humoral (e.g. neurotrophic factors, inflammatory cytokines) mechanisms and behavioural and socioemotional pathways (e.g., stress, sleep, depression, pain)¹² that explain PA effects on the brain and cognition.

Despite promising research showing that PA might improve cognitive functioning in people with MCI¹³⁻¹⁹, it is concerning that people with MCI are, compared to the general population, at a higher risk of not meeting the recommended 150 min/week moderate-to-vigorous intensity PA level (odds ratio=1.28; 95%CI=1.11-1.48)²⁰. To date, there is no research investigating factors that influence PA participation in this population.

Understanding the correlates of PA in MCI is important for multiple health reasons, including the enhancement of executive function²¹ for the maintenance of personal autonomy²² and reducing caregiver burden²³. In addition to the overall lack of information on factors influencing PA participation in people with MCI, very little is known about PA correlates in older people in LMICs in general. This is important as these countries are characterized by a suboptimal treatment of neurocognitive disorders and mental health problems²⁴, and often a lack of

knowledge regarding the benefits of PA ^{25, 26}. Information on PA correlates of older people with MCI in LMICs could guide the design and delivery of targeted public health interventions in these countries. It could also provide useful information that can support integration of PA into primary health care settings in many LMICs.

Thus, given the aforementioned gaps within the literature, we aimed to assess PA correlates among community-dwelling older people with MCI in six LMICs.

Methods

The survey

Data from the Study on Global Ageing and Adult Health (SAGE) survey was analysed. This dataset is publically available through the World Health Organization (WHO) website (<http://www.who.int/healthinfo/sage/en/>). The survey was undertaken in China, Ghana, India, Mexico, Russia, and South Africa between 2007 and 2010. These countries were all LMICs according to the World Bank classification at the time of the survey. Details of the survey methodology have been published elsewhere ²⁷. Following a common research protocol across countries, trained interviewers conducted face-to-face interviews using a standard questionnaire to collect information. Standard translation procedures for the questionnaires were undertaken to ensure comparability between countries. A stadiometer and a routinely calibrated electronic weighing scale were used to measure height and weight respectively. Blood pressure was measured three times with a one-minute interval with the use of a wrist blood pressure monitor. Grip strength was measured twice for both hands with the use of the Smedley's hand dynamometer. If the participant had any surgery in the last three months or arthritis or pain in the hand/wrist/arm, grip strength was not measured for that hand. Gait speed was based on a 4-m timed walk and was measured by asking the participant to walk at a usual pace. A cane or other walking aids were allowed. The interviewer recorded the time to completion of the 4-m walk.

Respondents who were unable to undertake the interview because of limited cognitive function were not included in the analysis. The survey response rate ranged from 51% (Mexico) to 93% (China). Sampling weights were constructed to adjust for the population structure as reported by the United Nations Statistical Division. Ethical approval was obtained from the WHO Ethical Review Committee and local ethics research review boards. Written informed consent was obtained from all participants.

Assessments

Mild cognitive impairment

MCI was ascertained based on the recommendations of the National Institute on Aging-Alzheimer's Association ²⁸. We applied the identical algorithms used in previous publications using a dataset with the same survey questions to identify MCI ^{29, 30}. Briefly, individuals fulfilling all of the following conditions were considered to have MCI:

(a) Concern about a change in cognition: Individuals who replied 'bad' or 'very bad' to the question "How would you best describe your memory at present?" and/or those who answered 'worse' to the question "Compared to

12 months ago, would you say your memory is now better, the same or worse than it was then?" were considered to have this condition.

(b) Objective evidence of impairment in one or more cognitive domains: was based on a <-1 SD cut-off after adjustment for level of education (completed secondary or less), age, and country. Cognitive function was assessed through the following performance tests: word list immediate and delayed verbal recall from the Consortium to Establish a Registry for Alzheimer's Disease ³¹, which assessed learning and episodic memory; digit span forward and backwards from the Wechsler Adult Intelligence Scale ³², that evaluated attention and working memory; and the animal naming task³¹, which assessed verbal fluency.

(c) Preservation of independence in functional abilities: was assessed by questions on self-reported difficulties with basic activities of daily living (ADL) in the past 30 days ³³. Specific questions were: "How much difficulty did you have in getting dressed?" and "How much difficulty did you have with eating (including cutting up your food)?". If the individual reported that ADL was intact or minimally impaired based on both questions, their ADL was considered to be preserved.

(d) No dementia: Individuals with a level of cognitive impairment severe enough to preclude the possibility to undertake the survey were not included in the current study.

Physical activity (PA)

Level of PA was assessed with the Global Physical Activity Questionnaire ³⁴. The total amount of moderate to vigorous PA in a typical week was calculated based on self-report. Those scoring ≥ 150 minutes of moderate to high intensity PA were classified as meeting the recommended guidelines (coded=0), and those scoring < 150 minutes (low PA) were classified as not meeting the recommended guidelines (coded=1) ³⁵.

Sociodemographic variables

These included age, sex, highest level of education achieved (completed secondary or less), wealth, marital status (married/cohabiting or else), household size (1, 2, ≥ 3), setting (urban or rural), and employment status (engaged in paid work ≥ 2 days in last 7 days: Yes/No). Wealth quintiles were created based on country-specific income.

Health behaviour

These comprised of current drinking [alcohol use in the past 30 days (Yes/No)], fruit and vegetable intake [≥ 2 (fruits) and ≥ 3 (vegetables) servings per day (adequate)]³⁶, and smoking (never, current, quit)³⁷.

Mental health

Anxiety was defined as having severe or extreme problems with worry or anxiety in the last 30 days³⁸. Questions based on the World Mental Health Survey version of the Composite International Diagnostic Interview³⁹ were used for the endorsement of past 12-month DSM-IV depression⁴⁰ (Details provided in **eTable 1**). Those having severe or extreme problems with sleeping, such as falling asleep, waking up frequently during the night or waking up too early in the morning, were considered to have sleep problems⁴¹.

Physical health

Body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared, and was categorized as <18.5 (underweight), 18.5-24.9 (normal), 25.0-29.9 (overweight), and ≥ 30 (obese) kg/m². Participants who had severe or extreme bodily aches or pains in the previous 30 days were considered to have bodily pain⁴². Chronic back pain was defined as having had back pain every day during the previous 30 days⁴³. Fall-related injuries in the previous 12 months were assessed with questions on the presence of bodily injury and cause⁴⁴. The participant was considered to have hearing problems if the interviewer observed this condition. Visual impairment was defined as having extreme difficulty in seeing and recognizing a person that the participant knows across the road⁴⁵.

Diabetes and stroke were solely based on lifetime self-reported diagnosis. Hypertension was defined as having at least one of: systolic blood pressure ≥ 140 mmHg; diastolic blood pressure ≥ 90 mmHg; or self-reported diagnosis. For angina, arthritis, asthma, and chronic lung disease, the participant was considered to have the condition in the presence of self-reported diagnosis and/or symptom-based diagnosis using algorithms. Specifically, the validated Rose questionnaire was used for angina⁴⁶, and other previously validated symptom-based algorithms were used for arthritis, asthma, and chronic lung disease^{47, 48} (See **eTable 2**).

Physical performance

Slow gait was defined as a gait speed of ≤ 1 m/sec⁴⁹. Weak handgrip was defined as <30 kg for men and <20 kg for women using the average value of the two handgrip measurements of the dominant hand⁵⁰.

Health status

Self-rated health was evaluated by the question 'In general, how would you rate your health today?' Those who answered 'bad' or 'very bad' were considered to have poor self-rated health. Self-rated health has been reported to be a valid indicator of overall health and a strong predictor of outcomes such as mortality⁵¹.

Social cohesion

The social cohesion index was built with the modified version of Ramlagan, Peltzer, and Phaswana-Mafuya (2013)⁵². It was created based on 9 questions on the participant's involvement in community activities in the past 12 months (e.g., attended religious services, club, society, union etc.) with answer options 'never (coded=1)', 'once or twice per year (coded=2)', 'once or twice per month (coded=3)', 'once or twice per week (coded=4)', and 'daily (coded=5)' (Actual questions are shown in **eTable 3**). The answers to these questions were summed (range=9-45) and later converted to a scale ranging from 0-100 with higher scores corresponding to higher levels of social cohesion (Cronbach's $\alpha=0.78$).

Statistical analysis

The statistical analysis was performed with Stata 14.1 (Stata Corp LP, College station, Texas). The analysis was restricted to individuals with MCI aged ≥ 50 years. The difference in sample characteristics between those with and without low PA was tested by Chi-squared tests and Student's *t*-tests for categorical and continuous variables respectively. The selection of the correlates of PA was based on past literature⁵³. We conducted multivariable logistic regression analyses using the overall and country-wise samples. First, we assessed the sociodemographic correlates of PA by constructing a model which includes all the sociodemographic variables (age, sex, education, wealth, marital status, household size, setting, employment status). Next, we assessed the association between each of the other health-related correlates with low PA while adjusting for all the sociodemographic variables mentioned above. We also conducted a sensitivity analysis to assess whether the correlates of MCI are similar among those with and without MCI using the overall sample including all countries. We included an interaction term of the correlate in question and MCI to test whether the association between the correlate and low PA differs by the presence or absence of MCI. The regression analyses using the overall sample were adjusted for country by including dummy variables for each country. All variables were included in the models as categorical variables with the exception of age and social cohesion (continuous variables). The sample weighting and the complex study design were taken into account in all analyses. Results from the

regression analyses are presented as odds ratios (ORs) with 95% confidence intervals (CIs). The level of statistical significance was set at $P < 0.05$.

Results

The prevalence of MCI in the general population aged ≥ 50 years was 14.8% (China 24.1%; Ghana 7.3%; India 9.1%; Mexico 16.9%; Russia 9.3%; South Africa 8.2%). A total of 4,854 individuals aged ≥ 50 years with MCI constituted the final sample (China $n=2,802$; Ghana $n=307$; India $n=646$; Mexico $n=329$; Russia $n=473$; South Africa $n=297$). The mean (SD) age of the sample was 64.4 (17.0) years and 55.1% were females. The prevalence of low PA was 27.4% (95%CI=25.0-30.0). In terms of the sociodemographic correlates, in the unadjusted analysis, individuals with low PA were significantly older, were part of a larger household, were more likely to live in urban areas and were unemployed, while they were also less likely to be married/cohabiting (**Table 1**). In terms of other correlates, those not meeting PA guidelines had a higher rate of non-alcohol consumption, inadequate fruit and vegetable consumption, non-smoking, depression, sleep problems, were underweight, had a high BMI, bodily pain, angina, asthma, chronic lung disease, hearing problems, stroke, visual impairment, slow gait, weak grip strength, poor self-rated health, and low levels of social cohesion ($P < 0.05$) (**Table 2**). In the multivariable analysis, the only sociodemographic correlates significantly associated with low PA were older age and unemployment (**Table 3**). The significant positive correlates of low PA in other domains based on adjusted estimates included depression, underweight, obesity, asthma, chronic lung disease, hearing problems, visual impairment, slow gait, weak grip strength, poor self-rated health, and lower levels of social cohesion (**Table 4**). Alcohol consumption and smoking were negatively associated with low PA ($P < 0.05$). There were some differences between the countries in terms of the correlates of low PA among those with MCI (see **eTables 4 and 5**). For example, regarding socio-demographic correlates, only Ghanaian men with MCI had higher odds for low PA. People in China who completed their secondary education had higher odds for low PA, but in Ghana, an opposite trend was observed. Wealthier people had higher odds for not complying with the PA guidelines only in Russia. As for health-related factors, some were significant correlates only in selected countries. For example, depression and chronic lung disease were significant positive correlates of low PA only in China and India, low BMI and visual impairment only in India, arthritis and asthma only in China, fall-related injuries only in Ghana, sleep problems only in Mexico, and a weak grip strength only in India and South Africa. Finally, we also assessed whether the correlates of PA differ among those with and without MCI. The results showed that the correlates were similar overall but with some differences. See **eTables 6 and 7** published as

supplementary material online attached to the electronic version of this paper. Older age, being underweight, fall-related injury, slow gait, and the presence of depression were significantly more strongly associated with not meeting PA guidelines among those with MCI (significant interaction $P < 0.05$).

Discussion

General findings

To the best of our knowledge, the current study is the first multinational study to explore PA correlates in older people with MCI. Based on the adjusted analysis, the only sociodemographic correlates significantly associated with low PA (i.e. not complying with the recommendation of 150 min/week moderate-to-vigorous intensity PA) were older age and being unemployed. Significant correlates of low PA in other domains included lower levels of social cohesion, the presence of depression, underweight or obesity, asthma, chronic lung disease, hearing problems, visual impairment, slow gait, weak grip strength, and poor self-rated health. At the behavioural level, alcohol consumption and smoking were negatively associated with low PA. Compared to those without MCI, older age, being underweight, fall-related injury, slow gait, and the presence of depression were stronger correlates for not meeting PA guidelines among those with MCI.

We found that participants who were employed were more physically active which is perhaps unsurprising. Nonetheless, the current study adds further evidence to a recent review demonstrating that there are important health benefits of continuing to engage in meaningful occupation in people with MCI⁵⁴. Next to the physically demanding labour, active transport to and from work might be an underlying reason for the more active lifestyle of those who work. Speculatively, employment may offer opportunities for older people to connect socially, enhance social functioning and consequently have more opportunity to be physically active⁵⁵⁻⁵⁷. Less social cohesion was in our study related to less likelihood of complying with the PA recommendation. Social cohesion and compliance with PA guidelines may be linked to each other in various ways⁵⁸. At the neighbourhood level, a high level of social cohesion is associated with less crime, and low neighbourhood crime rates tend to be associated with greater PA engagement⁵⁹. Also, socially cohesive neighbourhoods may be more likely to organize community activities that present opportunities for older people to engage in physical activities⁶⁰. At the individual level, it is important to feel socially connected to the community because it may increase the likelihood that one will take advantage of local opportunities to engage in PA.

In our multivariable analysis increasing age was another important socio-demographic correlate of PA participation in people MCI, which is in line with the wider literature. Therefore, in preventive public health

campaigns and rehabilitation programs special attention and, if required, additional support should be given to people at the oldest ages. A possible underlying reason is that those people are at an even higher risk for chronic conditions and worse cognition scores. In terms of chronic physical conditions most previous studies in people with MCI have focused on cardio-metabolic diseases ^{61, 62}, partly motivated by the theoretical link between cognitive decline via atherosclerosis, microvascular changes, and inflammatory processes ⁶³. The current study shows that in motivating people with MCI towards an active lifestyle, the presence of chronic lung diseases should be considered as a barrier. Vice versa, in people with chronic lung diseases MCI should be taken into account. For example, one in four people with COPD has MCI ⁶⁴ and the presence of MCI in this population is associated with a higher risk for developing depression, which is an important barrier for complying with PA prescriptions in this population ⁶⁵. Also, older people with asthma have a higher risk for MCI ⁶⁶. The presence of hearing problems ⁶⁷ and visual impairments ⁶⁸ should, on their turn, be considered as important barriers for being physically active due to perceived fear or risk of injury in old people in LMICs. Stigma and discrimination associated with it may further complicate PA participation in older people in these countries ⁶⁹.

The presence of chronic co-morbidities also results in a higher likelihood of experiencing pain. A recent study showed that mild to moderate pain and MCI are independently associated with poor mobility, and the presence of both was associated with the poorest status ⁷⁰. Therefore, health care professionals who encounter older adults in need of mobility rehabilitation should consider screening them for pain and MCI to better inform subsequent therapeutic interventions. Collaborative interventions that seek to reduce chronic conditions and pain and improve mobility among those who present with MCI might on its turn be an important strategy to increase PA levels.

Next to this, the present study demonstrated that a slow gait speed and low grip strength are associated with lower PA in older people with MCI. These factors may be reflecting the presence of a chronic physical illness, and the difficulty to engage in PA due to these conditions. Alternatively, the observation that a low grip strength and a slow gait, but also a low BMI was associated with less PA might be due to the fact that these are measures-of-proxy of frailty and sarcopenia (pathological loss of skeletal muscle mass and function) ^{71, 72}. People with MCI have a high risk for sarcopenia ⁷³, while the presence of sarcopenia is also associated with a higher risk for developing MCI ⁷⁴. A number of recent studies have demonstrated that slow gait speed is a strong predictor of future incident cognitive decline and mortality ^{75, 76}. Experiencing difficulty undertaking a physical performance measure assessing gait speed may therefore be an important sign that an older person is at risk of MCI and also physically inactive. Next to this, in people with MCI a worse performance on handgrip strength

and gait speed tests is associated with a higher risk of falling ⁷⁷. Risk of falls in older people is associated with a fear of falling again and avoidance of PA and increased sedentary behaviour ⁷⁸.

Next, our data demonstrate that depression should be considered as an important correlate of low PA in people with MCI. Depression is very common in people with MCI with a recent meta-analysis including 57 studies and 20 892 patients showing an overall pooled prevalence of 32% (95%CI=27%-37%) ⁷⁹. Depression is well-known barrier for complying with international PA standards. ⁸⁰. Given the evidence that depression can lead also to (further) cognitive decline ⁸¹, being physically active, despite the lack of energy and motivation and a low self-efficacy, may reduce the depressive symptoms in older people ⁸² and consequently enhance cognitive functioning. It is possible that PA has an indirect effect on cognitive functioning through its effect on older adults' mental health. Longitudinal mediation models are however needed to further untangle the complex relationship between PA, depression, and cognition.

Finally, while we identified some variation in the correlates across the different countries, the precise reasons for such differences and inconsistencies are not clear. For example, increasing levels of wealth were only associated with lower odds for low PA in Russia but not in other countries. One potential reason might be that in Russia, increasing levels of wealth correspond more strongly to less labor-demanding jobs, which means less occupational PA. The inconsistent patterns of sociodemographic PA correlates points to apparently contrasting socio-economic patterns of risk factors for low PA. A possible reason for the differences in health-related factors may be due to differences in access to health care and awareness about prevention and control of risk factors for low PA. More research is however needed to clarify these differences, which should inform country specific interventions.

Strengths, limitations and future research

The strength of the study includes the large sample size and the use of nationally representative samples from six countries, which comprise nearly half of the worldwide population ²⁷. Nevertheless, the study results should be interpreted in the light of several limitations. First, PA was measured with a self-report questionnaire, which is known to be less accurate than objective assessments ^{83, 84} and can overestimate PA levels ⁸⁵. Second, we did not assess cognitive deficits beyond attention/working memory, learning/episodic memory and language dysfunction which might have resulted in an underestimation of the number of people with MCI. However, it is reassuring that the prevalence of MCI in our study was within previously reported figures.⁸⁶ Third, because this was a cross-sectional study, causality cannot be inferred. Fourth, we were not able to explore differences in PA

participation in people with MCI living in urban versus rural settings in more detail. Future research could explore the role of specific environmental attributes relevant to PA on mental health populations (including those who present with MCI) in LMICs such as the availability and quality of sidewalks, pedestrian zones, bicycle facilities, and factors affecting intersection quality (e.g., crosswalks, pedestrian signals). In addition, future studies would benefit from assessing to what extent macro-level environmental factors such as food insecurity, civil conflicts, and extreme weather in LMICs are linked to physical inactivity in the older population in these countries.

Health policy-related implications

Despite the limitations, our data provides some guidance for health policy makers. First of all, employment and or volunteering work, at least when of high quality (i.e. without stressful or adverse psychosocial work conditions)⁸⁷ may serve as an opportunity for older people to engage in meaningful PA. Thus, public interventions seeking to improve the likelihood that older people with MCI can be involved in high quality employment or volunteering work may serve as an opportunity for older people to have a valuable role in society as well as achieve recommended PA targets. The current results also suggest that also in LMICs, among older adults health campaigns should consider somatic co-morbidities, reduced muscle strength, slow gait and depression as these might be important barriers for PA. On the other hand, in older adults with somatic co-morbidities, pain, frailty or sarcopenia, cognitive impairments should be taken into account as potential barriers for a more active lifestyle. However, increasing occupational PA may not be a feasible intervention target when a shift occurs from manual jobs to service-based industries in more urbanized areas. Therefore policy makers should focus on other identified PA correlates as well. For example, our study shows that the public health policy in LMICs should focus on social cohesion in order to facilitate older people to become more active. This can be done in several ways. In neighbourhoods where only a few older residents feel that they belong to the community, PA could be increased by promoting social engagement and participation among community members by organizing social and PA activities targeting older people. In neighbourhoods where the overall level of social cohesion is already high, further benefits may be achieved by socially integrating older residents who do not already feel that they belong.

In conclusion, our data illustrate that a number of sociodemographic, physical and mental health and behavioural factors are associated with PA levels among older people with MCI in six LMICs. These findings provide guidance for future public health interventions across LMICs to help older people with MCI, who are at risk for physical inactivity, to become more active.

Acknowledgements

None.

References

1. Wimo A, Guerchet M, Ali GC, Wu YT, Prina AM, Winblad B, et al. The worldwide costs of dementia 2015 and comparisons with 2010. *Alzheimer's & Dementia*. 2017; 13:1-7.
2. Prince MJ. World Alzheimer Report 2015: the global impact of dementia: an analysis of prevalence, incidence, cost and trends: Alzheimer's Disease International; 2015.
3. Christensen K, Doblhammer G, Rau R, Vaupel JW. Ageing populations: the challenges ahead. *Lancet*. 2009; 374:1196-208.
4. Beydoun MA, Beydoun H, Wang Y. Obesity and central obesity as risk factors for incident dementia and its subtypes: a systematic review and meta-analysis. *Obesity Reviews*. 2008; 9:204-18.
5. Covell GES, Hoffman-Snyder CR, Wellik KE, Woodruff BK, Geda YE, Caselli RJ, et al. Physical activity level and future risk of mild cognitive impairment or dementia: a critically appraised topic. *The Neurologist*. 2015; 19:89-91.
6. Alzheimer's Disease International. World Alzheimer Report 2015.
7. Cummings JL. Treatment of Alzheimer's disease: current and future therapeutic approaches. *Reviews in Neurological Diseases*. 2004; 1:60-9.
8. Kaduszkiewicz H, Zimmermann T, Beck-Bornholdt HP, van den Bussche H. Cholinesterase inhibitors for patients with Alzheimer's disease: systematic review of randomised clinical trials. *BMJ*. 2005; 331:321-7.
9. Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia*. 2011; 7:270-9.
10. Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C. Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. *The Lancet Neurology*. 2014; 13:788-94.
11. Sallis JF, Bull F, Guthold R, Heath GW, Inoue S, Kelly P, et al. Progress in physical activity over the Olympic quadrennium. *The Lancet*. 2016; 388:1325-36.
12. Stillman CM, Cohen J, Lehman ME, Erickson KI. Mediators of physical activity on neurocognitive function: a review at multiple levels of analysis. *Frontiers in Human Neuroscience*. 2016; 10.

13. Ahlskog JE, Geda YE, Graff-Radford NR, Petersen RC, editors. Physical exercise as a preventive or disease-modifying treatment of dementia and brain aging. Mayo Clinic Proceedings; 2011: Elsevier.
14. Cammisuli D, Innocenti A, Franzoni F, Pruneti C. Aerobic exercise effects upon cognition in Mild Cognitive Impairment: A systematic review of randomized controlled trials. Archives Italiennes de Biologie. 2017; 155:55-62.
15. Barber SE, Clegg AP, Young JB. Is there a role for physical activity in preventing cognitive decline in people with mild cognitive impairment? Age and Ageing. 2011; 41:5-8.
16. Blondell SJ, Hammersley-Mather R, Veerman JL. Does physical activity prevent cognitive decline and dementia?: A systematic review and meta-analysis of longitudinal studies. BMC Public Health. 2014; 14:510.
17. Lautenschlager NT, Cox K, Kurz AF. Physical activity and mild cognitive impairment and Alzheimer's disease. Current Neurology and Neuroscience Reports. 2010; 10:352-8.
18. Sofi F, Valecchi D, Bacci D, Abbate R, Gensini GF, Casini A, et al. Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. Journal of Internal Medicine. 2011; 269:107-17.
19. Gallaway PJ, Miyake H, Buchowski MS, Shimada M, Yoshitake Y, Kim AS, et al. Physical activity: a viable way to reduce the risks of mild cognitive impairment, Alzheimer's disease, and vascular dementia in older adults. Brain Sciences. 2017; 7:22.
20. Vancampfort D, Stubbs B, Lara E, Vandenbulcke M, Swinnen N. Mild cognitive impairment and physical activity in the general population: findings from six low- and middle-income countries. Experimental Gerontology. 2017; 100: 100-105.
21. Kobayashi Y, Takahashi Y, Seki T, Kaneta T, Amarume K, Kasai M, et al. Decreased Physical Activity Associated with Executive Dysfunction Correlates with Cognitive Impairment among Older Adults in the Community: A Retrospective Analysis from the Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra. 2016; 6: 350-60.
22. Tortosa-Martínez J, Beltrán-Carrillo VJ, Caus N, Iglesias-Martínez MJ, Lozano-Cabezas I, Jimenez-Hernández S, et al. Psychosocial benefits of exercise for older adults with amnesic Mild Cognitive Impairment: Innovative practice. Dementia. 2017: 1471301217725895.
23. Holthoff VA, Marschner K, Scharf M, Steding J, Meyer S, Koch R, et al. Effects of physical activity training in patients with Alzheimer's dementia: results of a pilot RCT study. PloS One. 2015; 10: e0121478.
24. Patel V, Araya R, Chatterjee S, Chisholm D, Cohen A, De Silva M, et al. Treatment and prevention of mental disorders in low-income and middle-income countries. The Lancet. 2007; 370:991-1005.

25. Pengpid S, Peltzer K, Kassean HK, Tsala JPT, Sychareun V, Müller-Riemenschneider F. Physical inactivity and associated factors among university students in 23 low-, middle-and high-income countries. *International Journal of Public Health*. 2015; 60:539-49.
26. Vancampfort D, Stubbs B, De Hert M, du Plessis C, Gbiri CAO, Kibet J, et al. A systematic review of physical activity policy recommendations and interventions for people with mental health problems in Sub-Saharan African countries. *Pan African Medical Journal*. 2017; 26.
27. Kowal P, Chatterji S, Naidoo N, Biritwum R, Fan W, Lopez Ridaura R, et al. Data resource profile: the World Health Organization Study on global AGEing and adult health (SAGE). *International Journal of Epidemiology*. 2012; 41:1639-49.
28. Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia*. 2011; 7:270-9.
29. Lara E, Koyanagi A, Domenech-Abella J, Miret M, Ayuso-Mateos JL, Haro JM. The Impact of depression on the development of mild cognitive impairment over 3 years of follow-up: a population-based study. *Dementia and Geriatric Cognitive Disorders*. 2017; 43:155-69.
30. Lara E, Koyanagi A, Olaya B, Lobo A, Miret M, Tyrovolas S, et al. Mild cognitive impairment in a Spanish representative sample: prevalence and associated factors. *International journal of Geriatric Psychiatry*. 2016; 31:858-67.
31. Morris JC, Heyman A, Mohs RC, Hughes JP, van Belle G, Fillenbaum G, et al. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology*. 1989; 39:1159-65.
32. The Psychological Corporation. The Psychological Corporation: The WAIS III-WMS III Updated Technical Manual. San Antonio 2002.
33. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of illness in the aged. The index of ADL: A standardized measure of biological and psychosocial function. *JAMA*. 1963; 185:914-9.
34. Bull FC, Maslin TS, Armstrong T. Global physical activity questionnaire (GPAQ): nine country reliability and validity study. *Journal of Physical Activity & Health*. 2009; 6:790-804.
35. World Health Organization. Global recommendations on Physical Activity for health: World Health Organization; 2010.

36. Joint FAO/WHO Expert Consultation. Joint WHO/FAO expert consultation on diet, nutrition and the prevention of chronic diseases. Geneva 2003.
37. Koyanagi A, Stickley A, Garin N, Miret M, Ayuso-Mateos JL, Leonardi M, et al. The association between obesity and back pain in nine countries: a cross-sectional study. *BMC Public Health*. 2015; 15:123.
38. Stubbs B, Koyanagi A, Hallgren M, Firth J, Richards J, Schuch F, et al. Physical activity and anxiety: A perspective from the World Health Survey. *Journal of affective disorders*. 2017; 208:545-52.
39. Kessler RC, Ustun TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int J Methods Psychiatry Research*. 2004; 13:93-121.
40. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (4th ed.) 2000.
41. Koyanagi A, Stickley A. The Association between Sleep Problems and Psychotic Symptoms in the General Population: A Global Perspective. *Sleep*. 2015; 38:1875-85.
42. Koyanagi A, Stickley A. The association between psychosis and severe pain in community-dwelling adults: Findings from 44 low- and middle-income countries. *Journal of Psychiatric Research*. 2015; 69:19-26.
43. Stubbs B, Koyanagi A, Thompson T, Veronese N, Carvalho AF, Solomi M, et al. The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low- and middle-income countries. *General Hospital Psychiatry*. 2016; 43:63-70.
44. Stewart Williams J, Kowal P, Hestekin H, O'Driscoll T, Peltzer K, Yawson A, et al. Prevalence, risk factors and disability associated with fall-related injury in older adults in low- and middle-income countries: results from the WHO Study on global AGEing and adult health (SAGE). *BMC Medicine*. 2015; 13:147.
45. Freeman EE, Roy-Gagnon MH, Samson E, Haddad S, Aubin MJ, Vela C, et al. The global burden of visual difficulty in low, middle, and high income countries. *PloS one*. 2013; 8:e63315.
46. Rose GA. The diagnosis of ischaemic heart pain and intermittent claudication in field surveys. *Bulletin of the World Health Organization*. 1962; 27:645-58.
47. Arokiasamy P, Uttamacharya, Kowal P, Capistrant BD, Gildner TE, Thiele E, et al. Chronic Noncommunicable Diseases in 6 Low- and Middle-Income Countries: Findings From Wave 1 of the World Health Organization's Study on Global Ageing and Adult Health (SAGE). *American Journal of Epidemiology*. 2017; 185:414-28.
48. Moussavi S, Chatterji S, Verdes E, Tandon A, Patel V, Ustun B. Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet*. 2007; 370:851-8.

49. Abellan van Kan G, Rolland Y, Andrieu S, Bauer J, Beauchet O, Bonnefoy M, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people an International Academy on Nutrition and Aging (IANA) Task Force. *The Journal of Nutrition, Health & Aging*. 2009; 13:881-9.
50. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age and Ageing*. 2010; 39:412-23.
51. Schnittker J, Bacak V. The increasing predictive validity of self-rated health. *PloS one*. 2014; 9:e84933.
52. Ramlagan S, Peltzer K, Phaswana-Mafuya N. Social capital and health among older adults in South Africa. *BMC geriatrics*. 2013; 13:100.
53. Stubbs B, Eggermont L, Soundy A, Probst M, Vandenbulcke M, Vancampfort D. What are the factors associated with physical activity (PA) participation in community dwelling adults with dementia? A systematic review of PA correlates. *Archives of Gerontology and Geriatrics*. 2014; 59:195-203.
54. McCulloch S, Robertson D, Kirkpatrick P. Sustaining people with dementia or mild cognitive impairment in employment: A systematic review of qualitative evidence. *British Journal of Occupational Therapy*. 2016; 79:682-92.
55. Suetani S, Waterreus A, Morgan V, Foley DL, Galletly C, Badcock JC, et al. Correlates of physical activity in people living with psychotic illness. *Acta Psychiatrica Scandinavica*. 2016; 134:129-37.
55. Sarkar S, Taylor WC, Lai D, Shegog R, Paxton RJ. Social support for physical activity: Comparison of family, friends, and coworkers. *Work*. 2016; 55:893-9.
57. Böhm AW, Mielke GI, da Cruz MF, Ramires VV, Wehrmeister FC. Social Support and Leisure-Time Physical Activity Among the Elderly: A Population-Based Study. *Journal of Physical Activity and Health*. 2016; 13:599-605.
58. Daskalopoulou C, Stubbs B, Kralj C, Koukounari A, Prince M, Prina A. Physical activity and healthy ageing: a systematic review and meta-analysis of longitudinal cohort studies. *Ageing Research Reviews*. 2017.
59. Sampson RJ, Raudenbush SW, Earls F. Neighborhoods and violent crime: A multilevel study of collective efficacy. *Science*. 1997; 277:918-24.
60. Cohen DA, Finch BK, Bower A, Sastry N. Collective efficacy and obesity: the potential influence of social factors on health. *Social Science & Medicine*. 2006; 62:769-78.

61. Ng TP, Feng L, Nyunt MSZ, Feng L, Gao Q, Lim ML, et al. Metabolic syndrome and the risk of mild cognitive impairment and progression to dementia: follow-up of the Singapore Longitudinal Ageing Study Cohort. *JAMA Neurology*. 2016; 73:456-63.
62. DeCarlo CA, MacDonald SW, Vergote D, Jhamandas J, Westaway D, Dixon RA. Vascular health and genetic risk affect mild cognitive impairment status and 4-year stability: Evidence from the Victoria Longitudinal Study. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2015; 71:1004-14.
63. Biessels GJ, Staekenborg S, Brunner E, Brayne C, Scheltens P. Risk of dementia in diabetes mellitus: a systematic review. *The Lancet Neurology*. 2006; 5:64-74.
64. Yohannes AM, Chen W, Moga AM, Leroi I, Connolly MJ. Cognitive impairment in chronic obstructive pulmonary disease and chronic heart failure: A systematic review and meta-analysis of observational studies. *Journal of the American Medical Directors Association*. 2017.
65. Pierobon A, Bottelli ES, Ranzini L, Bruschi C, Maestri R, Bertolotti G, et al. COPD patients' self-reported adherence, psychosocial factors and mild cognitive impairment in pulmonary rehabilitation. *International Journal of Chronic Obstructive Pulmonary Disease*. 2017; 12:2059.
66. Caldera-Alvarado G, Khan D, Defina L, Pieper A, Brown E. Relationship between asthma and cognition: the Cooper Center Longitudinal Study. *Allergy*. 2013; 68:545-8.
67. Gispén FE, Chen DS, Genther DJ, Lin FR. Association between hearing impairment and lower levels of physical activity in older adults. *Journal of the American Geriatrics Society*. 2014; 62:1427-33.
68. Marmeleira J, Laranjo L, Marques O, Pereira C. Physical activity patterns in adults who are blind as assessed by accelerometry. *Adapted Physical Activity Quarterly*. 2014; 31:283-96.
69. Wallhagen MI. The stigma of hearing loss. *The Gerontologist*. 2010; 50:66-75.
70. Schepker CA, Leveille SG, Pedersen MM, Ward RE, Kurlinski LA, Grande L, et al. Effect of pain and mild cognitive impairment on mobility. *Journal of the American Geriatrics Society*. 2016; 64:138-43.
71. Pessanha F, Lustosa L, Carneiro J, Pfrimer K, Fassini P, Alves N, et al. Body Mass Index and Its Relationship with Disability, Chronic Diseases and Frailty in Older People: A Comparison of the Lipschitz and Who Classifications. *The Journal of Frailty & Aging*. 2017; 6:24.
72. Kojima G. Frailty as a predictor of disabilities among community-dwelling older people: a systematic review and meta-analysis. *Disability and Rehabilitation*. 2016:1-12.

73. Sugimoto T, Ono R, Murata S, Saji N, Matsui Y, Niida S, et al. Prevalence and associated factors of sarcopenia in elderly subjects with amnesic mild cognitive impairment or Alzheimer disease. *Current Alzheimer Research*. 2016; 13:718-26.
74. Moon J, Kim K, Choi S, Lim S, Park K, Kim KW, et al. Sarcopenia as a predictor of future cognitive impairment in older adults. *The journal of Nutrition, Health & Aging*. 2016; 20:496-502.
75. Veronese N, Stubbs B, Fontana L, Trevisan C, Bolzetta F, Rui MD, et al. A comparison of objective physical performance tests and future mortality in the elderly people. *The Journals of Gerontology: Series A*. 2017; 72:362-8.
76. Veronese N, Stubbs B, Trevisan C, Bolzetta F, De Rui M, Solmi M, et al. Poor Physical Performance Predicts Future Onset of Depression in Elderly People: Pro. VA Longitudinal Study. *Physical Therapy*. 2017:pzx017.
77. Tyrovolas S, Koyanagi A, Lara E, Santini ZI, Haro JM. Mild cognitive impairment is associated with falls among older adults: findings from the Irish Longitudinal Study on Ageing (TILDA). *Experimental Gerontology*. 2016; 75:42-7.
78. Shimada H, Park H, Makizako H, Tsutsumimoto K, Uemura K, Nakakubo S, et al. Cognitive function and falling among older adults with mild cognitive impairment and slow gait. *Geriatrics & Gerontology International*. 2015; 15:1073-8.
79. Ismail Z, Elbayoumi H, Fischer CE, Hogan DB, Millikin CP, Schweizer T, et al. Prevalence of depression in patients with mild cognitive impairment: A systematic review and meta-analysis. *JAMA psychiatry*. 2017; 74:58-67.
80. Schuch F, Vancampfort D, Firth J, Rosenbaum S, Ward P, Reichert T, et al. Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. *Journal of Affective Disorders*. 2017; 210:139-50.
81. Jorm AF. Is depression a risk factor for dementia or cognitive decline? *Gerontology*. 2000; 46:219-27.
82. Schuch FB, Vancampfort D, Rosenbaum S, Richards J, Ward PB, Veronese N, et al. Exercise for depression in older adults: a meta-analysis of randomized controlled trials adjusting for publication bias. *Revista Brasileira de Psiquiatria*. 2016:0-.
83. Soundy A, Roskell C, Stubbs B, Vancampfort D. Selection, use and psychometric properties of physical activity measures to assess individuals with severe mental illness: a narrative synthesis. *Archives of Psychiatric Nursing*. 2014; 28:135-51.

84. Stubbs B, Firth J, Berry A, Schuch FB, Rosenbaum S, Gaughran F, et al. How much physical activity do people with schizophrenia engage in? A systematic review, comparative meta-analysis and meta-regression. *Schizophrenia Research*. 2016.
85. Ainsworth BE, Macera CA, Jones DA, Reis JP, Addy CL, Bowles HR, et al. Comparison of the 2001 BRFSS and the IPAQ Physical Activity Questionnaires. *Medicine and Science in Sports and Exercise*. 2006; 38:1584-92.
86. Petersen RC. Mild Cognitive Impairment. *Continuum (Minneapolis, Minn)*. 2016; 22:404-18.
87. Leach LS, Butterworth P, Strazdins L, Rodgers B, Broom DH, Olesen SC. The limitations of employment as a tool for social inclusion. *BMC Public Health*. 2010; 10:621.