King's Research Portal

DOI:

[10.1017/S1352465817000765](https://doi.org/10.1017/S1352465817000765)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Fairbrother, N., Thordarson, D. S., Challacombe, F. L., & Sakaluk, J. K. (2018). Correlates and Predictors of New Mothers' Responses to Postpartum Thoughts of Accidental and Intentional Harm and Obsessive Compulsive Symptoms. *Behavioural and Cognitive Psychotherapy*. <https://doi.org/10.1017/S1352465817000765>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Correlates and Predictors of New Mothers' Responses to Postpartum Thoughts of Accidental and Intentional Harm and Obsessive Compulsive Symptoms

Nichole Fairbrother

*Island Medical Program/Psychiatry, University of British Columbia, Room 002, Pearkes Building,
Queen Alexandra Centre for Children's Health, 2400 Arbutus Road, Victoria, British Columbia,
V8N 4V7, Canada*

Dana S. Thordarson

*Division of Midwifery, Department of Family Practice, University of British Columbia, 320–5950
University Boulevard, Vancouver, British Columbia, V6T 1Z3, Canada*

Fiona L. Challacombe

*Centre for Anxiety Disorders and Trauma, South London and Maudsley NHS Trust, 99 Denmark Hill,
London SE5 8AZ, UK*

John K. Sakaluk

*Department of Psychology, University of Victoria Room A240, Cornett Building, 3800 Finnerty Road
(Ring Road), Victoria, British Columbia, V8P 5C2, Canada*

Background: Unwanted, intrusive thoughts of infant-related harm are a normal, albeit distressing experience for most new mothers. The occurrence of these thoughts can represent a risk factor for the development of obsessive compulsive disorder (OCD). As the early postpartum period represents a time of increased risk for OCD development, the transition to parenthood provides a unique opportunity to better understand OCD development. **Aims:** The purpose of this study was to assess components of cognitive behavioural conceptualizations of postpartum OCD in relation to new mothers' thoughts of infant-related harm. **Method:** English-speaking pregnant women ($n = 100$) participated. Questionnaires were completed at approximately 36 weeks of gestation, and at 4 and 12 weeks postpartum. An interview to assess postpartum harm thoughts was administered at 4 and 12 weeks postpartum. Questionnaires assessed OC symptoms, OC-related beliefs, fatigue, sleep difficulties and negative mood.

Correspondence concerning this manuscript should be addressed to Nichole Fairbrother, Room 002, Pearkes Building, Queen Alexandra Centre for Children's Health, 2400 Arbutus Road, Victoria, BC, V8N 4V7, Canada. E-mail: nicholef@uvic.ca

Results: Prenatal OC-related beliefs predicted postpartum OC symptoms, as well as harm thought characteristics and behavioural responses to harm thoughts. The severity of behavioural responses to early postpartum harm thoughts did not predict later postpartum OC symptoms, but did predict frequency and time occupation of *accidental* harm thoughts, and interference in parenting by *intentional* harm thoughts. Strong relationships between OC symptoms and harm thought characteristics, and concurrent sleep difficulties, negative mood and fatigue were also found. **Conclusions:** Findings provide support for cognitive behavioural conceptualizations of postpartum OCD and emphasize the importance of maternal sleep, fatigue and negative mood in the relationship between OC-related beliefs and maternal cognitive and behavioural responses to postpartum harm thoughts.

Keywords: intrusive thoughts, obsessive compulsive disorder, new mothers

Introduction

Studies of violent, infant-related intrusive thoughts experienced by mothers (especially first-time mothers) during the early postpartum period make it clear that they are a very common and distressing experience, affecting more than 90% of new mothers (Abramowitz et al., 2005; Abramowitz and Deacon, 2006; Ahmad et al., 1994; Kleinknecht and Thorndike, 1990; Mann et al., 2010; Meyer et al., 1990; Wynter et al., 2013). Such thoughts may pertain to *accidental* or *intentional* harm to the infant. Given the proportion of new mothers who report thoughts of infant-related harm, it appears that these thoughts are a normal postpartum experience (Kleinknecht and Thorndike, 1990; Mann et al., 2010; Meyer et al., 1990; Wynter et al., 2013).

One of the possible negative consequences of new mothers' experience of infant-related harm thoughts is the development of obsessive compulsive disorder (OCD).

OCD is a mental health condition characterized by obsessions (recurrent, unwanted and distressing thoughts, images or impulses) and/or compulsions (repetitive mental or behavioural acts that the person engages in, in order to decrease the distress associated with the obsessions). OCD is a psychiatric disorder associated with high levels of distress and impairment (Monk, 2001; Nerum et al., 2013; Newham et al., 2014; Nieminen et al., 2009, 2016). For pregnant and postpartum women, OCD has negative implications for fetal and newborn health, mothering and infant development (Challacombe et al., 2016; Saisto et al., 2001; Salomonsson et al., 2013; Stoll et al., 2016; Straub et al., 2012; Sydsjö et al., 2014; Sydsjö et al., 2015; Takegata et al., 2015). OCD sufferers are frequently motivated to conceal their obsessions from others, including health-care providers, due to the often socially unacceptable nature of them and fears about what they might mean (Nilsson et al., 2012). This is especially true for women suffering from postpartum OCD (ppOCD) and experiencing thoughts of harming their infant. They may be fearful that their infant will be removed from their care should they disclose their thoughts and, therefore, are highly motivated to conceal their obsessions.

Based on the now significant number of studies demonstrating an increased risk of OCD onset and exacerbation during the perinatal period (Russell et al., 2013), and borrowing heavily from these existing cognitive behavioural theories of OCD, Fairbrother and Abramowitz (2007) proposed a cognitive conceptualization of postpartum OCD. They hypothesized that the dramatic objective increase in responsibility (i.e., I am responsible for my infant's care and wellbeing) and threat (i.e., my infant is vulnerable to harm and highly cherished) that accompanies the perinatal period, lowers the threshold for OCD development or exacerbation. According to Fairbrother and Abramowitz, cognitive vulnerability to OCD increases the likelihood that normally occurring, albeit upsetting, intrusive infant-related thoughts will be

interpreted as highly significant, threatening, and as otherwise requiring attention in order to prevent some feared negative behavioural or moral consequence. Furthermore, overt and covert behavioural patterns that develop in response to the intrusive thoughts (e.g. checking on the infant, avoidance of the infant, attempts to suppress or counteract intrusive ideation) contribute to the maintenance of the obsessional distress.

A number of studies of expectant parents, and parents of infants, provide preliminary support of for this conceptualization. For example, prenatal measures of dysfunctional OCD-related beliefs have been found to predict postpartum OC symptoms even after controlling for pre-existing OC symptoms, anxiety and depression (Abramowitz et al., 2006). The tendency to negatively interpret the presence and meaning of early postpartum unwanted intrusive infant-related thoughts has also been shown to be predictive of later postpartum OC symptoms, even after controlling for baseline levels of OC symptoms (Abramowitz et al., 2007). Among mothers and fathers who reported intrusive obsessional thoughts about their infant, thought control strategies have been positively associated with obsessional symptoms (Larsen et al., 2006).

In a comparison with pregnant, postpartum and non-childbearing women, general beliefs about responsibility were not found to differ between groups (Barrett et al., 2016). However, among pregnant women, baby-related intrusions were interpreted as implying more responsibility, and among mothers, only baby-related responsibility interpretations were related to OC symptomatology.

As pregnancy and the first 6 months postpartum is a unique time in a woman's life, it provides a brand-new source of unwanted intrusive thoughts, namely thoughts of harm coming to the baby, and therefore represents a naturally occurring experiment that allows us to better understand OCD development. The purpose of the present research was to take advantage of the natural changes in the occurrence of unwanted intrusive thoughts, and symptoms of OCD that occur during the perinatal period to assess components of cognitive behavioural conceptualizations of ppOCD. The novel contribution of this research is our focus on new mothers' thoughts of infant-related harm, and the distinction between thoughts of *accidental* and *intentional* harm. In addition, although associations have been established in non-perinatal OCD (Nota et al., 2016) it is the first study to assess the impact of sleep difficulties on perinatal OC symptoms. Specific study hypotheses are:

Hypothesis 1: Prenatal OC-related beliefs are predictive of the development of postpartum OC symptoms (i.e. OBQ-44 subscale scores will predict postpartum OCI-R obsessing subscale scores, at 4 and 12 weeks postpartum, over and above the contribution of prenatal OCI-R obsessing subscale scores).

Hypothesis 2: Prenatal OC-related beliefs are predictive of maternal postpartum harm thought characteristics and maternal behavioural responses to the thoughts, for both *accidental* and *intentional* harm thoughts (i.e. prenatal OBQ-44 subscale scores will predict: (a) infant-related harm thought characteristics, and (b) the severity of behavioural responses to thoughts, for both *accidental* and *intentional* harm thoughts, at 4 and 12 weeks postpartum).

Hypothesis 3: The use of safety-seeking or other strategies to neutralize postpartum intrusive thoughts is predictive of later ppOC symptoms. Specifically:

- (1) The severity of postpartum behavioural responses to early (4 week) postpartum intrusive thoughts of infant-related harm (*accidental* and *intentional*) predict later (12 week)

postpartum OCI-R obsession subscale scores, controlling for 4 week postpartum OCI-R obsession subscale scores.

- (2) The severity of postpartum behavioural responses to early (4 week) postpartum intrusive thoughts of infant-related harm (*accidental* and *intentional*) predict later harm thought characteristics, controlling for 4 week postpartum harm thought characteristics.

Hypothesis 4: Concurrent negative mood, maternal fatigue and self-reported sleep difficulties are associated with higher levels of OC symptoms and harm thought characteristics. Specifically, 4 week postpartum BDI, MAF and PSQI scores will correlate positively with 4 week postpartum OCI-R obsessing subscale scores and harm thought characteristics. This will be true for 12 week measures also.

Method

Design

This is a prospective cohort study of postpartum maternal affective, cognitive and behavioural responses to postpartum thoughts of infant-related harm. Additional findings from this research have been presented in an earlier publication (Fairbrother and Woody, 2007).

Participants

Participants were 100 English-speaking pregnant women experiencing a medically uncomplicated pregnancy and expecting a healthy first child. They were recruited via two major hospitals, midwifery and physician offices, and prenatal classes in Vancouver, British Columbia, Canada.

The average age of participants was 32.0 years ($SD = 4.3$) and ranged from 23 to 41 years. Participants reported an average of 16.7 ($SD = 2.4$) years of formal education. The majority of the sample was married or living with a romantic partner (97%), was employed (79%), and reported their ethnicity as either Caucasian (76%) or Asian (13%). More than two-thirds of the sample (69%) reported a family income of \$50,000 or greater.

Procedures

Participants were recruited in the third trimester of pregnancy and completed questionnaires prenatally (at approximately 36 weeks gestation), and again at 4 and 12 weeks postpartum. At 4 and 12 weeks postpartum, participants were also administered a semi-structured interview to assess infant-related thoughts of harm.

All but two of the 100 participants completed the prenatal questionnaire package. At 4 weeks postpartum, 91 women completed the questionnaires, and 84 completed the interview. A total of 81 women completed the 12 week questionnaires, and 83 completed the 12 week interview. Altogether, this represents a 16% loss to follow-up. Two women entered the study after they had given birth and therefore provided postpartum data only.

Measures

The *Obsessional Beliefs Questionnaire* (OBQ-44; Obsessive Compulsive Cognitions Working Group (OCCWG), 2005) is a 44-item self-report measure designed to

measure beliefs associated with OCD including: responsibility and threat estimation, perfectionism and need for certainty, and importance and control of thoughts. Psychometric properties of the OBQ-44 are good, although discrimination between subscales is low (OCCWG, 2005).

The *Postpartum Intrusions Interview* (PPII; Fairbrother and Woody, 2007) is a semi-structured interview designed to assess new mothers' thoughts of *accidental* and *intentional* harm related to the newborn. The interview includes an assessment of the content and characteristics of (i.e., frequency of the thoughts, time occupied by the thoughts, distress associated with the thoughts, perceived interference with parenting and other aspects of one's life by the thoughts, and difficulty distracting oneself from the thoughts), emotional and behavioural responses to, and the natural history of *accidental* and *intentional* harm thoughts. The PPII was designed specifically for use in this project, the details of which are described in full in an earlier publication (Fairbrother and Woody, 2008).

To date, studies of postpartum intrusive thoughts and ppOCD have focused on the assessment of obsessive compulsive symptoms assessed using self-report measures of symptoms of OCD, validated in non-perinatal samples. We were concerned that some elements of typical self-report measures of OC symptoms might not be recognized by new mothers as relevant to their infant-related harm thoughts, and consequently underestimate their emotional and behavioural responses to infant-related harm thoughts. Specifically, symptoms of ordering and arranging counting or hoarding may be less relevant to women whose OC symptoms primarily pertain to infant-related harm thoughts. Beyond this, references to unpleasant or nasty thoughts, or difficulties controlling one's thoughts may not be perceived by new mothers as pertaining to their thoughts of infant-related harm. Given this, we developed a series of questions similar to those used to assess obsession severity in the Y-BOCS. Specifically, we asked participants about harm thought frequency, distress and perceived interference as well as how time consuming the thoughts are, and how much control they have over the thoughts. We believe this assessment provides a clearer view of the severity and scope of new mothers' cognitive and emotional responses to early postpartum harm thoughts compared with more standard symptom measures of OCD.

The *Beck Depression Inventory-II* (BDI-II; Beck et al., 1996) is a 21-item self-report measure of depression. Scale items are scored on a 4-point scale. Psychometric evaluations of the BDI-II suggest a two-factor solution and indicate that the scale possesses excellent reliability and validity (Beck et al., 1996; Dozois et al., 1998).

The *Multidimensional Assessment of Fatigue Scale* (MAF; Belza, 1990) is a 16-item self-report measure of fatigue that produces a global fatigue index (GFI). Individual MAF items are scored on a Likert-type scale. GFI scores range in value from 0 (no fatigue) to 50 (severe fatigue). Cronbach alphas for the measure are strong (i.e. range from .88 to .96) (Belza et al., 1993; Belza, 1995; Bormann et al., 2001; Cella et al., 2005; Hall et al., 2006; Jump et al., 2004; Meek et al., 2000; Wambach, 1998; Williams et al., 1999; Winstead-Fry, 1998). The MAF is also stable over time (Belza, 1995; Bormann et al., 2001; Grady et al., 1998; Meek et al., 2000), and responsive to change (Belza et al., 2005; Hall et al., 2006; Kaltwasser et al., 2001; Roscoe et al., 2005; Wambach, 1998; Williams et al., 1999).

The *Pittsburgh Sleep Quality Index* (PSQI; Buysse et al., 1989) is a self-report measure of sleep quality and disturbance over the previous month, with higher scores indicating reduced sleep quality. This scale has shown good internal consistency and test-retest reliability as

well as good construct validity (Backhaus et al., 2002; Buysse et al., 1989; Carpenter and Andrykowski, 1998).

Data analytic approach

When possible, data were analysed using structural equation modeling (SEM) of observed variables, using the *lavaan* package (Rosseel, 2012) for *R* (R Core Team, 2016). We used full-information maximum likelihood estimation for models with variables that were normally distributed, while using diagonal weighted least-squares estimation for models with a presence of non-normal data (Rhemtulla et al., 2012). Otherwise, data were analysed using IBM, SPSS version 23. As the three OBQ-44 subscales were highly intercorrelated ($r = 0.64$ to 0.76), collinearity becomes an issue. In order to avoid collinearity problems, analyses involving the OBQ-44 were carried out using total scale scores only. Descriptive findings are presented in the form of means and standard deviations.

Hypothesis 1 was tested using SEM. Specifically, we fitted a model with prenatal OCI-R obsessing subscale scores and prenatal OBQ-44 total scale scores predicting both 4 week and 12 week postpartum OCI-R obsessing subscale scale scores, while modelling the residual correlation between these similar outcomes and predictors.

As *Hypothesis 2* and its related analyses involve a single predictor only (i.e. OBQ-44 total scores), bivariate correlations were calculated for the relationship between prenatal OBQ-44 total scores and infant-related harm thought characteristics for both *accidental* and *intentional* harm thoughts, at both 4 and 12 weeks postpartum.

Hypothesis 3 was tested using SEM for thoughts of accidental harm, whereas linear regression was used for analysing thoughts of intentional harm due to sparse data. For the SEM portion of this hypothesis, we fitted a model in which 12 week postpartum OCI-R obsessing subscale scores and 12 week postpartum thoughts of *accidental* harm were predicted by 4 week postpartum OCI-R obsessing scores and 4 week postpartum thoughts of *accidental* harm, respectively, as well as severity of behavioural responses to *accidental* infant-related harm thoughts.

For testing *Hypothesis 3* with respect to thoughts of intentional harm, we used linear regression models for each of the outcome variables (i.e. 12 week postpartum OCI-R obsessing subscale scores, and 12 week postpartum harm thought characteristics for thoughts of *intentional* harm). In step one of each regression, either 4 week postpartum OCI-R obsessing subscale scores or 4 week postpartum harm thought characteristics were entered. Severity of behavioural responses to the infant-related harm thoughts (*intentional*) was entered in Step 2.

Finally, *Hypothesis 4* was tested using bivariate correlations.

Results

Descriptive findings

Means and standard deviations for all scale scores are presented in [Table 1](#).

Hypothesis 1

Results of the SEM in which postpartum OCI-R obsession subscale scores were predicted by prenatal OCI-R obsessing scores and prenatal OBQ-44 total scores are presented in [Fig. 1](#).

Table 1. Descriptives: means and standard deviations

Variable	Prenatal	4 week	12 week
OBQ-44 total	132.2 (44.7)	128.0 (45.7)	127.6 (51.0)
OCI-R Obsessing Subscale total	2.0 (2.7)	1.4 (2.2)	1.5 (2.3)
Accidental harm thought characteristics			
Often	n/a	2.0 (1.0)	1.8 (1.0)
Time	n/a	1.3 (0.9)	0.9 (0.9)
Distress	n/a	1.3 (0.9)	0.8 (0.9)
Parent	n/a	0.2 (0.5)	0.1 (0.4)
Life	n/a	0.3 (0.7)	0.2 (0.5)
Distract	n/a	0.5 (0.7)	0.2 (0.5)
Intentional harm thought characteristics			
Often	n/a	0.7 (0.7)	0.9 (0.7)
Time	n/a	0.4 (0.6)	0.5 (0.6)
Distress	n/a	2.0 (1.0)	1.6 (0.9)
Parent	n/a	0.2 (0.4)	0.1 (0.3)
Life	n/a	0.0 (0.2)	0.1 (0.3)
Distract	n/a	0.4 (0.7)	0.4 (0.6)
Severity of behavioural responses to accidental harm thoughts	n/a	5.1 (2.8)	3.8 (2.3)
Severity of behavioural responses to intentional harm thoughts	n/a	3.5 (1.9)	3.4 (1.9)
PSQI	7.6 (3.6)	9.3 (3.5)	7.2 (3.3)
GAF	29.4 (9.6)	30.7 (9.8)	26.7 (11.9)
BDI	11.7 (8.3)	10.3 (6.3)	10.3 (8.3)

Figure 1. SEM of postpartum OCI-R Obsessing Subscale scores predicted by prenatal OCI-R Obsessing Subscale score and OBQ-44 total score. Directional pathways are standardized slopes. ** $p < .01$, *** $p < .001$.

Indexes of model fit are uninformative for this model, as it is fully saturated and therefore perfectly fits the data. As expected, prenatal OCI-R obsessing scores and prenatal OBQ-44 total scores were both uniquely and positively associated with greater 4 week and 12 week postpartum OCI-R obsessing scores. OBQ-44 total scores uniquely explained an additional

Table 2. Proportion of variance (r^2) in postpartum harm thought characteristics and behavioural responses to postpartum harm thoughts, at 4 and 12 weeks postpartum, predicted by prenatal OBQ-44 total scores

	r^2 values						Behavioural responses
	Often	Time	Distress	Parenting	Life	Distract	
4 weeks postpartum							
Accidental harm ($n = 87$)	.04	.13**	.31***	.11**	.08**	.16***	.23***
Intentional harm ($n = 43$)	.04	.13*	.18**	.00	.00	.00($n = 42$)	.16**
12 weeks postpartum							
Accidental harm ($n = 77$)	.03	.09**	.15**	.08*	.10**	.18***	.17***
Intentional harm ($n = 15$)	.03	.20	.50**	.36*	.03	.13	.41*

* $p < .05$, ** $p < .01$, *** $p < .001$.

6.60% of the variance in 4 week postpartum OCI-R obsessing subscale scores, and 7.80% of the variance in 12 week OCI-R obsessing subscale scores.

Hypothesis 2

The proportion of variance in infant-related harm thought characteristics, for both *accidental* and *intentional* harm thoughts, at both 4 and 12 weeks postpartum, accounted for by OBQ-44 total scores (i.e. squared bivariate correlations) are presented in Table 2.

Prenatal OBQ-44 total scores significantly predicted all 4 week and 12 week *accidental* harm thought characteristics (other than thought frequency), and the severity of behavioural responses to the thoughts. For 4 week *intentional* harm thoughts, prenatal OBQ-44 total scores predicted time occupied by the thoughts, distress associated with the thoughts, and the severity of behavioural responses to the thoughts. For 12 week *intentional* harm thought characteristics, OBQ-44 total scores predicted distress associated with the thoughts, perceived interference with parenting, and the severity of behavioural responses to the thoughts.

Hypothesis 3

We have reported the zero-order correlations and the beta-values for each analysis in Table 3. This model fitted the data extremely well: $\chi^2(42) = 10.60, p = 1.00$, CFI = 1.00, SRMR = .06. Although a number of zero-order correlations for severity of behavioural responses to both *accidental* and *intentional* harm thoughts were significant, once 4 week OC symptoms and harm thought characteristics were controlled for (either through SEM, for *accidental* harm thoughts, or through regression models, for *intentional* harm thoughts, see also Table 3), behavioural response severity only predicted 12 week postpartum *accidental* harm thought frequency and time occupied by the thoughts, as well as *intentional* harm thought perceived interference with parenting.

Table 3. Associations between 4 week severity of behavioural responses to harm thoughts and 12 week postpartum outcomes before (r) and after (β) controlling for 4 week postpartum outcomes

12 week postpartum outcome variables	Accidental ($n = 80$)		Intentional ($n = 12$)	
	r	β	r	β
OCI-R Obsessing Subscale scores	.51***	0.29	.42** ($n = 39$)	0.12
Often: thought frequency	.29**	0.28**	.55	0.41
Time: time occupied by the thoughts	.53***	0.66*	.58*	0.04
Distress: how upsetting the thoughts are	.39***	−0.05	.33	0.40
Parent: interference with parenting	.39***	0.24	.47	0.90*
Life: interference with other areas of life?	.40***	0.23	n/a	n/a
Distract: difficulty distracting oneself from the thoughts	.55***	0.15	.40	0.53

* $p < .05$, ** $p < .01$, *** $p < .001$. β for *accidental* variables estimated using SEM; β for *intentional* variables estimated using separate regression models.

Hypothesis 4

The majority of the significant correlations were for *accidental* harm thought characteristics, at both postpartum assessment points. For *intentional* harm thought characteristics, only two correlations were significant: time occupied by the thoughts and BDI-II scores at 4 weeks postpartum, and perceived interference with parenting and BDI-II scores at 12 weeks postpartum. Correlations are presented in Table 4.

Discussion

The primary purpose of this research was to assess elements of cognitive behavioural conceptualizations of ppOCD (Fairbrother and Abramowitz, 2007) and build on earlier findings pertaining to this conceptualization (Abramowitz et al., 2006, 2007; Barrett et al., 2016; Larsen et al., 2006). Ours is the first study to distinguish between, and explore, separately, postpartum thoughts of *accidental* versus *intentional* harm.

Hypothesis 1

Our prediction that prenatal OC-related beliefs would predict postpartum OC symptoms, even when controlling for the contribution of prenatal OC symptoms was supported by the data. Specifically, prenatal obsessive beliefs were found to predict unique variance in 4 week and 12 week postpartum OC symptoms, even after taking prenatal OC symptoms into account. These findings further support cognitive behavioural conceptualizations of ppOCD – namely that pre-existing, maladaptive beliefs about responsibility, uncertainty, and the importance and control of thoughts are a risk factor for OC symptom development. However, it is worth noting that the proportion of variance accounted for was small. This may be partially a function of standard OC symptom measures (e.g. the OCI-R) not being fully relatable to women

Table 4. Correlations between OCI-R Obsessing Subscale scores and harm thought characteristics (for both accidental and intentional harm thoughts), and measures of sleep quality, fatigue and depressed mood, at both 4 and 12 weeks postpartum

4 weeks postpartum			
	Global PSQI	GFI (MAF)	BDI
OCI-R	<i>n</i> = 74	<i>n</i> = 80	<i>n</i> = 86
Obsessing Subscale	.35**	.16	.47***
Accidental harm thought characteristics	<i>n</i> = 74	<i>n</i> = 80	<i>n</i> = 86
Often: thought frequency	-.03	.10	.11
Time: time occupied by the thoughts	.05	.08	.10
Distress: how upsetting the thoughts are	.17	.24*	.26*
Parent: interference with parenting	.39**	.34**	.51***
Life: interference with other areas of life?	.23	.28*	.34**
Distract: difficulty distracting oneself from the thoughts	.14	.23*	.21*
Intentional harm thought characteristics	<i>n</i> = 39	<i>n</i> = 39	<i>n</i> = 42
Often: thought frequency	-.12	-.01	.04
Time: time occupied by the thoughts	.22	.18	.38*
Distress: how upsetting the thoughts are	-.00	-.02	.14
Parent: interference with parenting	-.24	.04	.03
Life: interference with other areas of life?	-	.11	.05
Distract: difficulty distracting oneself from the thoughts	.02	.21	.06
12 weeks postpartum			
	Global PSQI	GFI (MAF)	BDI
OCI-R	<i>n</i> = 74	<i>n</i> = 81	<i>n</i> = 81
Obsessing Subscale	.18	.44***	.67***
Accidental harm thought characteristics	<i>n</i> = 69	<i>n</i> = 76	<i>n</i> = 76
Often: thought frequency	.01	.11	.08
Time: time occupied by the thoughts	.07	.22	.19
Distress: how upsetting the thoughts are	.32**	.33**	.32**
Parent: interference with parenting	.07	.26*	.48***
Life: interference with other areas of life?	.13	.21	.30*
Distract: difficulty distracting oneself from the thoughts	.13	.38**	.31**
Intentional harm thought characteristics	<i>n</i> = 15	<i>n</i> = 15	<i>n</i> = 15
Often: thought frequency	-.16	-.22	-.14
Time: time occupied by the thoughts	.34	.14	-.01
Distress: how upsetting the thoughts are	-.05	-.03	.12
Parent: interference with parenting	-.13	.45	.83***
Life: interference with other areas of life?	-.13	-.07	.09
Distract: difficulty distracting oneself from the thoughts	-.17	-.01	.37

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

experiencing postpartum OC symptoms specific to thoughts of infant-related harm. With more postpartum appropriate measurement of OC symptoms, we may see prenatal beliefs about thoughts predicting a greater proportion of unique variance in OC symptoms. Additionally, properties of the OBQ mean that we were not able to distinguish the relative contribution of the subscales.

Hypothesis 2

Our prediction that prenatal OC-related beliefs would predict maternal postpartum harm thought characteristics and maternal behavioural responses to the thoughts, for both *accidental* and *intentional* harm thoughts, was also supported by the data. Most of the significant findings were for thoughts of *accidental* harm, and provide empirical support for cognitive behavioural conceptualizations of OCD. Specifically, these conceptualizations posit that prenatal OC-related beliefs represent a risk factor for OC symptom development. Our findings provide specific support for the hypothesis that OC-related beliefs are predictive of postpartum, harm thought-related OC symptoms. Furthermore, these findings also provide support for the use of perinatal specific measures of OC symptoms (e.g. the PPII). It may be that measures specific to postpartum thoughts of harm (i.e. the PPII), similar to the Y-BOCS (Goodman et al., 1989) better capture OC symptoms in response to postpartum thoughts of infant-related harm than do more traditional OC symptom measures. It is also worth noting, however, that prenatal OC symptoms were not controlled for in the analyses involving measures of harm thought characteristics as there was not a prenatal parallel. Had this been possible, the variance accounted for above would most likely have been lower.

OC-related beliefs predicted a significant proportion of the variance in all accidental harm thought characteristics, with the exception of harm thought frequency. This finding is consistent with cognitive behavioural conceptualizations of OCD which posit that it is not the occurrence of unwanted, intrusive ideation that predict OC symptom development, but rather the appraisals that one makes of the occurrence of the thoughts (Rachman, 1998). OC-related beliefs most strongly predicted the following accidental harm thought characteristics, at both 4 and 12 weeks postpartum: distress associated with the thoughts, and one's ability to distract oneself from the thoughts. With respect to infant-related thoughts of *accidental* harm, negative beliefs about thoughts are predictive of, and may even increase how distressing these thoughts are and how mentally captivating they are.

Fewer significant relationships were found for thoughts of *intentional* harm. At 4 weeks postpartum, these analyses included only 43 women, and at 12 weeks only 15. It is clear that this analysis was underpowered, therefore reducing our ability to detect effects related to intentional harm thoughts, and compromising the precision of the effects that we were able to detect. It is interesting that, among significant variables, OC-related beliefs predicted the time occupied by the thoughts, for both *accidental* and *intentional* harm thoughts, but not the frequency with which the thoughts occurred. This finding is consistent with cognitive behavioural theories of OCD. Specifically, CBT theories of OCD posit that the occurrence of unwanted, intrusive thoughts about one's baby are normal, but maladaptive OC-related beliefs may significantly increase how time-consuming they are (Fairbrother and Abramowitz, 2007, Rachman, 1998). For example, maladaptive OC-related beliefs may result in time spent ruminating about the thoughts, or trying to neutralize them. Prenatal OC-related beliefs also

predicted half of the variance in the distress associated with 12 week *intentional* harm thoughts, and slightly more than a third of the variance in how much these thoughts interfered with parenting. One might imagine that the longer the thoughts persist, the more strongly OC-related beliefs might impact how distressing and impairing the thoughts become. Although the sample sizes for these analyses are very small, and therefore findings should be interpreted cautiously, the magnitude of the relationships makes it clear that they merit additional assessment in a larger sample.

Prenatal OC-related beliefs predicted significant variance in behavioural responses to both accidental and intentional harm, at both 4 and 12 weeks postpartum. The strongest effect was for 12 week postpartum thoughts of intentional harm. This suggests that perhaps negative beliefs about unwanted, intrusive thoughts trigger a stronger urge to engage in avoidance, reassurance seeking, checking or other behavioural and mental activities in response to unwanted thoughts of intentionally harming one's infant, compared with thoughts of accidental harm coming to one's infant. This is consistent with the finding that thoughts of intentional harm are reported to be more distressing than thoughts of accidental harm.

Hypothesis 3

We hypothesized that the use of safety-seeking or other strategies to neutralize postpartum intrusive thoughts would be predictive of later ppOC symptoms. This hypothesis was only weakly supported by the data. Although a number of zero order correlations between the severity of behavioural responses to *accidental* and *intentional* harm thoughts and 12 week postpartum OC symptoms, and harm thought characteristics were significant, few significant relationships remained once 4 week variables (i.e. OC symptoms and harm thought characteristics) were controlled for.

The severity of behavioural responses to harm thoughts did not predict unique variance in OCI-R obsessing subscale scores at 12 weeks postpartum, for either *accidental* or *intentional* harm thoughts. It may be that the OCI-R is neither specific enough (i.e. non-perinatal in focus), nor sensitive enough to detect the experiences of postpartum women, as it refers to more generic rather than postpartum-specific intrusive thoughts. Furthermore, mean scores on the OCI-R were low at all three time points, suggesting that perhaps there was insufficient variance in OC symptoms, as measured by the OCI-R, to detect the hypothesized relationships.

With respect to harm thought characteristics, a small number of significant relationships emerged. Specifically, for thoughts of *accidental* harm, behavioural response severity predicted the frequency of and time occupied by 12 week postpartum thoughts. With respect to thoughts of *intentional* harm, behavioural response severity predicted perceived interference with parenting only. What is most notable is that the severity of 4 week postpartum behavioural responses to *intentional* harm thoughts predicted close to half of the variance in perceived interference with parenting (from *intentional* harm thoughts) at 12 weeks postpartum, even after controlling for 4 week postpartum perceived interference with parenting. This finding, that new mothers perceive that their OC symptoms (i.e. behavioural responses to harm thoughts) significantly interfere with parenting is consistent with findings by others (Barrett et al., 2016). Due to the very small sample involved in this analysis ($n = 12$), the possibility that behavioural responses to intentional harm thoughts may contribute negatively and significantly to parenting deserves further exploration.

Hypothesis 4

Sleep. OC symptoms were positively associated with sleep difficulties at 4 weeks postpartum, but not 12 weeks postpartum. Perceived interference with parenting from 4 week *accidental* harm thoughts, and with distress associated with 12 week *accidental* harm thoughts were also positively associated with sleep difficulties. This is consistent with the fact that women reported the most sleep difficulties at 4 weeks postpartum. It may be that sleep has a stronger association with OC symptoms and perceived interference with parenting at higher levels of sleep interference. Perhaps, in part, sleep difficulties at 12 weeks postpartum are driven by distress associated with harm thoughts, more than by interference by one's infant. This would explain why sleep difficulties were associated with distress associated with *accidental* harm thoughts at 12 but not 4 weeks postpartum.

Fatigue. OC symptoms were positively associated with sleep difficulties, but at 12 weeks postpartum only. Maternal fatigue was also positively and significantly related to 4 and 12 week distress associated with harm thoughts, harm thought perceived interference with parenting, as well as difficulty distracting oneself from the thoughts, but only for *accidental* harm thoughts. At 4 weeks postpartum, maternal fatigue was also associated with interference (from thoughts of *accidental* harm only) with other areas of one's life. The strongest relationships were for perceived interference with parenting at 4 weeks, and difficulty distracting oneself from the thoughts at 12 weeks. No significant relationships between maternal fatigue and intentional harm thought characteristics were found.

Negative mood. Not surprisingly, OC symptoms were positively associated with negative mood at both 4 and 12 weeks postpartum. There is compelling evidence that OC symptoms and negative mood are correlated and frequently co-occur (Ruscio et al., 2010). At both 4 and 12 weeks postpartum, negative mood was also positively associated with distress associated with harm thoughts, perceived harm thought interference with parenting and other aspects of life, and difficulty distracting oneself with the thoughts, but only for *accidental* harm thoughts. For *intentional* harm thoughts, the only significant relationships were between negative mood and time occupied by the thoughts at 4 weeks postpartum and perceived interference with parenting at 12 weeks postpartum. The implication of this latter finding is that while the occurrence of postpartum thoughts of infant-related harm appears to be a normal experience for new mothers, being very distressed by them, or for them to interfere significantly with one's life may be indicative of significant mood difficulties. This also suggests that negative mood may be an important mechanism via which OCD interferes with parenting.

Limitations and future directions

The primary limitation of this research is that it is based on a small, upper-middle class sample. Sample size limitations were particularly problematic for our analyses of thoughts of *intentional* harm. Studies involving large enough sample sizes to be able to speak more definitively about the relationship between postpartum thoughts of *intentional* harm and OC symptoms are required. Our team has recently completed data collection for a study in which we replicate many of the above findings in a sample of over 600 women. This much larger sample size will allow us to provide more definitive and stable estimates of the relationships described above, in particular with respect to thoughts of *intentional* harm. Further exploration of the impact of postpartum OC symptoms on perceived interference with parenting is also warranted.

Conclusions

Our study provides important support for CBT conceptualization of ppOCD and extends our understanding of the relationship between pre-existing OC-related beliefs, and OC symptoms and characteristics of new mothers' thoughts of infant-related harm. Prenatal OC-related beliefs predicted change in OC symptoms from prenatal to postpartum, as well as characteristics of and behavioural responses to postpartum intrusive thoughts of harm. This was particularly true for thoughts of *accidental* harm coming to infants, which were more common and therefore for which we had more power to detect an effect. Behavioural responses to early postpartum thoughts did not predict worsening of OC symptom thoughts later in postpartum, but did predict perceived interference with parenting from *intentional* harm thoughts. An issue deserving of consideration is that of disclosure. It may be that unwanted, intrusive thoughts of *intentional*, infant-related harm are only experienced by half of all new mothers. However, it may also be that, despite our efforts to normalize these thoughts and maximize disclosure, some women who experience them may nevertheless not report them. Depressed mood was associated with more OC symptoms postpartum. Poor sleep predicted more OC symptoms at 4 weeks, whereas fatigue predicted more OC symptoms at 12 weeks. Depressed mood, poor sleep and fatigue did not predict the occurrence or frequency of intrusive thoughts of harm, but did predict other harm thought characteristics such as distress and perceived interference with parenting.

Acknowledgements

We would like to express our gratitude to Dr Michael Papsdorf, Ms Elizabeth Horner, Ms Marion MacKay-Dunn, Ms Maria Watson, and our study participants for their invaluable contributions to this project.

Financial support: This work was supported by the Lion's Gate Healthcare Research Foundation (\$39,837; 2004–2006; Sheila Woody, PI; Nichole Fairbrother, Co-I).

Conflicts of interest: Nichole Fairbrother, Dana S. Thordarson, Fiona L. Challacombe and John K. Sakaluk have no conflicts of interest with respect to this publication.

Ethical statement: The authors have complied with the Ethical Principles of Psychologists and Codes of Conduct as set forth by the APA. This study was reviewed and approved by the University of British Columbia's Behavioural Research Ethics Board and the Children's and Women's Health Centre of British Columbia Research Review Committee (B04-0370).

References

- Abramowitz, J. S. and Deacon, B. J.** (2006). Psychometric properties and construct validity of the Obsessive Compulsive Inventory-Revised: replication and extension with a clinical sample. *Journal of Anxiety Disorders*, 20, 1016–1035. doi: [10.1016/j.janxdis.2006.03.001](https://doi.org/10.1016/j.janxdis.2006.03.001)
- Abramowitz, J. S., Khandker, M., Nelson, C. A., Deacon, B. J. and Rygwall, R.** (2006). The role of cognitive factors in the pathogenesis of obsessive-compulsive symptoms: a prospective study. *Behaviour Research and Therapy*, 44, 1361–1374. doi: [10.1016/j.brat.2005.09.011](https://doi.org/10.1016/j.brat.2005.09.011)
- Abramowitz, J. S., Nelson, C. A., Rygwall, R. and Khandker, M.** (2007). The cognitive mediation of obsessive-compulsive symptoms: a longitudinal study. *Journal of Anxiety Disorders*, 21, 91–104. doi: [10.1016/j.janxdis.2006.05.003](https://doi.org/10.1016/j.janxdis.2006.05.003)

- Abramowitz, J. S., Tolin, D. F. and Diefenbach, G. J.** (2005). Measuring change in OCD: Sensitivity of the obsessive-compulsive inventory-revised. *Journal of Psychopathology and Behavioral Assessment*, 27, 317–324. doi: [10.1007/s10862-005-2411-y](https://doi.org/10.1007/s10862-005-2411-y)
- Ahmad, T. F. E., Suleiman, R. and Shougah, D.** (1994). Mixed anxiety depression in pregnancy: the influence of misconceptions. *Arab Journal of Psychiatry*, 5, 103–113.
- Backhaus, J., Junghanns, K., Broocks, A., Riemann, D. and Hohagen, F.** (2002). Test–retest reliability and validity of the pittsburgh sleep quality index in primary insomnia. *Journal of Psychosomatic Research*, 53, 737–740. doi: [10.1016/S0022-3999\(02\)00330-6](https://doi.org/10.1016/S0022-3999(02)00330-6)
- Barrett, R., Wroe, A. L. and Challacombe, F. L.** (2016). Context is everything: an investigation of responsibility beliefs and interpretations and the relationship with obsessive-compulsive symptomatology across the perinatal period. *Behavioural and Cognitive Psychotherapy*, 44, 318. doi: [10.1017/S1352465815000545](https://doi.org/10.1017/S1352465815000545)
- Beck, A. T., Steer, R. A. and Brown, G. K.** (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Belza, B.** (1990). Multidimensional Assessment of Fatigue (MAF) Scale: User's Guide. Available at: <http://www.son.washington.edu/research/maf/users-guide.asp>
- Belza, B. L.** (1995). Comparison of self-reported fatigue in rheumatoid arthritis and controls. *Journal of Rheumatology*, 22, 639.
- Belza, B. L., Henke, C. J., Yelin, E. H., Epstein, W. V. and Gilliss, C. L.** (1993). Correlates of fatigue in older adults with rheumatoid arthritis. *Nursing Research*, 42, 93–99. doi: [10.1097/00006199-199303000-00006](https://doi.org/10.1097/00006199-199303000-00006)
- Belza, B., Steele, B. G., Cain, K., Coppersmith, J., Howard, J. and Lakshminarayan, S.** (2005). Seattle obstructive lung disease questionnaire: sensitivity to outcomes in pulmonary rehabilitation in severe pulmonary illness. *Journal of Cardiopulmonary Rehabilitation*, 25, 107–114. doi: [10.1097/00008483-200503000-00011](https://doi.org/10.1097/00008483-200503000-00011)
- Bormann, J., Shively, M., Smith, T. L. and Gifford, A. L.** (2001). Measurement of fatigue in HIV-positive adults: reliability and validity of the global fatigue index. *Journal of the Association of Nurses in AIDS Care*, 12, 75–83. doi: [10.1016/S1055-3290\(06\)60146-5](https://doi.org/10.1016/S1055-3290(06)60146-5)
- Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R. and Kupfer, D. J.** (1989). The pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Research*, 28, 193–213. doi: [10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4)
- Carpenter, J. S. and Andrykowski, M. A.** (1998). Psychometric evaluation of the pittsburgh sleep quality index. *Journal of Psychosomatic Research*, 45, 5–13. doi: [10.1016/S0022-3999\(97\)00298-5](https://doi.org/10.1016/S0022-3999(97)00298-5)
- Cella, D., Yount, S., Sorensen, M., Chartash, E., Sengupta, N. and Grober, J.** (2005). Validation of the functional assessment of chronic illness therapy fatigue scale relative to other instrumentation in patients with rheumatoid arthritis. *Journal of Rheumatology*, 32, 811–819.
- Challacombe, F. L., Salkovskis, P. M., Woolgar, M., Wilkinson, E. L., Read, J. and Acheson, R.** (2016). Parenting and mother–infant interactions in the context of maternal postpartum obsessive-compulsive disorder: effects of obsessional symptoms and mood. *Infant Behavior and Development*, 44, 11–20.
- Dozois, D. J. A., Dobson, K. S. and Ahnberg, J. L.** (1998). A psychometric evaluation of the Beck Depression Inventory-II. *Psychological Assessment*, 10, 83–89. doi: [10.1037//1040-3590.10.2.83](https://doi.org/10.1037//1040-3590.10.2.83)
- Fairbrother, N. and Abramowitz, J. S.** (2007). New parenthood as a risk factor for the development of obsessional problems. *Behaviour Research and Therapy*, 45, 2155–2163. doi: [10.1016/j.brat.2006.09.019](https://doi.org/10.1016/j.brat.2006.09.019)
- Fairbrother, N. and Woody, S. R.** (2007). Fear of childbirth and obstetrical events as predictors of postnatal symptoms of depression and post-traumatic stress disorder. *Journal of Psychosomatic Obstetrics and Gynecology*, 28, 239–242. doi: [10.1080/01674820701495065](https://doi.org/10.1080/01674820701495065)
- Fairbrother, N. and Woody, S. R.** (2008). New mothers' thoughts of harm related to the newborn. *Archives of Women's Mental Health*, 11, 221–229. doi: [10.1007/s00737-008-0016-7](https://doi.org/10.1007/s00737-008-0016-7)

- Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Fleischmann, R. L., Hill, C. L. et al. (1989). The Yale-Brown Obsessive Compulsive Scale I. Development, use and reliability. *Archives of General Psychiatry*, 46, 1006–1011. doi: [10.1001/archpsyc.1989.01810110048007](https://doi.org/10.1001/archpsyc.1989.01810110048007)
- Grady, C., Anderson, R. and Chase, G. A. (1998). Fatigue in HIV-infected men receiving investigational interleukin-2. *Nursing Research*, 47, 227–234. doi: [10.1097/00006199-199807000-00008](https://doi.org/10.1097/00006199-199807000-00008)
- Hall, W. A., Clauson, M., Carty, E. M., Janssen, P. A. and Saunders, R. A. (2006). Effects on parents of an intervention to resolve infant behavioral sleep problems. *Pediatric Nursing*, 32, 243.
- Jump, R. L., Fifield, J., Tennen, H., Reisine, S. and Giuliano, A. J. (2004). History of affective disorder and the experience of fatigue in rheumatoid arthritis. *Arthritis Care and Research*, 51, 239–245. doi: [10.1002/art.20243](https://doi.org/10.1002/art.20243)
- Kaltwasser, J. P., Kessler, U., Gottschalk, R., Stucki, G. and Möller, B. (2001). Effect of recombinant human erythropoietin and intravenous iron on anemia and disease activity in rheumatoid arthritis. *Journal of Rheumatology*, 28, 2430–2436.
- Kleinknecht, R. A. and Thorndike, R. M. (1990). The mutilation questionnaire as a predictor of blood/injury fear and fainting. *Behaviour Research and Therapy*, 2, 429–437. doi: [10.1016/0005-7967\(90\)90163-D](https://doi.org/10.1016/0005-7967(90)90163-D)
- Larsen, K. E., Schwartz, S. A., Whiteside, S. P. and Khandker, M. (2006). Thought control strategies used by parents reporting postpartum obsessions. *Journal of Cognitive Psychotherapy*, 20, 435.
- Mann, R., Gilbody, S. and Adamson, J. (2010). Prevalence and incidence of postnatal depression: what can systematic reviews tell us? *Archives of Women's Mental Health*, 13, 295–305. doi: [10.1007/s00737-010-0162-6](https://doi.org/10.1007/s00737-010-0162-6)
- Meek, P. M., Nail, L. M., Barsevick, A., Schwartz, A. L., Stephen, S., Whitmer, K. et al. (2000). Psychometric testing of fatigue instruments for use with cancer patients. *Nursing Research*, 49, 181–190. doi: [10.1097/00006199-200007000-00001](https://doi.org/10.1097/00006199-200007000-00001)
- Meyer, T. J., Miller, M. L., Metzger, R. L. and Borkovec, T. D. (1990). Development and validation of the Penn State Worry Questionnaire. *Behaviour Research and Therapy*, 28, 487–495. doi: [10.1016/0005-7967\(90\)90135-6](https://doi.org/10.1016/0005-7967(90)90135-6)
- Monk, C. (2001). Stress and mood disorders during pregnancy: implications for child development. *Psychiatric Quarterly*, 72, 347–357. doi: [10.1023/A:1010393316106](https://doi.org/10.1023/A:1010393316106)
- Nerum, H., Halvorsen, L., Straume, B., Sørli, T. and Øian, P. (2013). Different labour outcomes in primiparous women that have been subjected to childhood sexual abuse or rape in adulthood: a case-control study in a clinical cohort. *BJOG: An International Journal of Obstetrics and Gynaecology*, 120, 487–495. doi: [10.1111/1471-0528.12053](https://doi.org/10.1111/1471-0528.12053)
- Newham, J. J., Wittkowski, A., Hurley, J., Aplin, J. D. and Westwood, M. (2014). Effects of antenatal yoga on maternal anxiety and depression: a randomized controlled trial. *Depression and Anxiety*, 31, 631–640. doi: [10.1002/da.22268](https://doi.org/10.1002/da.22268)
- Nieminen, K., Berg, I., Frankenstein, K., Viita, L., Larsson, K., Persson, U. et al. (2016). Internet-provided cognitive behaviour therapy of posttraumatic stress symptoms following childbirth: a randomized controlled trial. *Cognitive Behaviour Therapy*, 45, 287–306. doi: [10.1080/16506073.2016.1169626](https://doi.org/10.1080/16506073.2016.1169626)
- Nieminen, K., Stephansson, O. and Ryding, E. L. (2009). Women's fear of childbirth and preference for cesarean section – a cross-sectional study at various stages of pregnancy in Sweden. *Acta Obstetrica et Gynecologica Scandinavica*, 88, 807–813. doi: [10.1080/00016340902998436](https://doi.org/10.1080/00016340902998436)
- Nilsson, C., Lundgren, I., Karlstrom, A. and Hildingsson, I. (2012). Self reported fear of childbirth and its association with women's birth experience and mode of delivery: a longitudinal population-based study. *Women and Birth: Journal of the Australian College of Midwives*, 25, 114.
- Nota, J. A., Schubert, J. R. and Coles, M. E. (2016). Sleep disruption is related to poor response inhibition in individuals with obsessive compulsive and repetitive negative thought symptoms. *Journal of Behavior Therapy and Experimental Psychiatry*, 50, 23–32.

- Obsessive Compulsive Cognitions Working Group** (2005). Psychometric validation of the obsessive belief questionnaire and interpretation of intrusions inventory – part 2: Factor analyses and testing of a brief version. *Behaviour Research and Therapy*, 43, 1527–1542.
- R Core Team** (2016). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rachman, S.** (1998). A cognitive theory of obsessions: elaborations. *Behaviour Research and Therapy*, 36, 385–401. doi: [10.1016/s0005-7967\(97\)10041-9](https://doi.org/10.1016/s0005-7967(97)10041-9)
- Rhemtulla, M., Brosseau-Liard, P. É. and Savalei, V.** (2012). When can categorical variables be treated as continuous? A comparison of robust continuous and categorical SEM estimation methods under suboptimal conditions. *Psychological Methods*, 17, 354–373. doi: [10.1037/a0029315](https://doi.org/10.1037/a0029315).
- Roscoe, J. A., Morrow, G. R., Hickok, J. T., Mustian, K. M., Griggs, J. J., Matteson, S. E. et al.** (2005). Effect of paroxetine hydrochloride (Paxil®) on fatigue and depression in breast cancer patients receiving chemotherapy. *Breast Cancer Research and Treatment*, 89, 243–249. doi: [10.1007/s10549-004-2175-1](https://doi.org/10.1007/s10549-004-2175-1)
- Rosseel, Y.** (2012). lavaan: an R package for structural equation modeling. *Journal of Statistical Software*, 48, 1–36.
- Russell, E. J., Fawcett, J. M. and Mazmanian, D.** (2013). Risk of obsessive-compulsive disorder in pregnant and postpartum women: a meta-analysis. *Journal of Clinical Psychiatry*, 74, 377–385.
- Ruscio, A. M., Stein, D. J., Chiu, W. T. and Kessler, R. C.** (2010). The epidemiology of obsessive compulsive disorder in the National Comorbidity Survey Replication. *Molecular Psychiatry*, 15, 53–63.
- Saisto, T., Salmela-Aro, K., Nurmi, J. and Halmesmäki, E.** (2001). Psychosocial characteristics of women and their partners fearing vaginal childbirth. *BJOG: An International Journal of Obstetrics and Gynaecology*, 108, 492–498.
- Salomonsson, B., Gullberg, M. T., Alehagen, S. and Wijma, K.** (2013). Self-efficacy beliefs and fear of childbirth in nulliparous women. *Journal of Psychosomatic Obstetrics and Gynecology*, 34, 116–121.
- Stoll, K., Hauck, Y., Downe, S., Edmonds, J., Gross, M. M., Malott, A. et al.** (2016). Cross-cultural development and psychometric evaluation of a measure to assess fear of childbirth prior to pregnancy. *Sexual and Reproductive Healthcare*, 8, 49–54.
- Straub, H., Adams, M., Kim, J. J. and Silver, R. K.** (2012). Antenatal depressive symptoms increase the likelihood of preterm birth. *American Journal of Obstetrics and Gynecology*, 207, 329.
- Sydsjö, G., Bladh, M., Lilliecreutz, C., Persson, A., Vyöni, H. and Josefsson, A.** (2014). Obstetric outcomes for nulliparous women who received routine individualized treatment for severe fear of childbirth – a retrospective case control study. *BMC Pregnancy and Childbirth*, 14, 1.
- Sydsjö, G., Blomberg, M., Palmquist, S., Angerbjörn, L., Bladh, M. and Josefsson, A.** (2015). Effects of continuous midwifery labour support for women with severe fear of childbirth. *BMC Pregnancy and Childbirth*, 15, 1.
- Takegata, M., Haruna, M., Matsuzaki, M., Shiraishi, M., Okano, T. and Severinsson, E.** (2015). Does antenatal fear of childbirth predict postnatal fear of childbirth? A study of Japanese women. *Open Journal of Nursing*, 5, 144.
- Wambach, K. A.** (1998). Maternal fatigue in breastfeeding primiparae during the first nine weeks postpartum. *Journal of Human Lactation*, 14, 219–229. doi: [10.1177/089033449801400311](https://doi.org/10.1177/089033449801400311)
- Williams, A. R., Williams, P. D., Press, A., Piamjariyakul, U., Keeter, L. M., Schultz, J. and Hunter, K.** (1999). Fatigue in mothers of infants discharged to the home on apnea monitors. *Applied Nursing Research*, 12, 69–77. doi: [10.1016/S0897-1897\(99\)80342-5](https://doi.org/10.1016/S0897-1897(99)80342-5)
- Winstead-Fry, P.** (1998). Psychometric assessment of four fatigue scales with a sample of rural cancer patients. *Journal of Nursing Measurement*, 6, 111.
- Wynter, K., Rowe, H. and Fisher, J.** (2013). Common mental disorders in women and men in the first six months after the birth of their first infant: a community study in Victoria, Australia. *Journal of Affective Disorders*, 151, 980–985. doi: <http://dx.doi.org.ezproxy.library.ubc.ca/10.1016/j.jad.2013.08.021>