King's Research Portal

DOI:

[10.1055/s-0036-1591552](https://doi.org/10.1055/s-0036-1591552)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Corcoran, D., Picconi, P., Chui, C. K. F., Procopiou, G., Pysz, I., Rahman, K. M., & Thurston, D. E. (2018). Formation of a Novel C11-Acetone Adduct of a Pyrrolobenzodiazepine (PBD) with Loss of Cytotoxicity. *SYNLETT*, 29(8), 1112-1116. <https://doi.org/10.1055/s-0036-1591552>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Formation of a Novel C11-Acetone Adduct of a Pyrrolobenzodiazepine (PBD) With Loss of Cytotoxicity

David B. Corcoran^{a,b}
 Pietro Picconi^a
 Connie K. Chui^a
 George Procopiou^b
 Ilona Pysz^{a,b}
 Khondaker Miraz Rahman^{a,b}
 David E. Thurston^{*a,b}

^aInstitute for Pharmaceutical Science, King's College London, Faculty of Life Sciences and Medicine, Franklin Wilkins Building, London SE1 9NH, United Kingdom.

^bFemtoGenix Ltd., Biopark, Broadwater Road, Welwyn Garden City AL7 3AX, United Kingdom.

*Indicates the main/corresponding author.

david.thurston@kcl.ac.uk or
 david.thurston@femtoGenix.com

Received:
 Accepted:
 Published online:
 DOI:

Abstract The pyrrolidine-catalysed formation of novel diastereomeric C11-acetone adducts was observed during the chromatographic purification of pyrrolobenzodiazepine (PBD) compounds in the presence of acetone. The mechanism of this reaction was explored, and the adducts obtained fully characterised. Talirine, the cytotoxic payload element of the Antibody-Drug Conjugate (ADC) vadastuximab talirine, was also found to form this adduct under similar conditions. A cellular cytotoxicity evaluation of the modified PBD compounds confirmed their lack of cytotoxicity, consistent with loss of the DNA-interactive N10-C11 imine functionality. As well as the new chemistry reported here, given the number of PBD-based ADCs presently in the clinic, this observation may be important for the larger scale manufacture of PBD-based products.

Key words Pyrrolobenzodiazepines, PBDs, Stork Enamine, DNA-Binding, Talirine, ADC.

Originally discovered in *Streptomyces* bacteria, pyrrolo[2,1-*c*][1,4]benzodiazepines (PBDs) and their derivatives are of interest as chemotherapeutic agents¹. Examples of PBD structures include anthramycin, SJG-136 and talirine (Figure 1). One PBD-based agent, SJG-136², reached Phase II clinical trials in ovarian cancer and leukaemia, and a clinical trial in canine lymphoma is still underway³. PBD dimers are also being used as cytotoxic payloads to attach to antibodies to form Antibody-Drug Conjugates (ADCs), and examples containing the PBD dimers talirine and tesirine are presently in the clinic⁴⁻⁶. The presence of an electrophilic imine/carbinolamine at the N10-C11 position of the PBD structure is regarded as essential for their biological activity. This 'soft' imine electrophile undergoes nucleophilic attack by the nucleophilic C2-NH₂ groups of guanine bases within the DNA minor groove, forming covalent adducts (Figure 2)⁷⁻¹⁰. These adducts are thought to inhibit the activity of a number of DNA processing proteins including RNA polymerase,

endonucleases and transcription factors, ultimately leading to cell death^{11,12,13, 14}.

Figure 1 Structures of Anthramycin (1), a naturally occurring PBD monomer, and the synthetic PBD dimers SJG-136 (2) and talirine (3).

Figure 2 Mechanism of binding of a PBD to the C2-amino group of a guanine base within the DNA minor groove.

In addition to the significance of the N10-C11 imine moiety of the PBD core from a biological perspective, this inherently reactive functional group can also be manipulated synthetically in order to enhance stability or alter toxicity profiles. For example, water or methanol can add to the imine to form carbinolamine (4) or carbinolamine methyl ether (5) species, respectively. The former is thought to form in aqueous solution and has been postulated

to be the ultimate biologically active species for all PBD-based molecules (Figure 3)¹⁵. The imine, carbinolamine and carbinolamine methyl ether forms have been shown to be interconvertible, and some PBDs are more stable in their methyl ether forms when stored in powder form (*e.g.*, anthramycin methyl ether¹⁶).

Recently, several groups have sought to reversibly modify the N10-C11 imine to develop more selectively targeted PBD pro-drugs. Examples include the conjugation of enzymatically cleavable agents such as sugars¹⁷ and nitroimidazole carbamate¹⁸ moieties to the N10-position of the imine. Furthermore, in the construction of ADCs containing PBD-based payloads, control of the reactivity of the N10-C11 imine is often important to improve overall stability and toxicity of the ADC. Accordingly, the antibody is sometimes connected to the PBD through a carbamate linkage to the N10-position, allowing release of the electrophilic N10-C11 imine moiety only upon carbamate cleavage¹⁹.

Irreversible modifications of the N10-C11 imine have also been reported. For example, thiol nucleophiles such as thiophenol and glutathione have also been shown to add to the C11-position of the imine functionality (*e.g.*, **6**)^{20, 21}. In the case of carbon nucleophiles, the stereoselective addition of an indole to the C11-position of a PBD core structure has been reported by Mori and co-workers²² to produce the natural product tilivalline (**7**), a good example of the potential for modification and diversification at this position. Recently, the biosynthesis of this compound by *Klebsiella oxytoca*¹⁶ has been implicated as a possible cause of human antibiotic-associated colitis²³. However, in comparison with traditional, imine-containing PBDs, **7** displays relatively weak cytotoxicity²⁴ which may derive from a non-DNA-binding mechanism. The similarly synthesised C11-cyano derivative (**8**)²⁵ appears to be more cytotoxic, and may be able to convert to the N10-C11 imine form under biological conditions, although this has not been demonstrated experimentally. Therefore, based on the above, the formation of potentially irreversible C-C bonds at the C11-position of the PBD structure has been avoided in the development of PBD-based therapeutic agents in an effort to maintain DNA-binding properties and cytotoxicity.

Here, we report that acetone can add to the C11-position of a PBD core *via* an apparent Stork enamine reaction. This was initially observed to occur during the purification of PBD compounds by column chromatography when acetone was used as a component of the solvent system. The acetone adducts are devoid of

cytotoxicity due to the loss of the DNA-interactive N10-C11 imine moiety. Therefore, the observation of this reaction has implications for the manufacture of PBD-based therapeutic agents, as well as being of chemical interest.

PBDs are mostly synthesized *via* a multi-step route starting from commercially available building blocks such as vanillin and pyrrolidinemethanol, and many typical routes have been reported and reviewed¹⁵. The N10-C11 imine moiety (in its carbinolamine form) is protected during a typical synthesis using Alloc (at the N10) and THP (at the C11) as protecting groups (*e.g.*, **9**, Scheme 1). In particular, the C11-O-THP protection avoids loss of the relatively vulnerable C11-OH group when exposed to numerous reagents during multistep synthetic routes^{1, 15}.

The final step of the synthesis involves removal of the N10-Alloc group *via* a palladium catalysed Tsuji-Trost reaction, using pyrrolidine as a nucleophile. This leads to a concerted loss of the THP group to yield the imine **10** with typical yields of 70-80% (Scheme 1).

Progress of the reaction is typically monitored through TLC or HPLC-MS and, in order to remove the palladium catalyst and other impurities, flash column chromatography is frequently used. In the observation reported here, a dichloromethane/acetone solvent system was used for the final purification of a PBD of type **9**. However, following purification of the concentrated reaction mixture after chromatography, an increase in mass was observed by HPLC-MS analysis, with two new compounds produced for which NMR analysis indicated a loss of the imine functionality. Further chromatography allowed the two products to be isolated and characterised as the diastereomeric C11-acetone adducts **16** and **17**²⁶ (Scheme 2).

This proposed Stork enamine reaction mechanism is feasible because the unpurified PBD imine was exposed to residual

pyrrolidine during the final deprotection step. The combination of acetone (used as a component of the eluent during chromatography) and pyrrolidine could generate a reactive enamine **14** which may act as a carbon nucleophile, attacking the PBD imine at the C11-position and forming intermediate **15**. Subsequent hydrolysis of the enamine would yield the final products **16** and **17** as a mixture of two diastereomers. One diastereomer (**16**) formed preferentially (*i.e.*, ~70% of total product), although it was possible to separate both diastereomers by further flash column chromatography, obtaining compounds of identical molecular weight as identified by UPLC-MS (see *S1, Supporting Information*). It proved difficult to conclusively confirm the diastereomeric ratio through NMR due to the complexity of the spectra. However, based upon COSY NMR and the observed multiplicity of the C11a-proton (ddd, $J = 10.6, 8.9, 4.9$ Hz, see *S2, Supporting Information*) in the NMR spectrum of the dominant diastereomer, it was evident that the C11(*S*)/C11a(*S*)-diastereomer forms preferentially. This would involve attack of the enamine from the most sterically-favoured direction, *trans* to the pyrrolidine ring of the PBD (see Figure 4).

Figure 4 Rationalisation of the stereochemistry of the dominant diastereomer **16** based on the most sterically-favoured approach of the nucleophile to PBD **10** ($R = \text{CH}_2\text{CH}_2\text{CH}_2\text{-COOCH}_3$). The dotted line in the Newman Projection represents the pyrrolidine ring of the PBD.

To provide more evidence for the proposed mechanism (*i.e.*, Scheme 2), pyrrolidine (0.1 equiv.) was added to the purified PBD imine **10** dissolved in acetone. After 72 hours at room temperature, conversion to **16** and **17** in good yield (67% overall) was observed (see *S3, Supporting Information*). The experiment was repeated in the presence of silica gel (0.05 equiv.) to simulate the chromatographic conditions, however no significant difference in the rate or extent of formation of **16** and **17** was observed, indicating that only pyrrolidine and acetone are required for reaction to occur.

To investigate the vulnerability of more clinically-relevant PBD compounds to enamine attack, talirine (**3**, Figure 1), a PBD-based ADC payload of significant current clinical interest⁴⁻⁶, was exposed to similar acetone/pyrrolidine conditions. Evidence of mono and *bis* acetone adducts were observed using LC-MS (see *S4, Supporting Information*). Given the asymmetric nature of this dimer molecule, up to eight possible adducts with different stereochemistries may have formed, and this mixture was too complex to resolve chromatographically. An example of a potentially preferred adduct is shown in Figure 5.

Figure 5 One of eight potential *bis*-acetone adducts of talirine (**3**) formed upon treatment with acetone/pyrrolidine. Acetone adducts may form at both N10-C11 loci in either *S* or *R* configurations.

Formation of **16** and **17** was similarly observed when piperidine was used in place of pyrrolidine, albeit in more modest yield (16%). To eliminate the possibility that this reaction proceeded through the base catalysed enolization of acetone, pyrrolidine was substituted with the hindered base 2,2,6,6-tetramethylpiperidine (TMP) of equal basicity. No formation of **16** and **17** was observed in the presence of this base, suggesting that an enamine intermediate is required for adduct formation. Further reactions were attempted using more complex ketone structures as well as diethylamine and morpholine as base (see Table 1). However, these reactions were unsuccessful, suggesting that acetone and pyrrolidine may be the optimal reagents. The lack of reactivity of the more complex ketones could be due to their increased steric bulk and/or lower nucleophilicity of their corresponding bases. For example, it has been reported that morpholine-containing enamines consistently show poorer nucleophilicity compared to their pyrrolidine-containing counterparts²⁷.

The major isolated diastereomer **16** and a 50:50 mixture of both diastereomers (**16/17**) were evaluated for cytotoxicity in a breast tumour cell line (MDA-MB-231) using the MTT assay (48 hours incubation). The unmodified imine-containing PBD **10** was also evaluated as a control (Table 2; and *S5, Supporting Information*). PBD **10** was cytotoxic in the low micromolar range, consistent with DNA-binding due to the presence of the N10-C11 imine. In contrast, the C11-modified PBD **16**, and the mixture of **16/17**, were non-cytotoxic at concentrations of up to 100 μM (the maximum concentration tested), consistent with the absence of the electrophilic N10-C11 imine moiety and an inability to react covalently with DNA. The lack of cytotoxicity for both **16** and **17** provides further evidence for the importance of the N10-C11 imine for the biological activity of PBD molecules.

Table 1 Investigation of the use of various ketones and different bases for the formation of **16** and **17**.

Ketone	Base	Adduct Formation	Yield (%)
Acetone	Pyrrolidine	Yes	67
Acetone	Piperidine	Yes	16
Acetone	TMP	No	N/A
Acetone	Diethylamine	No	N/A
Acetone	Morpholine	No	N/A
Butan-2-one	Pyrrolidine	No	N/A
Pentan-3-one	Pyrrolidine	No	N/A
Acetophenone	Pyrrolidine	No	N/A

Table 2 Cytotoxicity of the unmodified PBD (**10**) and the C11-acetone adducts (**16** and **17**) in the MDA-MB-231 breast tumour cell line.

Compound	IC ₅₀ (μM) ^a
10	6.67 (±1.08)
16	>100
16 & 17 (1:1)	>100

^aCytotoxicity toward the MDA-MB-231 breast tumour cell line measured using an MTT assay after 48 hours incubation.

In summary, the formation of C11-acetone adducts of PBD molecules is reported, most likely mediated through a Stork enamine-type mechanism. Interest in PBDs as therapeutic agents has increased in recent years due to the development of highly potent PBD-based transcription factor inhibitors¹⁴ and Antibody-Drug Conjugates (ADCs) containing PBD-based payloads^{4, 5, 28}. Therefore, the observation reported here that a biologically inactive C11-acetone adduct can form during chromatographic purification has potential implications for the manufacture of PBD-based therapeutic agents. Awareness of the possibility of this reaction may allow its prevention by using alternative amines to pyrrolidine for protecting group removal during the final synthetic steps. From a medicinal chemistry perspective, development of novel non-DNA-interactive PBD-based libraries could also be produced through this reaction for other purposes. Finally, the cytotoxicity evaluation carried out on compounds **16** and **17** provides further evidence for the importance of the N10-C11 imine functionality for the biological activity of PBDs.

Funding Information

This work was supported by a grant (12-1263/JGATCBBR) from Worldwide Cancer Research (WCR; formerly AICR) to DET, KMR and CP (2012). WCR is thanked for a PhD studentship to DBC.

Acknowledgment

Dr Julia Mantaj is thanked for providing preliminary UPLC data for acetone adducts **16** and **17**.

Supporting Information

YES

Primary Data

NO

References and Notes

- Gerratana, B., Biosynthesis, Synthesis, and Biological Activities of Pyrrolobenzodiazepines. *Medicinal Research Reviews* 2012, 32 (2), 254-293.
- Gregson, S. J.; Howard, P. W.; Hartley, J. A.; Brooks, N. A.; Adams, L. J.; Jenkins, T. C.; Kelland, L. R.; Thurston, D. E., Design, Synthesis, and Evaluation of a Novel Pyrrolobenzodiazepine

DNA-Interactive Agent with Highly Efficient Cross-Linking Ability and Potent Cytotoxicity. *J. Med. Chem.* 2001, 44 (5), 737-748.

- Mellinas-Gomez, M.; Spanswick, V. J.; Paredes-Moscossa, S. R.; Robson, M.; Pedley, R. B.; Thurston, D. E.; Baines, S. J.; Stell, A.; Hartley, J. A., Activity of the DNA minor groove cross-linking agent SG2000 (SJG-136) against canine tumours. *BMC veterinary research* 2015, 11 (1), 215.
- Beck, A.; Goetsch, L.; Dumontet, C.; Corvaia, N., Strategies and Challenges for the Next Generation of Antibody-Drug Conjugates. *Nature reviews. Drug discovery* 2017.
- Mantaj, J.; Jackson, P. J.; Rahman, K. M.; Thurston, D. E., From Anthramycin to Pyrrolobenzodiazepine (PBD)-Containing Antibody-Drug Conjugates (ADCs). *Angewandte Chemie International Edition English* 2017, 56 (2), 462-488.
- Stein, A. S.; Walter, R. B.; Erba, H. P.; Fathi, A. T.; Advani, A. S.; Lancet, J. E.; Ravandi, F.; Kovacs, T. J.; DeAngelo, D. J.; Bixby, D., A phase 1 trial of SGN-CD33A as monotherapy in patients with CD33-positive acute myeloid leukemia (AML). *Am Soc Hematology*: 2015.
- Rahman, K. M.; James, C. H.; Bui, T. T. T.; Drake, A. F.; Thurston, D. E., Observation of a Single-Stranded DNA/Pyrrolobenzodiazepine Adduct. *J. Am. Chem. Soc.* 2011, 133 (48), 19376-19385.
- Rahman, K. M.; Mussa, V.; Narayanaswamy, M.; James, C. H.; Howard, P. W.; Thurston, D. E., Observation of a Dynamic Equilibrium Between DNA Hairpin and Duplex Forms of Covalent Adducts of a Minor Groove Binding Agent. *Chemical Communications* 2009, (2), 227-229.
- Rahman, K. M.; Thompson, A. S.; James, C. H.; Narayanaswamy, M.; Thurston, D. E., The Pyrrolobenzodiazepine Dimer SJG-136 Forms Sequence-Dependent Intrastrand DNA Cross-Links and Monoalkylated Adducts in Addition to Interstrand Cross-Links. *J. Am. Chem. Soc.* 2009, 131 (38), 13756-13766.
- Rahman, K. M.; Thurston, D. E., Effect of Microwave Irradiation on Covalent Ligand-DNA Interactions. *Chemical Communications* 2009, (20), 2875-2877.
- Puvvada, M. S.; Forrow, S. A.; Hartley, J. A.; Stephenson, P.; Gibson, I.; Jenkins, T. C.; Thurston, D. E., Inhibition of bacteriophage T7 RNA polymerase in vitro transcription by DNA-binding pyrrolo [2, 1-c][1, 4] benzodiazepines. *Biochemistry* 1997, 36 (9), 2478-2484.
- Puvvada, M. S.; Hartley, J. A.; Jenkins, T. C.; Thurston, D. E., A quantitative assay to measure the relative DNA-binding affinity of pyrrolo [2, 1-c][1, 4] benzodiazepine (PBD) antitumour antibiotics based on the inhibition of restriction endonuclease BamHI. *Nucleic acids research* 1993, 21 (16), 3671-3675.
- Kotecha, M.; Kluza, J.; Wells, G.; O'Hare, C. C.; Forni, C.; Mantovani, R.; Howard, P. W.; Morris, P.; Thurston, D. E.; Hartley, J. A., Inhibition of DNA binding of the NF- κ B transcription factor by the pyrrolobenzodiazepine-polyamide conjugate GWL-78. *Molecular cancer therapeutics* 2008, 7 (5), 1319-1328.
- Rahman, K. M.; Jackson, P. J. M.; James, C. H.; Basu, B. P.; Hartley, J. A.; De La Fuente, M.; Schatzlein, A.; Robson, M.; Pedley, R. B.; Pepper, C.; Fox, K. R.; Howard, P. W.; Thurston, D. E., GC-Targeted C8-linked pyrrolobenzodiazepine-biaryl conjugates

with femtomolar in vitro cytotoxicity and in vivo antitumor activity in mouse models. *J. Med. Chem.* 2013, 56 (7), 2911-2935.

15. Antonow, D.; Thurston, D. E., Synthesis of DNA-interactive pyrrolo [2, 1-c][1, 4] benzodiazepines (PBDs). *Chemical reviews* 2010, 111 (4), 2815-2864.

16. Dornisch, E.; Pletz, J.; Glabonjat, R. A.; Martin, F.; Lembacher-Fadum, C.; Neger, M.; Hoegenauer, C.; Francesconi, K.; Kroutil, W.; Zangger, K., Biosynthesis of the enterotoxic pyrrolobenzodiazepine natural product tilivalline. *Angewandte Chemie* 2017.

17. Adiyala, P. R.; Tekumalla, V.; Sayeed, I. B.; Nayak, V. L.; Nagarajan, A.; Shareef, M. A.; Nagaraju, B.; Kamal, A., Development of pyrrolo [2, 1-c][1, 4] benzodiazepine β -glucoside prodrugs for selective therapy of cancer. *Bioorganic chemistry* 2018, 76, 288-293.

18. Dragovich, P. S.; Broccatelli, F.; Chen, J.; Fan, P.; Le, H.; Mao, W.; Pillow, T. H.; Polson, A. G.; Wai, J.; Xu, Z., Design, synthesis, and biological evaluation of pyrrolobenzodiazepine-containing hypoxia-activated prodrugs. *Bioorganic & medicinal chemistry letters* 2017, 27 (23), 5300-5304.

19. Tiberghien, A. C.; Levy, J.-N.; Masterson, L. A.; Patel, N. V.; Adams, L. R.; Corbett, S.; Williams, D. G.; Hartley, J. A.; Howard, P. W., Design and synthesis of tesirine, a clinical antibody-drug conjugate pyrrolobenzodiazepine dimer payload. *ACS Medicinal Chemistry Letters* 2016, 7 (11), 983-987.

20. Morris, S. J.; Thurston, D. E.; Nevell, T. G., Evaluation of the electrophilicity of DNA-binding pyrrolo [2, 1-c][1, 4] benzodiazepines by HPLC. *The Journal of antibiotics* 1990, 43 (10), 1286-1292.

21. Zioga, G.; Howard, P. W.; Canfield, L. M.; Gescher, A.; Thurston, D. E., Interaction of the pyrrolobenzodiazepine antitumour agent anthramycin with glutathione: A possible role in metabolism. *Pharmacy and Pharmacology Communications* 1996, 2 (1), 39-42.

22. Mori, S.; Ohno, T.; Harada, H.; Aoyama, T.; Shioiri, T., New methods and reagents in organic synthesis. 92. A stereoselective synthesis of tilivalline and its analogs. *Tetrahedron* 1991, 47 (27), 5051-5070.

23. Schneditz, G.; Rentner, J.; Roier, S.; Pletz, J.; Herzog, K. A. T.; Bucker, R.; Troeger, H.; Schild, S.; Weber, H.; Breinbauer, R., Enterotoxicity of a nonribosomal peptide causes antibiotic-associated colitis. *Proceedings of the National Academy of Sciences* 2014, 111 (36), 13181-13186.

24. Shioiri, T.; Aoyama, T.; Yamagami, N.; Shimizu, N.; Mori, N.; Kohda, K., STRUCTURE-CYTOTOXICITY RELATIONSHIP OF

TILIVALLINE DERIVATIVES. *Anti-Cancer Drug Des.* 1995, 10 (2), 167-176.

25. Kohda, K.; Yamagami, N.; Kasamatsu, T.; Aoyama, T.; Shioiri, T., Wide difference between the cytotoxicity of the 11- α - and 11- β -cyano analogues of tilivalline and their epimeric conversion. *Biochemical pharmacology* 1995, 49 (8), 1063-1068.

26. **Experimental procedure for the synthesis of 16 and 17.** Pyrrolidine (2.3 μ l, 0.1 equiv.) was added to a solution of **10** (100 mg, 1 equiv.) in acetone (1 ml), and the solution allowed to stir at room temperature for 72 hours. The excess acetone and pyrrolidine were removed under reduced pressure, and the resulting oil purified using flash column chromatography (20:80 acetone/dichloromethane to resolve the two diastereomers **16** and **17** (both yellow oils) in ~67% overall yield.

Methyl 4-(((11S,11aS)-7-methoxy-5-oxo-11-(2-oxopropyl)-2,3,5,10,11,11a-hexahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)butanoate (16; Major Diastereomer): ^1H NMR (400 MHz, CDCl_3): 7.29 (s, 1H), 6.19 (s, 1H), 4.02 (t, 2H, $J = 6.2$ Hz), 3.84 (s, 3H), 3.79-3.66 (m, 2H), 3.70 (s, 3H), 3.69-3.65 (m, 1H), 3.44 (ddd, 1H, $J = 10.6, 8.9, 4.9$ Hz), 2.57-2.52 (m, 4H), 2.24 (s, 3H) 2.19-2.07 (m, 2H), 2.05-1.92 (m, 3H), 1.75-1.64 (m, 1H). ^{13}C NMR (100 MHz, CDCl_3): 208.3, 173.6, 168.4, 151.2, 144.6, 138.3, 119.4, 113.1, 106.7, 67.8, 61.7, 60.0, 56.2, 53.7, 51.7, 46.9, 30.8, 30.3, 30.0. 24.7, 23.2.

Methyl 4-(((11R,11aS)-7-methoxy-5-oxo-11-(2-oxopropyl)-2,3,5,10,11,11a-hexahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)butanoate (17, Minor Diastereomer): ^1H NMR (400 MHz, CDCl_3): 7.45 (s, 1H), 5.97 (s, 1H), 4.05-3.99 (m, 2H), 3.95 (t, 2H, $J = 6.3$ Hz), 3.92-3.85 (m, 1H), 3.77 (s, 3H), 3.64 (s, 3H), 3.61-3.52 (m, 1H), 2.79 (dd, 1H, $J = 18.0, 10.4$ Hz), 2.51-2.44 (m, 3H), 2.29-2.18 (dt, 1H, $J = 13.2, 7.2$ Hz) 2.15-2.04 (m, 5H), 1.93-1.80 (m, 2H), 1.72-1.61 (m, 1H).

27. Kempf, B.; Hampel, N.; Ofial, A. R.; Mayr, H., Structure-Nucleophilicity Relationships for Enamines. *Chemistry-A European Journal* 2003, 9 (10), 2209-2218.

28. Jeffrey, S. C.; Burke, P. J.; Lyon, R. P.; Meyer, D. W.; Sussman, D.; Anderson, M.; Hunter, J. H.; Leiske, C. I.; Miyamoto, J. B.; Nicholas, N. D., A potent anti-CD70 antibody-drug conjugate combining a dimeric pyrrolobenzodiazepine drug with site-specific conjugation technology. *Bioconjugate chemistry* 2013, 24 (7), 1256-1263.