King's Research Portal

DOI:

[10.1002/eat.22938](https://doi.org/10.1002/eat.22938)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Matheson, E., Wade, T. D., & Yiend, J. (2018). A New Cognitive Bias Modification Technique to Influence Risk Factors for Eating Disorders. *International Journal of Eating Disorders*. <https://doi.org/10.1002/eat.22938>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Number of Words Abstract = 224
Number of Words Manuscript =4,496
Total Number of Words =4,720
Number of Tables= 3
Number of Figures=1
Number of pages=26

Revised: 17th July 2018

A New Cognitive Bias Modification Technique to Influence Risk Factors for Eating Disorders

Emily Matheson, BPsycHons¹

Tracey D. Wade, PhD¹

Jenny Yiend, PhD²

¹ School of Psychology, Flinders University, Adelaide, South Australia, Australia

² Institute of Psychiatry, King's College London

Acknowledgements and conflict of interest: The authors whose names are listed certify that they have no conflicts of interests to disclose.

Address correspondence to: Emily Matheson, School of Psychology, Flinders University, GPO Box 2100, Adelaide, South Australia, 5001, Australia; Phone +61-8-8201 7993; Email: emily.matheson@flinders.edu.au

Submitted to: International Journal of Eating Disorders

Abstract

Objective: Eating disorder psychopathology is associated with a propensity to interpret ambiguous stimuli to be negatively related to one's appearance and self-worth. The relative impact of modifying interpretation bias for these respective stimuli is unknown. Hence the main aim of the current study was to compare two cognitive bias modification protocols targeting interpretation bias (CBM-I), one focused on appearance and the other on self-worth, in terms of impacting interpretation bias, body dissatisfaction and negative affect. The appearance-based CBM-I protocol was developed for the current study. **Method:** Female university students (N=123) were randomised into one of three CBM-I conditions: appearance, self-worth or control. Immediately following a negative induction that significantly increased body dissatisfaction and negative affect, participants underwent their respective CBM-I training. **Results:** the CBM-I for appearance produced significant changes in the targeted bias, as well as significant improvements (moderate effect sizes) in appearance satisfaction, relative to the CBM-I for self-worth and control conditions. **Discussion:** The results support the usefulness of the CBM-I for appearance protocol, and suggests that this technique warrants further investigation with respect to modifying interpretation bias and risk factors associated with eating disorder psychopathology. Null effects of CBM-I for self-worth should be interpreted in light of study limitations, including the potential unsuitability of training material for young women. CBM-I for both types of interpretation bias should be evaluated in future research.

Keywords: cognitive bias modification; interpretation bias; body dissatisfaction; appearance

Cognitive theories propose that the development and maintenance of various psychopathologies can be partially attributed to the tendency to preferentially process disorder-salient stimuli above all other information types, resulting in an interpretation bias. Research consistently shows that eating disorder risk is associated with the perception of ambiguous stimuli to be negatively related to one's appearance (Brockmeyer et al., 2018; Rodgers & Dubois, 2016) and self-worth (e.g., Cooper, 2005; Cooper & Cowen, 2009; Pringle, Harmer, & Cooper 2010). For instance, a friend stating they joined a gym would be interpreted by the person at risk as evidence that they too should exercise and improve their weight and shape (interpretation bias related to appearance), or that they are lazy for not doing so (interpretation bias related to self-worth), rather than considering a more adaptive resolution (e.g., a friend's pride in their new found motivation).

The use of cognitive bias modification targeting interpretation bias (CBM-I) for therapeutic purposes aims to train individuals to adopt adaptive explanations for ambiguity. Techniques typically involve presenting individuals with a series of ambiguous scenarios that consistently yield an adaptive resolution once disambiguated. In consistently constraining individuals' interpretations to one theme, a new "production rule" is formed (Hoppitt, Mathews, Yiend &, Mackintosh, 2010; Mackintosh, Mathews, Yiend, Ridgeway, & Cook, 2006). Subsequently, CBM-I efficacy is determined by individuals' ability to apply this newly formed rule to new ambiguous information, as well as the degree to which targeted symptomatology is impacted.

Given that individuals with greater psychopathology are likely to have more difficulty in generating positive interpretations, standardized CBM-I paradigms have proven more effective relative to self-generated CBM-I (Rohrbacher, Blackwell, Holmes, & Reinecke, 2014). Four studies utilizing such standardized paradigms relevant to eating disorders exist. The first study trained a subclinical eating disorder sample to interpret emotionally

ambiguous scenarios that were consistent with either a positive/neutral, or negative self-worth (Yiend, Parnes, Shepherd, Roche, & Cooper, 2014). Both forms of CBM-I produced significant bias change congruent with the training valence. The negative CBM-I training had a significant impact on eating disorder symptomatology, with participants demonstrating a significant increase in depression and intrusive thoughts, related to weight and shape, during a mirror exposure task, as well as food restriction. Conversely, the positive/neutral form of CBM-I significantly reduced anxiety, depression and intrusive thoughts during two behavioural tasks (i.e., mirror exposure and weighing).

Two studies sought to modify interpretation bias associated with interpersonal difficulties in women with anorexia nervosa that can damage self-worth. The first study explored multiple sessions of CBM, targeting attentional (CBM-A) and interpretation bias for negative social stimuli (Cardi et al., 2015). Over a two-week period the women completed five sessions of CBM-A (direct attention toward positive social cues and away from negative cues) and CBM-I (beginning interpretation training of social-relevant scenarios). At post-intervention, the multi-session training significantly modified attentional and interpretation bias, as well as ameliorated anxiety and self-compassion. There was no impact on eating disorder symptoms. A more recent study found a single session of CBM-I to be comparably effective to a CBM-I control condition, with respects to modifying interpretation biases. Furthermore, the training had no impact on eating disorder behaviour or stress levels (Turton, Cardi, Treasurer, & Hirsch, 2018).

A final study successfully used an appearance-based CBM-I to modify social- and appearance-related interpretation bias in those with heightened body dysmorphia symptomatology (Summers & Cougle, 2016). CBM-I significantly reduced self-reported bulimia symptoms in those with high pre-treatment symptomatology; however there was no impact on drive for thinness (Summers & Cougle, 2018). No study has directly investigated

the effects of an appearance-based CBM-I on eating disorder psychopathology. An important and novel contribution of the current study is the development of a new appearance-based CBM-I protocol, which is the first approach to both *assess* and *modify* appearance-related interpretations biases.

In a review of CBM procedures, MacLeod (2012) noted that the effectiveness of CBM-I procedures beyond anxiety and depression was largely uncertain. The small body of literature summarized above, emerging since this review, suggests that CBM-I shows therapeutic potential in eating disorders. Direct comparisons of CBM-I for appearance and self-worth stimuli may indicate which protocol shows most promise, or whether both are worth pursuing, thus helping to efficiently shape future evaluations in the field. Therefore, the primary objective of the current study was to examine the effects of two CBM-I protocols, one targeting bias related to appearance and the other targeting bias related to self-worth, with respect to modifying disorder-salient bias and improving two risk factors for eating disorder psychopathology, body dissatisfaction and negative affect (Jacobi & Fittig, 2011). The cognitive-behavioural model would suggest that self-worth is a central maintaining factor of appearance concern and disordered eating (Fairburn, 2007), so we hypothesised that CBM-I for self-worth would be more effective at modifying the targeted interpretation bias, as well as improving risk factors (body dissatisfaction and negative affect), than CBM-I for appearance and control.

Method

Participants

One hundred and forty-five university students were recruited from a volunteer research pool, where participation earned course credit. To have a homogenous sample and thus increase power, the inclusion criteria required participants to be female, aged between 17

and 25, and have a body mass index (BMI) between 18.5 and 29.9 (i.e., not underweight nor obese; World Health Organisation, 2017). Of the 145 participants recruited, the data of 123 participants were included in the analyses; $n = 3$ were excluded for falling outside the age range, and $n = 19$ were excluded for not meeting weight criteria ($n = 10$ were underweight and $n = 9$ were obese). Prior to commencement, ethics approval for the study was obtained from the Human Research Ethics Committee and informed consent was obtained from each participant.

Materials

Body dissatisfaction induction

To induce body dissatisfaction and negative affect participants viewed 16 sequential images of thin women from contemporary fashion advertisements. Using a modified version of the Consumer Response Questionnaire (Mills, Polivy, Herman, & Tiggemann, 2002) to enhance comparisons, participants were instructed to compare their own appearance to that of the women they viewed on the computer screen. Using a 100-pixel VAS, participants rated their agreeability (0 being '*strongly disagree*' and 100 being '*strongly agree*') with the following statements: '*I would like my body to look like this woman's body*'; '*This woman is thinner than me*'; '*In a busy clothes shop, I would not like to try on bathing suits if this woman was also trying on bathing suits in the same change room*'. Inductions are routinely used in unselected samples to reduce variation in mood state levels (Segal & Ingram, 1994) and thus increase effect sizes for any subsequent improvements. The current induction has been shown to reliably induce body dissatisfaction and negative affect in unselected samples (Atkinson & Wade, 2012).

Cognitive bias modification targeting interpretation (CBM-I) training

The CBM-I training took the previously reported (e.g. Yiend et al., 2014) form of word completion and question tasks. Word completion tasks, first described by Matthews and

Mackintosh (2000), have shown to be effective at modifying biases in subclinical levels of anxiety and depression (Bowler et al., 2012) and eating disorders (Yiend et al., 2014). The training material used in the CBM-I for self-worth and CBM-I control conditions, was sourced from Yiend and colleagues (2014). To our knowledge, there has not been a CBM-I training specifically designed to modify appearance-related interpretation bias using the word completion task.

Therefore, using the framework described in the aforementioned studies, the authors developed the training task ‘CBM-I for appearance’. Training stimuli were informed by appearance-based feedback and rejection sensitivity scales (Altabe et al., 2004; Park, 2013; Park, Calogero, Young, & DiRaddo, 2010; Tantleff-Dunn, Thompson &, Dunn, 1995), as well as a pilot study conducted with 21 women aged between 21-27 years ($M = 24.35$, $SD = 1.33$). The women were asked to rate appearance-related terms (e.g., fit) on two 9-point Likert scales, which assessed relatedness to appearance (1 being ‘*completely unrelated*’ and 9 being ‘*completely related*’) and affective valence (1 being ‘*completely unpleasant*’ and 9 being ‘*completely pleasant*’). Based on these ratings, target words were chosen according to the degree to which they related to appearance and were positively endorsed.

The CBM-I training comprised 67 trials, each consisting of two consecutive components: an ambiguous scenario (including word completion) and a comprehension question. First, participants were presented with a 3-line ambiguous scenario, where the last word was purposely incomplete. Scenarios retained their emotional ambiguity until the final word was formed, which then disambiguated the meaning in a positive direction towards either one’s appearance or self-worth. Lastly, to reinforce the positive meaning of the disambiguated passage, participants were presented with a comprehension question that required the completion of the words ‘*Yes*’ or ‘*No*’. Control scenarios related to imperative (e.g., making a cup of tea) and declarative (e.g., facts about butterflies) knowledge and

177 retained neutrality when disambiguated. Training was delivered through an online survey
 178 program, and incorporated 4 initial practice trials. Both CBM-I for appearance (Matheson,
 179 Yiend, & Wade, 2018) and self-worth (Houlihan, Yiend, & Cooper, 2017) training materials
 180 are available via Open Science Framework. A sample training item from the three CBM-I
 181 conditions follow:

182 Appearance: “*Your friend is a very keen hiker and persuades you to join her and a*
 183 *group of friends on their next hike. You are apprehensive, given how far the hike was going*
 184 *to be. During the hike you realise that you are f-t*” (fit). The trained interpretation is
 185 reinforced by the comprehension question (with feedback given reflecting whether
 186 participants have responded ‘correctly’ or not) here: “*Are you surprised by your level of*
 187 *fitness?* ‘Correct’ answer: *No*. There are an equal number of randomly distributed ‘yes and
 188 ‘no’ responses required.

189 Self-worth: “*Your partner has been acting distant. You seek to reassure yourself that*
 190 *they are not annoyed with you for doing something wrong. You call them twice in quick*
 191 *succession. In your view you are being l-v-ng*” (loving). Comprehension question: *Are you*
 192 *too dependent on your partner?* *No*.

193 Control: “*You turn the kettle on and wait for the water to boil. You get a teabag out of the*
 194 *tin, which you put into a mug, and pour the boiling water onto the teabag. Next, you add the m- - k*”
 195 (milk). Comprehension question: *Have you made a cup of tea?* *Yes*.

196 **Similarity Rating Task**

197 The similarity ratings task (SRT) assessed modification of interpretation bias between
 198 pre- and post-training (Eysenck, Mogg, May, Richards, & Mathews, 1991; Mathews &
 199 Mackintosh, 2000). Similarly, the SRT assessing self-worth related bias was sourced from
 200 Yiend and colleagues (2014). Meanwhile, the authors developed corresponding tasks for the
 201 newly developed CBM-I for appearance condition. The SRTs comprised of two consecutive

subtasks: 1) a word completion task and 2) a recognition test. Together, the two SRTs consisted of 40 word completion scenarios (20 appearance-relevant and 20 self-worth relevant), which were separated into two parallel sets and their presentation counterbalanced between pre- and post-training.

The word completion task appeared in a similar format to that described in the CBM-I training; however the aim of the SRT was to assess, rather than to modify, biases. Thus, when the fragmented word was complete, each scenario and comprehension question retained emotional ambiguity, rather than reflecting and reinforcing positive interpretations, respectively. Further, each scenario was presented with a corresponding title, such as the “Family Christmas Card”: *Every year your mother organises a family portrait to use for Christmas cards. The photographer places you front and centre. You start to think about how many people will see the c-rd. (card). Is the card for celebrating Easter?*” N- (No).

In the recognition task, test sentences appeared beneath a title that corresponded with the previously encoded assessment scenarios. Participants were instructed to rate how similar in meaning the sentence was to the original passage on a scale between 1 (*very different*) and 4 (*very similar*). Each scenario had four corresponding test sentences; two target sentences and two foils sentences. Target items reflected either a positive or negative interpretation of the scenario. While, foil sentences were unrelated to appearance or self-worth and assessed participants’ general response bias (i.e., the tendency to respond to ambiguity in a positive or negative manner). Thus, the inclusion of both target and foil items, allowed for the distinction between modifying interpretation bias and more general priming effects of training (Mathews & Mackintosh, 2000). Test sentences were programmed to appear individually and at random. A sample set of test sentences for the “*Family Christmas Card*” scenario follow:

People will enjoy seeing your photo on the Christmas card (positive target)

People will dislike your appearance in the photo (negative target)

227 *The photographer was kind (positive foil)*

228 *The photographer was rude (negative foil)*

229 **Measures**

230 **Interpretation bias** Similarity rating scores were used to assess the changes in
 231 interpretation and response bias, using target and foil items respectively (Yiend et al, 2014).
 232 Interpretation bias indices were calculated separately for appearance and self-worth, at pre-
 233 and post-training, by subtracting the mean negative target rating from the mean positive
 234 target rating. Meanwhile, general response bias indices (mean positive foil rating minus mean
 235 negative foil rating) captured participants' tendency to respond in a more positive or negative
 236 manner. Bias scores ranged between -4 and 4, with 0 indicating no bias, and positive and
 237 negative values indicating a positive or negative interpretation bias, respectively.

238 **Trait measures.** The 9-item Body Dissatisfaction subscale from the Eating Disorder
 239 Inventory (Garner, Olmsted, & Polivy, 1983) was used to measure body dissatisfaction.
 240 Participants were asked to indicate how often the statement was true for them on a 6-point
 241 Likert scale ranging from 1 (*never*) to 6 (*always*). Responses to item numbers 3, 5, 6, 7 and 9
 242 were reverse-coded and the total score on the nine items were converted to a mean score,
 243 with higher values indicating a greater level of body dissatisfaction. The internal reliability of
 244 the questionnaire in this population was .85, and the range of corrected item-total correlations
 245 was .4 to .78.

246 Depression and anxiety were assessed using the two relevant subscales from the
 247 Depression, Anxiety and Stress Survey (Lovibond & Lovibond, 1995). Participants rated the
 248 applicability of statements as having occurred in the past week. Responses were scored on a
 249 4-point Likert scale ranging from 0 (*did not apply to me at all*) to 3 (*applied to me very much,*
 250 *or most of the time*), with higher scores indicating higher levels of depression and anxiety.
 251 Total mean subscale scores were multiplied by two for comparison to normative data, where

higher scores indicate a higher psychopathology. The internal reliability for the depression and anxiety subscales was .92 and .81, respectively, with corrected item-total correlations ranging from .55 to .83 and .3 to .65, respectively.

State measures. Visual analogue scales (VAS; Heinberg & Thompson, 1995) were used to assess participants' state level of appearance and weight satisfaction, by indicating a response to the following questions: (1) "*How satisfied do you feel about your appearance right now?*" (2) "*How satisfied do you feel about your weight right now?*" Participants indicated their level of satisfaction by dragging a slider along a 100-pixel VAS, which was fixed with two extreme values (0 indicating *extreme dissatisfaction* and 100 indicating *extreme satisfaction*). Lower scores indicated a lower level of satisfaction.

Negative affect was assessed using the Negative Affect subscale from the Positive Affect and Negative Affect Schedule (Watson, Clarke, & Tellegen, 1988). The measure was comprised of ten words relating to negative feelings (e.g., *distressed* or *jittery*) and required participants to indicate the level to which they were experiencing this feeling on a 5-point Likert scale ranging from 1 (*very slightly or not at all*) to 5 (*extremely*). Internal consistency for the Negative Affect subscale for the present study was .91, and the range of corrected item-total correlations was .6 to .81.

Procedure

The procedure is depicted in **Figure 1**. Participants attended a single session with six sequential phases lasting a total of 90 minutes. After data collection, participants were formally debriefed about the study objectives and provided with referrals for any concerns regarding body dissatisfaction.

Results

Participant characteristics and baseline measures

As shown in **Table 1** the three groups did not differ on any baseline variables. Negative affect was severely positively skewed, however results remained unchanged when inverse transformations were applied; thus original scores are reported.

Body Dissatisfaction Induction

A manipulation check was conducted using 3 (Training Group) x 2 (Time) mixed analysis of variance (ANOVA) to assess the changes in the three state variables between pre and post-induction. Across all three groups, the induction significantly exacerbated body dissatisfaction and negative affect as indicated by a main effect of Time (appearance satisfaction, $F[1, 120] = 29.16, p < .001$; weight satisfaction, $F[1, 120] = 19.61, p < .001$; negative affect $F[1, 120] = 21.66, p < .001$). There were no significant main effects of Training Group for any of the three dependent variables; nor significant interactions between Training Group and Time. After the induction, there was no difference between the three groups on any of the three variables (appearance satisfaction $F[2] = .94, p = .39$; weight satisfaction $F[2] = .76, p = .47$; negative affect $F[2] = .64, p = .53$). Hence all groups experienced commensurate changes on the outcome variables.

The impact of CBM-I on state variables

Changes in body dissatisfaction and negative affect between pre-training (i.e. post-induction) and post-training, across the three groups were assessed using 3 (Training Group) x 2 (Time) repeated measures ANOVA.

As shown in **Table 3**, appearance satisfaction was associated with a main effect of Time, with no main effect of Training Group and a significant interaction between Time and Training Group. A significant main effect of Time was observed for weight satisfaction and negative affect, no main effect of Training Group, nor was an interaction between Time and Training Group associated with either variable.

Post hoc analyses were conducted on the significant interaction between Time and Training Group for appearance satisfaction using Cohen's d within-group effect sizes and their 95% confidence intervals. Analyses revealed that CBM-I for appearance was associated with a significant medium sized improvement in appearance satisfaction, $d = .61$ [.18:1.04]; these effects were not mirrored in the CBM-I for self-worth ($d = .31$ [-.15:.77]) or CBM-I control ($d = .21$ [-.22:.64]) conditions.

Impact of CBM-I on modifying bias

Interpretation and response bias indices (calculated by collapsing the different directions of interpretation, see methods) were considered together in a three-way mixed ANOVA, to compare the specific interpretative consequences of CBM-I (indicated by responses to target items) to wider priming effects of training (indicated by responses to foil items), respectively. (Yiend et al., 2005; Yiend et al, 2014; Lee et al., 2016; Savulich et al., 2017). A mixed model ANOVA was conducted, with Training Group (CBM-I for appearance vs. CBM-I for self-worth vs. CBM-I control) as a between-subjects factor and Bias Type (target vs. foil) and Time (pre- vs. post-training) as within-subject factors.

As shown in **Table 3**, for self-worth bias indices, no main effects or three-way interaction were observed. Meanwhile, for appearance bias indices (see **Table 3**), significant main effects of Time, Bias Type and Training Group were observed. Accompanying the effects was a significant interaction between Training Group and Time. Post hoc analyses of the interaction were conducted using Cohen's d within-group effect sizes and their 95% confidence intervals. Analyses revealed a significant medium sized increase in positive bias in the CBM-I for appearance condition ($d = .72$ [.29:1.15]); however the CBM-I for self-worth ($d = .02$ [-.47:.44]) and CBM-I control ($d = .10$ [-.32:.54]) conditions were not associated with such effects. No significant three-way interaction between Time, Training

Group and Bias Type was observed, indicating no significant difference in the trajectory of interpretation vs. response bias between the three training groups over time.

Directions of change associated with CBM-I bias

Using pre- and post-training interpretation bias indices for appearance and self-worth, we examined the direction of change and whether this was congruent with the training that participants received. Of the 44 participants in the CBM-I for appearance condition, 29 (66%) showed a change in the predicted direction (i.e., increased positive interpretations towards appearance), 12 (27%) showed a change in the adverse direction (i.e., reduced positive interpretations), and 3 (7%) showed no change in bias. After removing the data of unresponsive participants (i.e. those who showed no change in bias), the difference in proportions of congruent and incongruent bias change was significant, $\chi^2(1, N = 41) = 7.05, p < .01$. For the 37 participants in the CBM-I for self-worth condition, 15 (40.5%) demonstrated a bias change congruent with training (i.e., increased positive interpretations towards self-worth), while the scores of 21 (56.8%) participants were incongruent (i.e., reduced positive interpretations) and 1 (2.7%) showed no change. The difference in proportions of congruent and incongruent bias change was not significant, $\chi^2(1, N = 36) = 1, p < .32$.

Discussion

The current research sought to comparatively examine two CBM-I approaches and their influence on modifying interpretation bias and two risk factors for eating disorder psychopathology. In contrast to the original hypothesis, results supported the newly developed CBM-I for appearance protocol, with the approach proving to be more effective at modifying the targeted bias and improving symptomatology than CBM-I for self-worth. Specifically, CBM-I increased positive-appearance related interpretations of ambiguity and produced significant medium sized improvements in appearance satisfaction. No significant

350 changes to bias or symptomatology were observed in those who completed CBM-I for self-
351 worth.

352 Our results are inconsistent with those of Yiend and colleagues (2014) who found
353 CBM-I for self-worth to be an effective approach for retraining disorder-salient bias, and
354 reducing anxiety, depression and intrusive thoughts related to appearance. The
355 inconsistencies may relate to differences in sample demographics, namely age and clinical
356 severity. Specifically, the current study used an unselected sample with varying levels of
357 psychopathology and a mean age of 19 years, relative to a subclinical sample with a mean
358 age of 29 years. Subsequently, age and clinical severity may determine the suitability of
359 training material and the degree of pre-existing bias, respectively. First, the content of the
360 self-worth training material may have been somewhat unsuitable for the younger sample.
361 Self-worth in younger female populations is likely to hinge on appearance, studies and
362 dating, as opposed to marriage, children and a full time career, which were reoccurring
363 themes in the existing self-worth training scenarios. Exposure to scenarios that one has yet to
364 experience or achieve may foster negative self-worth bias and subsequently feelings of failure
365 or discontent. In future studies of similar aged females to the present sample, researchers
366 should review the CBM-I for self-worth material and modify training scenarios to be more
367 reflective of younger female life domains. More generally, researchers using CBM should
368 place close attention to the suitability of the training content and seek to match or adapt items
369 to be as relevant as possible to the concerns of the sample. In the wider interpretation bias
370 literature the importance of the content match with the concerns of the sample has been
371 termed ‘content specificity’ and has been the subject of specific investigation in some
372 vulnerable populations (e.g. Savulich, Freeman, Shergill & Yiend, 2015). Second, prior to the
373 intervention, the current sample reported high levels of general response bias and positive
374 self-worth bias, indicating a propensity to respond to ambiguity in an optimistic manner, both

375 generally and regarding self-worth. Therefore, modifying bias in a direction that is already
376 congruent with participants' cognitions is likely to reduce the potency of the intervention.

377 An important contribution of the current research was the development of a CBM-I
378 for appearance protocol. Although there was a specific effect of CBM-I on interpretation bias
379 (target items), a similar pattern of results also emerged for general response bias (foil items).
380 Matthews and Mackintosh (2000) propose that the current assessment of bias may be
381 sensitive to experimental noise, resulting in target and foil items being equally encoded and
382 considered similar in meaning to the original message. Specifically, the text method assesses
383 bias on the assumption that the individual will consistently respond to ambiguity with one
384 form of interpretation (e.g., positive target), therefore rejecting the three alternative
385 interpretations (e.g., positive foil, negative target and negative foil). For example, an
386 interpretation such as, "*People will enjoy seeing your photo on the Christmas card*" (positive
387 target) leads to the correct rejection of the positive foil "*the photographer was kind*".
388 However, when encoding the original passage, these specific interpretations are not visible to
389 the individual. As such, more generic interpretations may have been generated and encoded
390 into memory, such as "*the photographer found me appealing*", "*I felt accepted*" or "*the*
391 *experience was enjoyable*". In this case, both the positive target and foil item would be
392 considered as accurate representations of the outcome, thus leading the individual to rate both
393 items as similar in meaning. We can conclude that we induced an interpretation bias as well
394 as a more general positive bias and our ability to distinguish between these two effects
395 represents a limitation of the current CBM-I protocol which may require further modification.

396 Current findings should be interpreted in the context of limitations additional to those
397 already mentioned. Firstly, the design did not include qualitative assessments at debriefing,
398 thus participants' awareness of the study's intentions is unknown. The impact of participants'
399 awareness of intervention intentions on CBM-I efficacy remains unclear, with some evidence

400 suggesting this knowledge enhances bias modification and symptom change (Mobini et al.,
401 2014), while others found it to hinder treatment effects (Orchard, Apetroaia, Clarke, &
402 Creswell, 2017). Future efforts should look to include quantitative and/or qualitative
403 awareness checks to determine the relationship between awareness and CBM-I efficacy.
404 Second, despite null effects of CBM-I for self-worth on bias and symptomatology (i.e.,
405 weight satisfaction and negative affect), the approach should not be considered ineffective.
406 Current findings are likely to be indicative of ceiling effects. The current unselected sample
407 were positively biased at baseline, both generally and towards self-worth, and as such
408 participants bias is likely to be less amenable to positive manipulation. These findings are
409 consistent with previous studies, which found an adaptive interpretation bias in healthy
410 populations prior to completing CBM-I training (e.g. Yiend, Savulich, Coughtrey, & Shafran,
411 2011; Hirsch & Mathews, 2000). Applying CBM-I for self-worth to a subclinical or clinical
412 sample, with maladaptive biases and higher levels of trait body dissatisfaction and negative
413 affect, may elicit changes in a positive direction (Yiend et al., 2014). Therefore, future efforts
414 should seek to compare the two CBM-I approaches in a subclinical or clinical sample to
415 determine whether the current findings were due to varying degrees of psychopathology.

416 Overall, development of a CBM-I approach that assesses and modifies appearance-
417 related interpretation bias is an important contribution to the currently limited understanding
418 of the role of CBM-I in eating disorders. Given that state variables of an unselected sample
419 were impacted after a single session of CBM-I for appearance, long term effects of the
420 approach in a subclinical population should be explored. Specifically, it would be of value to
421 investigate whether multi-session training generates a more pronounced improvement in bias
422 and symptomatology, and whether these changes persist over time.

References

- Altabe, M., Wood, K., Herbozo, S., & Thompson, J. (2004). The physical appearance ambiguous feedback scale (PAAFS): A measure for indexing body image related cognitive bias. *Body Image, 1*, 299–304. doi: 10.1016/j.bodyim.2004.05.004
- Atkinson, M. J., & Wade, T. D. (2012). Impact of metacognitive acceptance on body dissatisfaction and negative affect: Engagement and efficacy. *Journal of Consulting and Clinical Psychology, 80*(3), 416–425. doi:10.1037/a0028263
- Bowler, J. O., Mackintosh, B., Dunn, B., Mathews, A., Dalgleish, T., & Hoppitt, L. (2012). A comparison of cognitive bias modification for interpretation and computerised cognitive behaviour therapy: Effects on anxiety, depression, attentional control, and interpretive bias. *Journal of Consulting and Clinical Psychology, 80*, 1021–1033. doi:10.1037/a0029932
- Brockmeyer, T., Anderle, A., Schmidt, H., Febr, S., Wunsch-Leiteritz, W., Leiteritz, A., & Friederich, H. (2018). Body image related negative interpretation bias in anorexia nervosa. *Behaviour Research and Therapy, 104*, 69-73. doi:10.1016/j.brat.2018.03.003
- Cardi, V., Esposito, M., Bird, G., Rhind, C., Yiend, J., Schifano, S., . . . Treasure, J. (2015). A preliminary investigation of a novel training to target cognitive biases towards negative social stimuli in anorexia nervosa. *Journal of Affective Disorders, 188*, 188-193.
- Cooper, M. J. (2005). Cognitive theory in anorexia nervosa and bulimia nervosa: Progress, development and future directions. *Clinical Psychology Review, 25*, 511–531. doi:10.1016/j.cpr.2005.01.003
- Cooper, M. J., & Cowen, P. J. (2009). Negative self-beliefs in relation to eating disorder and depressive symptoms: Different themes are characteristic of the two sets of symptoms

- 448 in those with eating disorders and/or depression. *Journal of Cognitive Psychotherapy*,
449 23, 147–159. doi:10.1891/0889-8391.23.2.147
- 450 Eysenck, M. W., Mogg, K., May, J., Richards, A., & Mathews, A. (1991). Bias in
451 interpretation of ambiguous sentences related to threat in anxiety. *Journal of*
452 *Abnormal Psychology*, 100, 144–150.
- 453 Garner, D. M., Olmstead, M. P., & Polivy, J. (1983). Development and validation of a
454 multidimensional eating disorder inventory for anorexia nervosa and bulimia.
455 *International Journal of Eating Disorders*, 2(2), 15–34.
- 456 Heinberg, L. J., & Thompson, J. K. (1995). Body image and televised images of thinness and
457 attractiveness: A controlled laboratory investigation. *Journal of Social and Clinical*
458 *Psychology*, 14(4), 325–338.
- 459 Hirsch, C.R., & Mathews, A. (2000). Impaired positive inferential bias in social phobia.
460 *Journal of Abnormal Psychology*, 109, 705-712. doi: 10.1037//0021-843X.109.4.705
- 461 Hoppitt, L., Mathews, A., Yiend, J., & Mackintosh, B. (2010). Cognitive bias modification:
462 The critical role of active training in modifying emotional responses. *Behavior*
463 *Therapy*, 41(1), 73-81. doi: 10.1016/j.beth.2009.01.002
- 464 Houlihan, C., Yiend, J., & Cooper, M. (2017). Cognitive bias modification for negative self-
465 beliefs in eating disorders. Retrieved from osf.io/h3dur
- 466 Jacobi, C., & Fittig, E. (2010). Psychosocial risk factors for eating disorders. In W. Agras
467 (Ed.), *The Oxford Handbook of Eating Disorders* (pp. 123–136). New York: Oxford
468 University Press.
- 469 Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the depression anxiety stress scales*
470 (2nd ed). Sydney: Psychology Foundation of Australia.
- 471 MacLeod, C., Mathews, A., & Tata, P. (1986). Attentional bias in emotional disorders.
472 *Journal of Abnormal Psychology*, 95(1), 15–20. doi:10.1037/0021-843X.95.1.15

- 473 Matheson, E., Yiend, J., & Wade, T. (2018). Cognitive bias modification for appearance
474 technique for risk factors for eating disorders. Retrieved from osf.io/tzyfp
- 475 Mathews, A., & Mackintosh, B. (2000). Induced emotional interpretation bias and anxiety.
476 *Journal of Abnormal Psychology*, 109, 602–615. doi: 10.1037/0021-843X.109.4.602
- 477 Mills, J. S., Polivy, J., Herman, C. P., & Tiggemann, M. (2001). Effects of exposure to thin
478 media images: Evidence of self-enhancement among restrained eaters. *Personality
479 and Social Psychology Bulletin*, 28, 1687–1699.
- 480 Mobini, S., Mackintosh, B., Illingworth, J., Gega, L., Langdon, P., & Hoppitt, L. (2014).
481 Effects of standard and explicit cognitive bias modification and computer-
482 administered cognitive-behaviour therapy on cognitive biases and social anxiety.
483 *Journal of Behavior Therapy and Experimental Psychiatry*, 45(2), 272–279. doi:
484 10.1016/j.jbtep.2013.12.002
- 485 Orchard, F., Apetroaia, A., Clarke, K., Creswell, C. (2017). Cognitive bias modification of
486 interpretation in children with social anxiety disorder, *Journal of Anxiety Disorders*,
487 45, 1-8. doi:doi.org/10.1016/j.janxdis.2016.10.012.
- 488 Park, L. E. (2013). Appearance-based Rejection Sensitivity Scale [Measurement instrument].
489 Retrieved from [http://www.midss.org/content/appearance-based-rejection-sensitivity-
490 appearance-rs-scale-long-and-short-form](http://www.midss.org/content/appearance-based-rejection-sensitivity-appearance-rs-scale-long-and-short-form)
- 491 Park, L. E., Calogero, R. M., Young, A. F., & DiRaddo, A. (2010). Appearance-based
492 rejection sensitivity predicts body dysmorphic disorder symptoms and cosmetic
493 surgery acceptance. *Journal of Social and Clinical Psychology*, 29(5), 489-509
- 494 Pringle, A., Harmer, C. J., & Cooper, M. J. (2010). Biases in emotional processing are
495 associated with vulnerability to eating disorders over time. *Eating Behaviours*, 12,
496 56–59.doi:[10.1017/S0033291709990778](https://doi.org/10.1017/S0033291709990778)

- 497 Rodgers, R. F., & DuBois, R. H. (2016). Cognitive biases to appearance-related stimuli in
498 body dissatisfaction: A systematic review. *Clinical Psychology Review*, 46, 1-11.
499 doi:10.1016/j.cpr.2016.04.006.
- 500 Rohrbacher, H., Blackwell, S. E., Holmes, E. A., & Reinecke, A. (2014). Optimizing the
501 ingredients for imagery-based interpretation bias modification for depressed mood: Is
502 self-generation more effective than imagination alone? *Journal of Affective Disorders*,
503 212–218. doi: 10.1016/j.jad.2013.09.013
- 504 Savulich, G., Freeman, D., Shergill, S., & Yiend, J (2015). Interpretation biases in paranoia.
505 *Behavior Therapy*, 46, 110–124. doi:10.1016/j.beth.2014.08.002.
- 506 Savulich, G., Shergill, & Yiend, Y. (2017). Interpretation biases in clinical paranoia. *Clinical*
507 *Psychological Science*, 5(6), 985-1000. doi:10.1177/2167702617718180
- 508 Segal, Z. V., & Ingram, R. E. (1994). Mood priming and construct activation in tests of
509 cognitive vulnerability to unipolar depression. *Clinical Psychology Review*, 14, 663–
510 695. doi: 10.1016/0272-7358(94)90003-5
- 511 Summers, B. J. and Coughle, J. R. (2016). Modifying interpretation biases in body dysmorphic
512 disorder: Evaluation of a brief computerized treatment. *Behaviour Research and*
513 *Therapy*, 87, 117–127. doi: 10.1016/j.brat.2016.09.005
- 514 Summers, B., & Coughle, J. (2018). Effects of an Appearance-Focused Interpretation Training
515 Intervention on Eating Disorder Symptoms. *Behavioural and Cognitive*
516 *Psychotherapy*, 1-14. doi:10.1017/S1352465818000164
- 517 Tantleff-Dunn, S., Thompson, K. T., & Dunn, M. E. (1995). The feedback on physical
518 appearance scale (FOPAS): Questionnaire development and psychometric evaluation.
519 *Eating Disorders*, 3(4), 332-341, dio:10.1080/10640269508250063
- 520 Turton, R., Cardi, V., Treasure, J., Hirsch, C. R. (2018). Modifying a negative interpretation
521 bias for ambiguous social scenarios that depict the risk of rejection in women with

- 522 anorexia nervosa. *Journal of Affective Disorders*, 227, 705-712. doi:
- 523 10.1016/j.jad.2017.11.089
- 524 Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief
- 525 measures of positive and negative affect: The PANAS Scales. *Journal of Personality*
- 526 *and Social Psychology*, 54(6), 1063–1070
- 527 Williamson, D. A., Muller, S. L., Reas, D. L., Thaw, J.M. (1999). Cognitive bias in eating
- 528 disorders: implications for theory and treatment. *Behavior Modification*, 23(4), 556-
- 529 577. doi: 10.1177/0145445599234003
- 530 World Health Organisation (2017). *Global Database on Body Mass Index*. Retrieved from
- 531 http://apps.who.int/bmi/index.jsp?introPage=intro_3.html
- 532 Yiend, J., & Mathews, A. (2004). Biases in attention: Methods, mechanisms and meaning. In
- 533 A. Wenzel & D. C. Rubin (Eds.), *Cognitive methods in clinical research* (pp. 65–71).
- 534 Washington, DC: APA Books
- 535 Yiend, J., Parnes, C., Shepherd, K., Roche, M. K., & Cooper, M. J. (2014). Negative self-
- 536 beliefs in eating disorders: A cognitive bias modification Study. *Clinical*
- 537 *Psychological Science*, 2(6), 756-766.
- 538 Yiend, J., Savulich, G., Coughtrey, A., & Shafran, R. (2011). Biased interpretation in
- 539 perfectionism and its modification. *Behaviour Research and Therapy*, 49(12), 892-
- 540 900. doi: 10.1016/j.brat.2011.10.004
- 541
- 542
- 543
- 544
- 545
- 546

Figure 1. Study Design

567 Table 1

568 *Baseline demographics and dependent variables*

Variable	CBM-I Appearance (<i>n</i> = 44)		CBM-I Self-Worth (<i>n</i> = 37)		CBM-I Control (<i>n</i> = 42)		<i>Main effect of Group</i>	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>F</i> (2, 122)	<i>p</i>
Age	19.55	1.61	19.49	1.33	19.21	1.41	.62	.54
Body mass index	22.70	2.82	23.08	2.98	22.47	2.67	.46	.63
Body dissatisfaction	3.63	1.00	3.67	.92	3.52	.88	.26	.77
Anxiety	1.35	1.11	1.43	1.16	1.61	1.21	.57	.57
Depression	1.25	1.21	1.44	1.34	1.72	1.40	1.41	.25
Appearance satisfaction	49.41	23.95	45.43	25.14	52.88	26.17	.87	.42
Weight satisfaction	46.11	31.35	44.91	26.40	53.31	27.25	1.03	.36
Negative affect	1.33	.45	1.48	.53	1.49	.72	1.06	.35
Appearance interpretation bias	.15	.79	-.07	.69	-.12	.72	1.77	.17
Response bias (appearance-related foils)	.55	.43	.40	.42	.43	.52	1.20	.30
Self-worth interpretation bias	.48	.58	.43	.62	.25	.49	1.99	.14
Response bias (self-worth related foils)	.45	.58	.42	.47	.35	.48	.39	.68

569 *Note.* CBM-I = Cognitive bias modification targeting interpretation bias

570 Table 2

571

572 *Mean (standard deviation) state variables and raw target and foil scores across condition pre- and post CBM-I training*

	CBM-I Appearance		CBM-I Self-Worth		CBM-I Control	
	<i>Pre-Training M (SD)</i>	<i>Post-Training M(SD)</i>	<i>Pre-Training M(SD)</i>	<i>Post-Training M(SD)</i>	<i>Pre-Training M(SD)</i>	<i>Post-Training M(SD)</i>
Appearance satisfaction	39.84 (27.18)	55.75 (29.21)	37.05 (25.26)	45.14 (24.76)	44.95 (25.74)	50.36 (25.25)
Weight satisfaction	38.31 (30.90)	51.62 (32.59)	40.16 (26.19)	46.27 (24.66)	45.57 (27.00)	52.02 (25.04)
Negative affect	1.55(.12)	1.22 (.09)	1.74 (.13)	1.49 (.10)	1.61 (.12)	1.36 (.09)
Appearance Bias						
Positive target	2.58 (.45)	2.87 (.39)	2.47 (.38)	2.51 (.30)	2.46 (.35)	2.48 (.33)
Negative target	2.42 (.49)	2.17 (.51)	2.54 (.47)	2.53 (.50)	2.58 (.47)	2.36 (.56)
Positive foil	2.54 (.43)	2.64 (.39)	2.45 (.31)	2.45(.31)	2.38 (.38)	2.50 (.35)
Negative foil	1.99 (.32)	1.89 (.34)	2.05 (.36)	2.08 (.42)	1.95 (.45)	1.95 (.44)
Self-worth Bias						
Positive target	2.96 (.36)	3.0 (.35)	2.94 (.30)	2.86 (.41)	2.81 (.39)	2.85 (.34)
Negative target	2.49 (.38)	2.45 (.38)	2.51 (.50)	2.51 (.42)	2.56 (.48)	2.45 (.40)
Positive foil	2.52 (.41)	2.49 (.43)	2.47 (.28)	2.52 (.31)	2.38 (.38)	2.41 (.36)
Negative foil	2.07 (.35)	2.07 (.32)	2.06 (.40)	2.16 (.32)	2.02 (.36)	2.04 (.48)

573 *Note.* CBM-I = Cognitive bias modification targeting interpretation bias; Pre-training = post-induction assessment of outcome variables.

574 Table 3

575

576 *Two-way and Three-way ANOVA: Impact of CBM-I on State Variables and Bias between Pre- and Post-Training*

		<i>df</i>	<i>F</i>	<i>p</i>
State variables				
<i>Appearance satisfaction</i>				
	Time	1, 120	45.42	<.001
	CBM-I Training Group	2, 120	.90	.41
	Time × CBM-I Training Group	2, 120	4.95	.01
<i>Weight satisfaction</i>				
	Time	1, 120	35.10	<.001
	CBM-I Training Group	2, 120	.45	.64
	Time × CBM-I Training Group	2, 120	2.68	< .07
<i>Negative affect</i>				
	Time	1, 120	37.35	<.001
	CBM-I Training Group	2, 120	1.34	.27
	Time × CBM-I Training Group	2, 120	.42	.66
Bias Change Variables				
<i>Change in Bias for Appearance</i>				
	Time	1, 120	17.75	<.001
	Bias Type	1, 120	56.69	<.001
	CBM-I Training Group	2, 120	7.32	.001
	Time × Bias Type	1, 120	8.03	.01
	Time × CBM-I Training Group	2, 120	5.55	.01
	Bias Type × CBM-I Training Group	2, 120	2.69	.07
	Time × Bias Type × CBM-I Training Group	2, 120	1.79	.17
<i>Change in Bias for Self-Worth</i>				
	Time	1, 120	.05	.83
	Bias Type	1, 120	.18	.68
	CBM-I Training Group	2, 120	1.42	.25
	Time × Bias Type	1, 120	1.85	.18
	Time × CBM-I Training Group	2, 120	.75	.48
	Bias Type × CBM-I Training Group	2, 120	.75	.47
	Time × Bias Type × CBM-I Training Group	2, 120	.71	.50

577 *Note.* CBM-I = Cognitive bias modification targeting interpretation bias; Pre-training = post-induction assessment of outcome variables. Time =
578 Pre- and Post-training; Bias Type = Target and Foils; CBM-I Training Group = CBM-I for appearance, CBM-I for self-worth, CBM-I control