King's Research Portal

DOI:

[10.1016/j.euroneuro.2020.10.007](https://doi.org/10.1016/j.euroneuro.2020.10.007)

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Salazar de Pablo, G., De Micheli, A., Nieman, D. H., Correll, C. U., Kessing, L. V., Pfennig, A., Bechdolf, A., Borgwardt, S., Arango, C., Van Amelsvoort, T., Vieta, E., Solmi, M., Oliver, D., Catalan Alcantara, A., Verdino, V., Di Maggio, L., Bonoldi, I., Vaquerizo Serrano, J. D., Provenzani, U., ... Fusar-Poli, P. (2020). Universal and selective interventions to prevent poor mental health outcomes in young people: systematic review and meta-analysis. *European Neuropsychopharmacology*. <https://doi.org/10.1016/j.euroneuro.2020.10.007>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

UNIVERSAL AND SELECTIVE INTERVENTIONS TO PROMOTE GOOD MENTAL HEALTH IN YOUNG PEOPLE: SYSTEMATIC REVIEW AND META-ANALYSIS

Gonzalo Salazar de Pablo, MD^{1,2,3,#}, Andrea De Micheli, MD^{1,4,#}, Dorien H. Nieman, PhD⁵, Christoph U. Correll, MD⁶⁻⁹, Lars Vedel Kessing, MD, DMSc¹⁰, Andrea Pfennig, MD, PhD¹¹, Andreas Bechdolf, MD, PhD¹²⁻¹⁵, Stefan Borgwardt, MD, PhD¹⁶⁻¹⁷, Celso Arango, MD, PhD³, van Amelsvoort T, MD, PhD¹⁸, Eduard Vieta, MD, PhD¹⁹, Marco Solmi, MD, PhD^{1,20}, Dominic Oliver, MSc¹, Ana Catalan, MD, PhD^{1,21}, Valeria Verdino, MD^{1,22}, Lucia Di Maggio, MD^{1,20}, Ilaria Bonoldi, MD, PhD²³, Julio Vaquerizo-Serrano, MD^{2,23}, Ottone Baccaredda Boy, MD²³, Umberto Provenzano, MD¹, Francesca Ruzzi, MD²⁰, Federica Calorio, MD²⁰, Guido Nosari, MD²⁰, Benedetto Di Marco, MD²⁰, Irene Famularo, MD²⁰, Silvia Molteni, MD²⁰, Eleonora Filosi, MD²², Martina Mensi, MD²², Umberto Balottin, MD²², Pierluigi Politi, MD²², Jae Il Shin, MD, PhD²⁴, Paolo Fusar-Poli, MD, PhD^{1,4,20,25}

¹Early Psychosis: Interventions and Clinical-detection (EPIC) Lab, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK;

²Department of Child and Adolescent Psychiatry, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK;

³Institute of Psychiatry and Mental Health. Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón School of Medicine, Universidad Complutense, Instituto de Investigación Sanitaria Gregorio Marañón (IISGM), CIBERSAM, Madrid, Spain;

⁴National Institute for Health Research, Maudsley Biomedical Research Centre, South London and Maudsley NHS Foundation Trust, London, UK;

⁵Amsterdam University Medical Centers (location AMC), Department of Psychiatry, Amsterdam, the Netherlands;

⁶Department of Psychiatry, The Zucker Hillside Hospital, Northwell Health, Glen Oaks, NY, USA

⁷Department of Psychiatry and Molecular Medicine, Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Hempstead, NY, USA;

⁸Center for Psychiatric Neuroscience; The Feinstein Institutes for Medical Research, Manhasset, NY, USA;

⁹Department of Child and Adolescent Psychiatry, Charité Universitätsmedizin, Berlin, Germany

¹⁰Copenhagen Affective Disorder Research Center (CADIC), Psychiatric Center Copenhagen, Rigshospitalet, Copenhagen, Denmark;

¹¹Department of Psychiatry and Psychotherapy, Carl Gustav Carus University Hospital, Medical Faculty, Technische Universität Dresden, Germany;

¹²Department of Psychiatry, Psychotherapy and Psychosomatic Medicine with Early Intervention and Recognition Centre (FRITZ), Vivantes Klinikum Am Urban, Charité-Universitätsmedizin, Berlin, Germany;

¹³Vivantes Klinikum im Friedrichshain, Department of Psychiatry, Psychotherapy and Psychosomatic Medicine, Charité-Universitätsmedizin, Berlin, Germany;

¹⁴Department of Psychiatry and Psychotherapy, University of Cologne, Cologne, Germany

¹⁵ORYGEN, The National Centre of Excellence in Youth Mental Health, Melbourne, Victoria, Australia;

¹⁶Department of Psychiatry (UPK), University of Basel, Switzerland;

¹⁷Department of Psychiatry and Psychotherapy, University of Lübeck, Germany;

¹⁸Department of Psychiatry and Neuropsychology, Maastricht University, Maastricht, The Netherlands;

¹⁹Hospital Clinic, institute of Neuroscience, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain;

²⁰Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy;

²¹Mental Health Department. Biocruces Bizkaia Health Research Institute. Basurto University Hospital. Facultad de Medicina y Odontología, Campus de Leioa, University of the Basque Country, UPV/EHU, Spain;

²²Department of Molecular and Developmental Medicine, Division on Psychiatry, University of Siena, Italy;

²³Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK;

²⁴Department of Paediatrics, Yonsei University College of Medicine, Seoul, Republic of Korea;

²⁵OASIS service, South London and Maudsley NHS Foundation Trust, London, UK.

These authors have contributed equally and share the first position authorship

Number of words: 3816

Abstract: 250

References: 58

Tables: 1

Figures: 4

Supplementary tables: 8

Supplementary text: 1

Correspondence to: Dr Paolo Fusar-Poli MD PhD, Department of Psychosis Studies, 5th Floor, Institute of Psychiatry, Psychology & Neuroscience, PO63, 16 De Crespigny Park, SE5 8AF London, UK. E-mail: paolo.fusar-poli@kcl.ac.uk

Keywords: Prevention, promotion, good mental health, outcomes

ABSTRACT

Promotion of good mental health in young people is important. Our aim was to evaluate the consistency and magnitude of the efficacy of universal/selective interventions to promote good mental health. A systematic PRISMA/RIGHT-compliant meta-analysis (PROSPERO: CRD42018088708) search of Web of Science until 04/31/2019 identified original studies comparing the efficacy of universal/selective interventions for good mental health vs a control group, in samples with a mean age <35 years. Meta-analytical random-effects model, heterogeneity statistics, assessment of publication bias, study quality and sensitivity analyses investigated the efficacy (Hedges' g =effect size, ES) of universal/selective interventions to promote 14 good mental health outcomes defined a-priori. 276 studies were included (total participants:159,508, 79,142 interventions and 80,366 controls), mean age=15.0 (SD=7.4); female=56.0%. There was a significant overall improvement in 10/13 good mental health outcome categories that could be meta-analysed: compared to controls, interventions significantly improved (in descending order of magnitude) mental health literacy (ES=0.685, $p<0.001$), emotions (ES=0.541, $p<0.001$), self-perceptions and values (ES=0.49, $p<0.001$), quality of life (ES=0.457, $p=0.001$), cognitive skills (Hedge's g =0.428, $p<0.001$), social skills (ES=0.371, $p<0.001$), physical health (ES=0.285, $p<0.001$), sexual health (ES=0.257, $p=0.017$), academic/occupational performance (ES=0.211, $p<0.001$) and attitude towards mental disorders (ES=0.177, $p=0.006$). Psychoeducation was the most effective intervention for promoting mental health literacy (ES=0.774, $p<0.001$) and cognitive skills (ES=1.153, $p=0.03$). Physical therapy, exercise and relaxation were more effective than psychoeducation and psychotherapy for promoting physical health (ES=0.498, $p<0.001$). In conclusion, several universal/selective interventions can be effective to promote good mental health in young people. Future research should consolidate and extend these findings.

Keywords: Promotion, Good mental health, Universal, Selective, Intervention, Outcomes.

INTRODUCTION

Mental health issues are expected to become one of the five more common causes of dismallness, mortality and dysfunction among youths (Gore et al., 2011). The worldwide prevalence of mental disorders in children and adolescents is estimated to be 13.4%: anxiety disorders 6.5%, disruptive disorders 5.7%, attention-deficit hyperactivity disorder 3.4% and depressive disorders 2.6% (Polanczyk et al., 2015). Besides, mental disorders in young people have a great economic burden on society (Fatori et al., 2018)". These facts highlight the importance of preventing mental disorders, but also of promoting good mental health in young people. Good mental health is defined as a state of well-being that allows individuals to cope with the normal stresses of life and function productively (Fusar-Poli et al., 2019b). While the prevention of mental disorders in young people has entered clinical practice in psychiatry thanks to the psychosis prevention field (Fusar-Poli et al., 2013; Fusar-Poli et al., 2016), promotion of mental health has received less empirical research attention, mostly because research has been fragmented (Fusar-Poli et al., 2019b). Core domains of good mental health have recently been defined and encompass the 14 areas of mental health literacy, attitude towards mental disorders, self-perceptions and values, cognitive skills, academic/occupational performance, emotions, behaviours, self-management strategies, social skills, family and significant relationships, physical health, sexual health, meaning of life, and quality of life (Fusar-Poli et al., 2019b). Classification of interventions parallel those employed to classify the prevention of mental health; universal, selective or indicated, depending on the individuals they target (World Health Organization, 2004). Out of these, universal interventions target the general public or a whole population group that has not been identified on the basis of increased risk (World Health Organization, 2004) and selective interventions target individuals or subgroups of the population whose risk of developing a mental disorder is significantly higher than average, as evidenced by biological, psychological or social risk factors (Arango et al., 2018; World Health Organization, 2004). Both universal and selective interventions are particularly suited to promote good mental health (Fusar-Poli et al., 2019b). Indicated preventions, on the other hand, target high-risk people who are identified as having minimal but detectable signs or symptoms (including prodromal help-seekers and clinical high risk populations) foreshadowing mental disorder or biological markers indicating predisposition for mental disorder but do not meet diagnostic criteria for the disorder (World Health Organization, 2004). Preventive interventions in this group were not the focus of the current review.

The consistency and magnitude of available and potential interventions to promote mental health are patchy and conflicting. Different interventions have been tried to promote good mental health in healthy individuals, most frequently psychoeducation (Strunk et al., 2014; Witvliet et al., 2009) and psychotherapy (Collins et al., 2014; World Health Organization, 2002) and less frequently physical therapy (Sharp and Caperchione, 2016b), animal-assisted therapy (Pendry et al., 2014) and art therapy (Rousseau et al., 2009). To date, the most effective interventions for each of the core determinants of good mental health remain unknown (Fusar-Poli et al., 2019b). This study aimed to systematically review and meta-analyse the consistency and magnitude of the efficacy of universal and selective interventions to promote good mental health in young people compared to the control group, using a-priori operationalisation of good mental health.

METHODS

Protocol and registration

The study (study protocol registered on PROSPERO - CRD42018088708-), was conducted in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (Moher et al., 2009) (eTable I) and MOOSE checklist (eTable II) (Stroup et al., 2000), following EQUATOR Reporting Guidelines (Altman et al., 2008).

Eligibility criteria

Inclusion criteria were a) individuals with an average age <35 years, no lower age limit, in line with the upper age limit of the early psychosis paradigm (Fusar-Poli et al., 2019a) b) reporting on universal interventions or selective interventions (see definitions of classification of preventive approaches in psychiatry in eTable III (Fusar-Poli et al., 2019a)), c) with an intervention and a control group (randomized or non-randomized, and employing either double-blind, open-label or cross-over design), d) focusing on good mental health outcomes as defined in Table 1, with meta-analysable data (e.g. raw mean values and standard deviation, SD, before and after the intervention, Risk Ratios, Odds Ratios or T-Students and direction of effects), e) original studies written in English language.

Exclusion Criteria were a) individuals with a mean age ≥ 35 years, b) interventions conducted in patients with a known mental disorder or presenting with subthreshold psychiatric conditions (clinical or subclinical, indicated interventions), c) lack of control

group d) reporting other outcomes e) abstracts, conference proceedings, pilot data and reviews f) studies written in a language other than English. Overlap between studies was actively searched using the intervention name and content, the type of selected population and country in which the study was carried out. When overlap was found, we selected the study with the largest sample.

Search and study selection

A systematic search strategy to identify relevant articles, and a two-step literature search using the following keywords: “mental health” AND “promotion” was implemented by independent researchers (GSP, MS, OBB). As a first step, the Web of Science database (Clarivate Analytics) was searched, incorporating the Web of Science Core Collection, the BIOSIS Citation Index, the KCI-Korean Journal Database, MEDLINE®, the Russian Science Citation Index, and the SciELO Citation Index, from inception until 31st April 2019. Then, references of the articles retrieved during the literature search were searched manually (GSP, AdM, VV).

Data collection process

At least two researchers (VV, LdM, OBB, UP, FR, FC, GN, BdM, IF, SM, EF, MM) independently extracted data from all the included studies, into an electronic database. The database was then crosschecked by a third author (GSP, AC), and any discrepancies were resolved through consensus under the supervision of a senior researcher (PFP).

Data items

Variables included in the study were: study ID; country; type of primary intervention (universal, selective); intervention content (psychoeducation, psychotherapy, other, combined); intervention name; control group (active control, no intervention, treatment as usual); sample size; sex (% male); age (mean, SD); quality assessment (see below); outcomes.

Good mental health outcomes

A detailed discussion of 14 relevant outcome categories and outcomes within promotion of mental health is fully provided in a complementary publication by a group of experts (Table 1) (Fusar-Poli et al., 2019b). When >1 measure from the same mental health outcome was available in the same study, we extracted the measure most frequently

investigated across the existing literature. The study with the largest sample size was selected when the previous criterion was not discerning.

Risk of bias (quality) assessment

As per protocol, we rated randomization according to the following scoring system: 0 for studies with no randomization or in which this was not mentioned; 1 for studies in which there was some randomization, but the type was not specified; 2 for studies with clear method of randomization. We rated blinding according to the following scoring system: 0 for studies without blinding or in which this was not mentioned/specified; 1 for studies with single blinding; 2 for studies with double or triple blinding. We decided to use a simpler score in front of "Cochrane Collaboration's Tool for assessing risk of bias in randomised trials" because it would have been difficult to apply it to the current database.

Strategy for data synthesis, summary measures and analyses

The primary effect size (ES, Hedge's g) was estimated when three or more studies were available, with positives values reflecting an improvement at follow-up in the intervention group compared to the control group and negative values indicating a superior improvement in the control group. The Hedges' g is obtained with the difference between the means of the patient and control groups divided by the standard deviation (or from associated measures) and weighted for sample size. The meta-analysis included overall summary effects stratified by the different types of good mental health outcomes and sensitivity analyses, with each subgroup defined by the type of intervention that was provided (psychoeducation vs psychotherapy vs others vs combined interventions), by the type of primary intervention (universal vs selective) and by the age period (children and adolescents vs adults). Heterogeneity among study point estimates was assessed using the Q statistic, with the proportion of the total variability in ES estimates being evaluated using the I^2 index (with an $I^2 > 50\%$ representing significant heterogeneity) (Lipsey and Wilson, 2000). We used a random-effects model as heterogeneity was expected to be high (DerSimonian and Laird, 1986). The presence of publication bias in the results was assessed informally by visually inspecting funnel plots, complemented by the "trim and fill" method to investigate the effects of any publication bias that was detected. We further conducted meta-analytical regression analyses to estimate the association between the quality of the included studies and the effectiveness of the interventions. For these meta-regression analyses, scores from both randomization and blinding were added, resulting in a total value ranging from 0-4. Analyses were performed

with Comprehensive Meta-Analysis Version 3 (Biostat, Englewood, NJ, USA) two-sided and with $\alpha=0.05$.

RESULTS

Study selection

The literature search yielded 9,198 citations, which were screened for eligibility; 690 articles were assessed for eligibility and 276 studies were finally included in the meta-analysis (Figure 1 PRISMA flowchart, eTable IV main characteristics of included studies).

Study characteristics

Outcomes were targeted as follows: 45 (16.3%) articles focused on mental health literacy, 16 (5.8%) on attitude towards mental disorders, 92 (33.3%) on self-perceptions and values, 20 (7.2%) on cognitive skills, 30 (10.9%) on academic/occupational performance, 37 (13.4%) on emotions, 11 (4.0%) on behaviours, 86 (31.2%) on self-management strategies, 94 (34.1%) on social skills, 45 (16.3%) on family and significant relationships, 25 (9.1%) on physical health 13 (4.7%) on sexual health and 48 (17.4%) on quality of life (eTable V-VII). We did not find any studies on the promotion of meaning of life in young individuals, which is why we could not perform a meta-analysis on this 14th outcome domain. The type of intervention was universal in 171 (61.9%) studies, selective in 104 (37.7%) studies and both universal and selective in 1 (0.4%) (Hiscock et al., 2018) study. The overall database included 159,508 individuals (mean age range=0-34.3 years; female=56.0%, 79,142 interventions and 80,366 controls). The total sample size of the studies included ranged from 13 (Attwood et al., 2012) to 8,989 (Shinde et al., 2018).

Synthesis of results: Interventions to promote good mental health outcomes

Interventions were effective to promote mental health literacy ($k=45$, $n=37,533$; $ES = 0.685$; $p<0.001$), emotions ($k=37$, $n=7,593$ $ES=0.541$; $p<0.001$), self-perceptions and values ($k=92$, $n=37,183$ $ES=0.49$; $p<0.001$), quality of life ($k=48$, $n=31,276$; $ES=0.457$; $p=0.001$), cognitive skills ($k=20$, $n=12,368$; $ES=0.428$; $p<0.001$), social skills ($k=94$, $n=62,274$ $ES=0.371$; $p<0.001$), physical health ($k=25$, $n=7,642$; $ES=0.285$; $p<0.001$), sexual health ($k=13$, $n=14,202$; $ES=0.257$; $p=0.017$) academic/occupational performance ($k=30$, $n=19,324$ $ES=0.211$; $p<0.001$) and attitude towards mental disorders ($k=16$, $n=8,741$; $ES=0.177$; $p=0.006$), vs control conditions (eTable V-VII and

Figure 2). There was no evidence that interventions could significantly improve behaviours ($k=11$, $n=10,474$; $ES=0.065$; $p=0.23$) or family and significant relationships ($k=45$, $n=32,639$; $ES= -0.046$; $p=0.674$) (eTable V-VII and Figure 2). Once publication bias was corrected with the trim and fill method, self-management strategies became nonsignificant ($k=86$, $n=41,437$; $ES=0.517$; $p<0.001$ corrected to $ES=0.107$; $p=0.09$).

Additional analysis/ Sensitivity analyses: types of interventions and preventive approach

Among available comparisons, there were no differences in efficacy among the different intervention categories to improve emotions ($p=0.729$), self-perceptions and values ($p=0.309$), quality of life ($p=0.341$), social skills ($p=0.182$), sexual health ($p=0.718$), academic/occupational performance ($p=0.234$), behaviours ($p=0.622$), family and significant relationships ($p=0.327$) or self-management strategies ($p=0.281$). Psychoeducation was the most effective intervention for the promotion of mental health literacy ($p<0.001$) and cognitive skills ($p<0.001$). Physical therapy, exercise and relaxation were the most effective type of interventions for the promotion of physical health ($p<0.001$). All the interventions to improve attitude towards mental disorders consisted on psychoeducation, so differences could not be evaluated (eTable V, Figure 3).

For cognitive skills, universal interventions fared better than selective interventions ($p<0.001$). There were no differences between universal and selective interventions for mental health literacy ($p=0.699$), attitude towards mental disorders ($p=0.17$), self-perceptions and values ($p=0.063$), academic/occupational performance ($p=0.955$), emotions ($p=0.678$), behaviours ($p=0.832$), self-management strategies ($p=0.686$), social skills ($p=0.388$), family and significant relationships ($p=0.708$), physical health ($p=0.098$), sexual health ($p=0.148$) or quality of life ($p=0.889$) (eTable VI, Figure 4). For cognitive skills, interventions for children and adolescents fared better than interventions for adults ($p=0.007$). There were no differences between children and adolescents vs adults interventions for mental health literacy ($p=0.717$), attitude towards mental disorders ($p=0.127$), self-perceptions and values ($p=0.298$), academic/occupational performance ($p=0.143$), emotions ($p=0.440$), self-management strategies ($p=0.966$), social skills ($p=0.169$), family and significant relationships ($p=0.144$), physical health ($p=0.528$), sexual health ($p=0.908$) or quality of life ($p=0.578$) (eTable VII).

Heterogeneity and publication bias

Heterogeneity across the included studies was statistically significant for all outcomes ($p < 0.001$), ranging from 64.377 (behaviours) to 98.670 (quality of life). Publication bias was detected in two outcome categories and corrected with the trim and fill method: behaviours (ES 0.065 corrected to ES=0.099) and self-management strategies, which became non-significant after the correction (ES=0.517 corrected to ES=0.107) (eTable V).

Quality assessment

The quality of the included studies ranged from 0;0 (20.6% of the studies: no randomization, no blinding) to 2;2 (0.3% of the studies: clear method of randomization, double blinding) (eTable VIII). None of the studies employed triple blinding. The meta-regressions revealed no significant association between efficacy of the interventions to promote mental health and the quality of the studies (all $P > 0.05$) (eTable IX).

DISCUSSION

To our knowledge, this is the first systematic review and meta-analysis to comprehensively explore and summarize the impact of universal and selective interventions on the promotion of good mental health. The study demonstrated that universal and selective promotion of good mental health outcomes (mental health literacy, emotions, self-perceptions and values, quality of life, cognitive skills, social skills, physical health, sexual health, academic/occupational performance and attitude towards mental disorders) in young people is feasible and may be effective, although further research is needed.

Our review identified 276 individual intervention studies with 159,508 participants. We did not limit our search to interventions conducted in particular groups of individuals (e.g., health profession students (Lo et al., 2018) or nursing students (Li et al., 2018)) as in previous reviews. Furthermore, our 14 good mental health outcome categories were defined a-priori. We found a significant improvement in 10 out of the 13 meta-analyzable outcomes. Overall, the magnitude of the effect sizes for improving good mental health outcomes ranged from small (ES=0.177 to 0.49) regarding attitude towards mental disorders, self-perceptions and values, cognitive skills, academic/occupational performance, social skills, physical health, sexual health and quality of life, to medium (ES=0.541 to 0.685) for mental health literacy and emotions. There were no large effect sizes for improving good mental health and no effective interventions to improve self-management strategies, family and significant relationships or behaviours. Overall, the

ES were small to moderate, but they are comparable to ES found for the selective and universal prevention of poor mental health outcomes. Because universal and selective interventions typically target asymptomatic populations, variance in mental health outcomes is expected to be reduced and, consequently, the magnitude of the effect sizes for these interventions can only be expected to be relatively small. At the same time, since these samples are young and not yet affected by psychiatric conditions, small effect sizes can translate into relevant benefits in the longer term, some of which may extend to the whole general population.

Our meta-analysis also provided some insight on the comparability of universal vs selective or specific type of intervention categories that may be more effective to promote good mental health. The highest improvement in ES (0.685) was for mental health literacy, with no difference between universal or selective approaches. A previous systematic review has drawn attention into the difficulties observed in young individuals to recognize mental problems (Singh et al., 2019). Another systematic review suggested the potential utility of psychoeducation to improve mental health literacy (Tay et al., 2018), but we did not replicate this finding. We found that psychoeducation was the most effective intervention for the promotion of mental health literacy. Our results also support similar benefits of selective and universal interventions to improve emotion awareness and knowledge (ES=0.541). This finding is clinically relevant because it has the potential to enhance and to improve resilience and personal resources (Kushnir et al., 2012). We found that the category self-perceptions and values has been the most studied in the literature (92 studies). Previous meta-analytical evidence suggested that psychotherapy may improve this domain acting on self-esteem (from Cohen's $d=0.34$ for one-day workshops to $d=1.12$ for weekly sessions (Kolubinski et al., 2018)), but we did not confirm superiority of psychotherapy over other types of interventions. Although the ES of our findings was modest (ES=0.541), we demonstrate that universal and selective interventions to promote self-perceptions are equally effective and should not be reserved only for individuals who suffer from a mental disorder or subthreshold condition (Linardon et al., 2019). Our finding that universal and selective interventions can comparably improve quality of life (ES=0.457) extend previous systematic reviews and meta-analyses (Calver et al., 2018; Kolubinski et al., 2018) suggesting that this domain can be improved by psychosocial interventions in specific populations with mental and physical health problems. The only good mental health domain on which universal fared better than selective interventions related to cognitive skills (ES=0.428). Interventions for this outcome were particularly effective in children and adolescents, probably due to the maturational brain changes that occur during the early development and the increased

neuroplasticity in younger individuals (Fusar-Poli, 2019; Salazar de Pablo et al., 2019). While most meta-analyses in psychiatry focusing on interventions to improve social skills have focused on individuals with established mental disorders (e.g., psychosis (Turner et al., 2018) or autism-spectrum disorders (Gates et al., 2017)), with promising results, our study demonstrates that the acquisition of social skills can be improved through universal interventions aimed at the general population. This is clinically important because cognitive skills have shown to reduce psychological distress (Medalia et al., 2017), as well as to allow individuals to obtain better productivity and occupational outcomes (Chan et al., 2015; Kumar et al., 2017). Social skills may also lead to greater ability to initiate and maintain meaningful relationships (Fusar-Poli et al., 2019b). By training social skills, the ability of young children to cope with potentially difficult social situations can be strengthened (Sawyer et al., 1997). Psychoeducation was the best intervention for improving cognitive skills, which was associated with a large effect size (but see eLimitations). We also showed that it may be possible to use universal and selective interventions similarly, to promote physical health ($ES=0.285$) and sexual health ($ES=0.257$) in young individuals. Young individuals with mental health problems have poorer physical and sexual health than their general population counterparts (Kaushik et al., 2016). According to our meta-analysis, the category “other interventions”, which encompass physical exercise programs (Bang et al., 2017; Sharp and Caperchione, 2016a) is more beneficial than psychoeducation or psychotherapy for improving physical health in young people. This finding is consistent with guidelines for mentally ill patients (Stubbs et al., 2018). Improving physical health is an important goal in young people with emerging and manifest mental disorders, whose physical health is poorer than in the general population and associated with greater physical morbidity and mortality (Correll et al., 2015; Correll et al., 2017; De Hert et al., 2011a; DE Hert et al., 2011b; Firth et al., 2019; Stubbs et al., 2016). Universal and selective interventions are similarly effective to improve academic/occupational performance ($ES=0.21$), to a magnitude that is similar to that reported in a previous meta-analysis evaluating the effectiveness of physical activity ($ES=0.26$) (de Greeff et al., 2018) or psychoeducation ($ES=0.18$) (Conley et al., 2015) to boost academic performance. Finally, we found that universal and selective interventions are equally effective to improve attitude towards mental disorders ($ES=0.177$). A recent meta-analysis found that reductions in stigmatizing attitudes to prevent poor mental health are feasible in a larger scale with similar magnitude of the ES (Cohen’s $d=0.24$ to 0.40) (Morgan et al., 2018).

Despite these positive results, we found no evidence that interventions could improve behaviours, family and significant relationships or self-management strategies.

Furthermore, we did not find any studies about promotion of meaning of life in young people to be able to meta-analyse its interventions. Meaning of life has gained interest in the last years, and several selective interventions have been conducted, but, more frequently, in patients affected with medical conditions, particularly cancer (Henry et al., 2010; Park et al., 2019; Pelekasis et al., 2016). Meaning of life may be a domain that deserves more attention for the promotion of good mental health in young people as well, especially given the high suicide rates in that age group and, especially, those with emerging and manifest mental disorders.

This study has several limitations. First, the amount of evidence regarding some mental health outcomes was limited, including for meaning of life, sexual health and behaviours outcomes. However, the database was sufficiently large to robustly test most of the other outcomes. Second, we were unable to quantify the differential efficacy of specific interventions that were pooled in the “other” and “combined” subgroups or the differential efficacy of distinct types of psychotherapies. Also, the evidence on pharmacological interventions to promote good mental health was very limited and its efficacy remains unknown. We also did not evaluate the putative efficacy of diet or food supplements. Third, due to the limited granularity of information, we were unable to quantify the comparative efficacy for some of the intervention subgroups, for example attitude towards mental disorders, in which only psychoeducation interventions were found. Fourth, the quality of some studies was low, missing proper blinding and randomization, or at least did not sufficiently report these features. However, our meta-regression analyses found no associations between efficacy of interventions and quality of the studies. Fifth, we were not able to disentangle the optimal age range to deliver each of the interventions to promote good mental health except for cognitive skills, which seem more effective in children and adolescents. Sixth, the long-term efficacy of the interventions tested beyond the experimental period was not investigated. Finally, the available evidence is not stratified to specific subgroups of young people (stratified medicine) or tailored to the individual subject level (precision medicine (Fusar-Poli et al., 2018)). New collaborative multicentre studies (Las Hayas et al., 2019) and randomized controlled trials are needed to overcome these limitations. Importantly, it will be essential to achieve the development of more standardised instruments to measure a core outcome set of good mental health dimensions. This study paves the way towards the accomplishment of these goals over the next generation of research.

CONCLUSIONS

Several interventions seem effective to promote good mental health in young people. Although we have attempted to define the outcomes a-priori, there are no harmonised operationalisations and assessment instruments to ascertain them. Future research should develop innovative outcome sets and measurement criteria to consolidate and extend these findings.

Conflict of Interest Disclosures: Dr. Correll has been a consultant and/or advisor to or has received honoraria from: Alkermes, Allergan, Angelini, Boehringer-Ingelheim, Gedeon Richter, Gerson Lehrman Group, Indivior, IntraCellular Therapies, Janssen/J&J, LB Pharma, Lundbeck, MedAvante-ProPhase, Medscape, Merck, Neurocrine, Noven, Otsuka, Pfizer, Recordati, Rovi, Servier, Sumitomo Dainippon, Sunovion, Supernus, Takeda, and Teva. He has provided expert testimony for Bristol-Myers Squibb, Janssen, and Otsuka. He served on a Data Safety Monitoring Board for Boehringer-Ingelheim, Lundbeck, Rovi, Supernus, and Teva. He received royalties from UpToDate and grant support from Janssen and Takeda. He is also a stock option holder of LB Pharma. Dr. Kessing has been a consultant to or has received honoraria or grants from Lundbeck. Dr. Arango has been a consultant to or has received honoraria or grants from Acadia, Angelini, Gedeon Richter, Janssen Cilag, Lundbeck, Otsuka, Roche, Sage, Servier, Shire, Schering Plough, Sumitomo Dainippon Pharma, Sunovion and Takeda. Dr. Bechdolf has been a consultant and/or advisor to or has received honoraria or grants from: Janssen/J&J, LB Pharma, Lundbeck, Otsuka, Pfizer and Recordati. Dr Catalan has received personal fees from Janssen. Dr. Vieta has received grants and served as consultant, advisor or CME speaker for the following entities: Abbott, Allergan, Angelini, Dainippon Sumitomo Pharma, Janssen, Lundbeck, Novartis, Otsuka, Richter, Sage, Sanofi-Aventis, and Takeda. Dr Fusar-Poli has received grants from Lundbeck and personal fees from Menarini, Lundbeck and Angelini. No other disclosures were reported.

Funding/ Support: This study was supported by the European College of Neuropsychopharmacology Network Board. Dr Salazar de Pablo is supported by the Alicia Koplowitz Foundation. Dr Fusar-Poli is supported by the PSYSCAN project through the European Commission

Table 1. Empirical domains of good mental health (adapted from (Fusar-Poli et al., 2019b))

Domain	Definition
Mental health literacy	The ability to recognize and possess knowledge of a variety of different profiles of emerging and established mental disorders, factors and warning signs contributing to poor mental health as well as about the different mental health resources that can be accessed in case of need.
Attitude towards mental disorder	The way we react in front of someone with a mental disorder; Positive attitudes include understanding, being compassionate and empathic while decreasing discrimination towards people with poor mental health or with mental disorders. Negative attitude towards a person include stigma and stigmatizing behaviours, which leads to negative action or discrimination.
Self-perceptions and values	A collection of subjective beliefs, values and emotions about one's own internal and external characteristics, shaping one's attributional style, self-compassion and self-esteem, impacting awareness and acceptance leading to living a valued life.
Cognitive skills	The ability to pay attention, remember and organize information, while having a degree of cognitive flexibility, attention to enable decision making and solve problems.
Academic/occupational performance	Objective learning and knowledge, study achievements, attendance and behaviour as well as school adjustment and academic adaptation.
Emotions	Affective states with arousing or motivational properties that lead individuals to a certain response or behaviour.
Behaviours	Behaviours are the conducts in which a person proceeds when a stimulus is presented to them.
Self-management strategies	Practical, everyday skills needed to effectively and independently take care of oneself and to function and meet the demands of the environment: coping skills to deal with stress, problem solving and decision making to face the adversities that may appear.
Social skills	Social skills are abilities that allow young people to interact and communicate with each other in order to foster positive relations.
Family and significant relationships	The ability to establish meaningful relationships with other family members; healthy and positive relationships and connectedness with family members within a secure environment, which facilitates positive communication and interaction.
Physical health	Physical variables, visible symptoms and measures related to a positive physical status and dimensions of health.
Sexual health	A state of physical, emotional, mental and social well-being in relation to sexuality.
Meaning of life	Feeling that life has a purpose and a significance.
Quality of life	The general well-being of a person, defined in terms of health, happiness and satisfaction life.

Figure 1. Prisma flowchart

Figure 2. Overall efficacy of universal and selective interventions on good mental health outcomes

Figure 3. Comparative efficacy of type of preventive interventions

Figure 4. Comparative efficacy of universal vs selective interventions*

* Only mental health outcomes with both universal and selective interventions are represented in the graph

REFERENCES

- Altman, D.G., Simera, I., Hoey, J., Moher, D., Schulz, K., 2008. EQUATOR: reporting guidelines for health research. *Lancet* 371, 1149-1150.
- Arango, C., Díaz-Caneja, C.M., McGorry, P.D., Rapoport, J., Sommer, I.E., Vorstman, J.A., McDaid, D., Marín, O., Serrano-Drozdzowskyj, E., Freedman, R., Carpenter, W., 2018. Preventive strategies for mental health. *Lancet Psychiatry* 5, 591-604.
- Attwood, M., Meadows, S., Stallard, P., Richardson, T., 2012. Universal and targeted computerised cognitive behavioural therapy (Think, Feel, Do) for emotional health in schools: Results from two exploratory studies. *Child and Adolescent Mental Health* 17, 173-178.
- Bang, K.-S., Lee, I., Kim, S., Lim, C.S., Joh, H.-K., Park, B.-J., Song, M.K., 2017. The Effects of a Campus Forest-Walking Program on Undergraduate and Graduate Students' Physical and Psychological Health. *International Journal of Environmental Research and Public Health* 14.
- Calver, L., Tickle, A., Moghaddam, N., Biswas, S., 2018. The effect of psychological interventions on quality of life in patients with head and neck cancer: A systematic review and meta-analysis. *Eur J Cancer Care (Engl)* 27.
- Chan, J.Y., Hirai, H.W., Tsoi, K.K., 2015. Can computer-assisted cognitive remediation improve employment and productivity outcomes of patients with severe mental illness? A meta-analysis of prospective controlled trials. *J Psychiatr Res* 68, 293-300.
- Collins, S., Woolfson, L., Durkin, K., 2014. Effects on coping skills and anxiety of a universal school-based mental health intervention delivered in Scottish primary schools. 35(1), 85-100.
- Conley, C.S., Durlak, J.A., Kirsch, A.C., 2015. A Meta-analysis of universal mental health prevention programs for higher education students. *Prev Sci* 16, 487-507.
- Correll, C.U., Detraux, J., De Lepeleire, J., De Hert, M., 2015. Effects of antipsychotics, antidepressants and mood stabilizers on risk for physical diseases in people with schizophrenia, depression and bipolar disorder. *World Psychiatry* 14, 119-136.
- Correll, C.U., Solmi, M., Veronese, N., Bortolato, B., Rosson, S., Santonastaso, P., Thapa-Chhetri, N., Fornaro, M., Gallicchio, D., Collantoni, E., Pigato, G., Favaro, A., Monaco, F., Kohler, C., Vancampfort, D., Ward, P.B., Gaughran, F., Carvalho, A.F., Stubbs, B., 2017. Prevalence, incidence and mortality from cardiovascular disease in patients with pooled and specific severe mental illness: a large-scale meta-analysis of 3,211,768 patients and 113,383,368 controls. *World Psychiatry* 16, 163-180.
- de Greeff, J.W., Bosker, R.J., Oosterlaan, J., Visscher, C., Hartman, E., 2018. Effects of physical activity on executive functions, attention and academic performance in preadolescent children: a meta-analysis. *J Sci Med Sport* 21, 501-507.
- De Hert, M., Cohen, D., Bobes, J., Cetkovich-Bakmas, M., Leucht, S., Ndeti, D.M., Newcomer, J.W., Uwakwe, R., Asai, I., Möller, H.J., Gautam, S., Detraux, J., Correll, C.U., 2011a. Physical illness in patients with severe mental disorders. II. Barriers to care, monitoring and treatment guidelines, plus recommendations at the system and individual level. *World Psychiatry* 10, 138-151.

DE Hert, M., Correll, C.U., Bobes, J., Cetkovich-Bakmas, M., Cohen, D., Asai, I., Detraux, J., Gautam, S., Möller, H.J., Ndeti, D.M., Newcomer, J.W., Uwakwe, R., Leucht, S., 2011b. Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care. *World Psychiatry* 10, 52-77.

DerSimonian, R., Laird, N., 1986. Meta-analysis in clinical trials. *Control Clin Trials* 7, 177-188.

Fatori, D., Salum, G., Itria, A., Pan, P., Alvarenga, P., Rohde, L.A., Bressan, R., Gadelha, A., de Jesus Mari, J., Conceição do Rosário, M., Manfro, G., Polanczyk, G., Miguel, E.C., Graeff-Martins, A.S., 2018. The economic impact of subthreshold and clinical childhood mental disorders. *J Ment Health* 27, 588-594.

Firth, J., Siddiqi, N., Koyanagi, A., Siskind, D., Rosenbaum, S., Galletly, C., Allan, S., Cane, C., Carney, R., Carvalho, A.F., Chatterton, M.L., Correll, C.U., Curtis, J., Gaughran, F., Heald, A., Hoare, E., Jackson, S.E., Kisely, S., Lovell, K., Maj, M., McGorry, P.D., Mihalopoulos, C., Myles, H., O'Donoghue, B., Pillinger, T., Sarris, J., Schuch, F.B., Shiers, D., Smith, L., Solmi, M., Suetani, S., Taylor, J., Teasdale, S.B., Thornicroft, G., Torous, J., Usherwood, T., Vancampfort, D., Veronese, N., Ward, P.B., Yung, A.R., Killackey, E., Stubbs, B., 2019. The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *Lancet Psychiatry* 6, 675-712.

Fusar-Poli, P., 2019. Integrated Mental Health Services for the Developmental Period (0 to 25 Years): A Critical Review of the Evidence. *Front Psychiatry* 10, 355.

Fusar-Poli, P., Bauer, M., Borgwardt, S., Bechdolf, A., Correll, C.U., Do, K.Q., Domschke, K., Galderisi, S., Kessing, L.V., Koutsouleris, N., Krebs, M.O., Lennox, B., McGuire, P., Meyer-Lindenberg, A., Millan, M.J., Nieman, D., Pfennig, A., Sand, M., Whenert, A., van Amelsvoort, T., Arango, C., 2019a. European college of neuropsychopharmacology network on the prevention of mental disorders and mental health promotion (ECNP PMD-MHP). *Eur Neuropsychopharmacol* 29, 1301-1311.

Fusar-Poli, P., Borgwardt, S., Bechdolf, A., Addington, J., Riecher-Rossler, A., Schultze-Lutter, F., Keshavan, M., Wood, S., Ruhrmann, S., Seidman, L.J., Valmaggia, L., Cannon, T., Velthorst, E., De Haan, L., Cornblatt, B., Bonoldi, I., Birchwood, M., McGlashan, T., Carpenter, W., McGorry, P., Klosterkotter, J., McGuire, P., Yung, A., 2013. The Psychosis High-Risk State A Comprehensive State-of-the-Art Review. *Jama Psychiatry* 70, 107-120.

Fusar-Poli, P., Cappucciati, M., Rutigliano, G., Lee, T.Y., Beverly, Q., Bonoldi, I., Lelli, J., Kaar, S.J., Gago, E., Rocchetti, M., Patel, R., Bhavsar, V., Tognin, S., Badger, S., Calem, M., Lim, K., Kwon, J.S., Perez, J., McGuire, P., 2016. Towards a Standard Psychometric Diagnostic Interview for Subjects at Ultra High Risk of Psychosis: CAARMS versus SIPS. *Psychiatry J* 2016, 7146341.

Fusar-Poli, P., Hijazi, Z., Stahl, D., Steyerberg, E.W., 2018. The Science of Prognosis in Psychiatry: A Review. *JAMA Psychiatry* 75, 1289-1297.

Fusar-Poli, P., Salazar de Pablo, G., De Micheli, A., Nieman, D., Correll, C., Kessing, L.V., Pfennig, A., Bechdolf, A., Borgwardt, S., Arango, C., van Amelsvoort, T., 2019b. What is good mental health? A scoping review. *European Neuropsychopharmacology*.

Gates, J.A., Kang, E., Lerner, M.D., 2017. Efficacy of group social skills interventions for youth with autism spectrum disorder: A systematic review and meta-analysis. *Clin Psychol Rev* 52, 164-181.

Gore, F.M., Bloem, P.J., Patton, G.C., Ferguson, J., Joseph, V., Coffey, C., Sawyer, S.M., Mathers, C.D., 2011. Global burden of disease in young people aged 10-24 years: a systematic analysis. *Lancet* 377, 2093-2102.

Henry, M., Cohen, S.R., Lee, V., Sauthier, P., Provencher, D., Drouin, P., Gauthier, P., Gotlieb, W., Lau, S., Drummond, N., Gilbert, L., Stanimir, G., Sturgeon, J., Chasen, M., Mitchell, J., Huang, L.N., Ferland, M.K., Mayo, N., 2010. The Meaning-Making intervention (MMi) appears to increase meaning in life in advanced ovarian cancer: a randomized controlled pilot study. *Psychooncology* 19, 1340-1347.

Hiscock, H., Gulenc, A., Ukoumunne, O.C., Gold, L., Bayer, J., Shaw, D., Le, H., Wake, M., 2018. Preventing Preschool Mental Health Problems: Population-Based Cluster Randomized Controlled Trial. *Journal of Developmental and Behavioral Pediatrics* 39, 55-65.

Kaushik, A., Kostaki, E., Kyriakopoulos, M., 2016. The stigma of mental illness in children and adolescents: A systematic review. *Psychiatry Res* 243, 469-494.

Kolubinski, D.C., Frings, D., Nikčević, A.V., Lawrence, J.A., Spada, M.M., 2018. A systematic review and meta-analysis of CBT interventions based on the Fennell model of low self-esteem. *Psychiatry Res* 267, 296-305.

Kumar, K.S., Samuelkamaleshkumar, S., Viswanathan, A., Macaden, A.S., 2017. Cognitive rehabilitation for adults with traumatic brain injury to improve occupational outcomes. *Cochrane Database Syst Rev* 6, CD007935.

Kushnir, J., Friedman, A., Ehrenfeld, M., Kushnir, T., 2012. Coping with preoperative anxiety in cesarean section: physiological, cognitive, and emotional effects of listening to favorite music. *Birth* 39, 121-127.

Las Hayas, C., Izco-Basurko, I., Fullaondo, A., Gabrielli, S., Zwiefka, A., Hjemdal, O., Gudmundsdottir, D.G., Knoop, H.H., Olafsdottir, A.S., Donisi, V., Carbone, S., Rizzi, S., Mazur, I., Krolicka-Deregowska, A., Morote, R., Anyan, F., Ledertoug, M.M., Tange, N., Kaldalons, I., Jonsdottir, B.J., González-Pinto, A., Vergara, I., González, N., Mar Medina, J., de Manuel Keenoy, E., consortium, U., 2019. UPRIGHT, a resilience-based intervention to promote mental well-being in schools: study rationale and methodology for a European randomized controlled trial. *BMC Public Health* 19, 1413.

Li, C., Yin, H., Zhao, J., Shang, B., Hu, M., Zhang, P., Chen, L., 2018. Interventions to promote mental health in nursing students: A systematic review and meta-analysis of randomized controlled trials. *J Adv Nurs* 74, 2727-2741.

Linardon, J., Kothe, E.J., Fuller-Tyszkiewicz, M., 2019. Efficacy of psychotherapy for bulimia nervosa and binge-eating disorder on self-esteem improvement: Meta-analysis. *Eur Eat Disord Rev* 27, 109-123.

Lipsey, M., Wilson, D., 2000. *Practical Meta-analysis*. Sage Publications, Thousand Oaks, CA.

Lo, K., Waterland, J., Todd, P., Gupta, T., Bearman, M., Hassed, C., Keating, J.L., 2018. Group interventions to promote mental health in health professional education: a

systematic review and meta-analysis of randomised controlled trials. *Adv Health Sci Educ Theory Pract* 23, 413-447.

Medalia, A., Saperstein, A.M., Huang, Y., Lee, S., Ronan, E.J., 2017. Cognitive Skills Training for Homeless Transition-Age Youth: Feasibility and Pilot Efficacy of a Community Based Randomized Controlled Trial. *J Nerv Ment Dis* 205, 859-866.

Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., Group, P., 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 339, b2535.

Morgan, A.J., Reavley, N.J., Ross, A., Too, L.S., Jorm, A.F., 2018. Interventions to reduce stigma towards people with severe mental illness: Systematic review and meta-analysis. *J Psychiatr Res* 103, 120-133.

Park, C.L., Pustejovsky, J.E., Trevino, K., Sherman, A.C., Esposito, C., Berendsen, M., Salsman, J.M., 2019. Effects of psychosocial interventions on meaning and purpose in adults with cancer: A systematic review and meta-analysis. *Cancer* 125, 2383-2393.

Pelekasis, P., Zisi, G., Koumarianou, A., Marioli, A., Chrousos, G., Syrigos, K., Darviri, C., 2016. Forming a Stress Management and Health Promotion Program for Women Undergoing Chemotherapy for Breast Cancer: A Pilot Randomized Controlled Trial. *Integr Cancer Ther* 15, 165-174.

Pendry, P., Carr, A.M., Smith, A.N., Roeter, S.M., 2014. Improving adolescent social competence and behavior: a randomized trial of an 11-week equine facilitated learning prevention program. *The journal of primary prevention* 35, 281-293.

Polanczyk, G.V., Salum, G.A., Sugaya, L.S., Caye, A., Rohde, L.A., 2015. Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *J Child Psychol Psychiatry* 56, 345-365.

Rousseau, C., Benoit, M., Lacroix, L., Gauthier, M.F., 2009. Evaluation of a sandplay program for preschoolers in a multiethnic neighborhood. *J Child Psychol Psychiatry* 50, 743-750.

Salazar de Pablo, G., Vaquerizo Serrano, J.D., Gómez Vallejo, S., Sánchez Cerezo, J., Moreno Ruiz, C., 2019. Applications of Developmental Psychopathology. *Adv Exp Med Biol* 1192, 429-451.

Sawyer, M.G., MacMullin, C., Graetz, B., Said, J., Clark, J., Baghurst, P., 1997. Social skills training for primary school children: A 1-year follow-up study. *Journal of paediatrics and child health* 33, 378-383.

Sharp, P., Caperchione, C., 2016a. The effects of a pedometer-based intervention on first-year university students: A randomized control trial. *Journal of American college health* 64, 630-638.

Sharp, P., Caperchione, C., 2016b. The effects of a pedometer-based intervention on first-year university students: A randomized control trial. *J Am Coll Health* 64, 630-638.

Shinde, S., Weiss, H.A., Varghese, B., Khandeparkar, P., Pereira, B., Sharma, A., Gupta, R., Ross, D.A., Patton, G., Patel, V., 2018. Promoting school climate and health outcomes with the SEHER multi-component secondary school intervention in Bihar, India: a cluster-randomised controlled trial. *The Lancet* 392, 2465-2477.

Singh, S., Zaki, R.A., Farid, N.D.N., 2019. A systematic review of depression literacy: Knowledge, help-seeking and stigmatising attitudes among adolescents. *J Adolesc* 74, 154-172.

Stroup, D.F., Berlin, J.A., Morton, S.C., Olkin, I., Williamson, G.D., Rennie, D., Moher, D., Becker, B.J., Sipe, T.A., Thacker, S.B., 2000. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 283, 2008-2012.

Strunk, C.M., King, K.A., Vidourek, R.A., Sorter, M.T., 2014. Effectiveness of the surviving the Teens® suicide prevention and depression awareness program: an impact evaluation utilizing a comparison group. *Health Educ Behav* 41, 605-613.

Stubbs, B., Koyanagi, A., Veronese, N., Vancampfort, D., Solmi, M., Gaughran, F., Carvalho, A.F., Lally, J., Mitchell, A.J., Mugisha, J., Correll, C.U., 2016. Physical multimorbidity and psychosis: comprehensive cross sectional analysis including 242,952 people across 48 low- and middle-income countries. *BMC Med* 14, 189.

Stubbs, B., Vancampfort, D., Hallgren, M., Firth, J., Veronese, N., Solmi, M., Brand, S., Cordes, J., Malchow, B., Gerber, M., Schmitt, A., Correll, C.U., De Hert, M., Gaughran, F., Schneider, F., Kinnafick, F., Falkai, P., Möller, H.J., Kahl, K.G., 2018. EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and Position Statement from the European Psychiatric Association (EPA), supported by the International Organization of Physical Therapists in Mental Health (IOPTMH). *Eur Psychiatry* 54, 124-144.

Tay, J.L., Tay, Y.F., Klainin-Yobas, P., 2018. Effectiveness of information and communication technologies interventions to increase mental health literacy: A systematic review. *Early Interv Psychiatry* 12, 1024-1037.

Turner, D.T., McGlanaghy, E., Cuijpers, P., van der Gaag, M., Karyotaki, E., MacBeth, A., 2018. A Meta-Analysis of Social Skills Training and Related Interventions for Psychosis. *Schizophrenia Bulletin* 44, 475-491.

Witvliet, M., van Lier, P.A., Cuijpers, P., Koot, H.M., 2009. Testing links between childhood positive peer relations and externalizing outcomes through a randomized controlled intervention study. *J Consult Clin Psychol* 77, 905-915.

World Health Organization, 2002. Prevention and promotion in mental health. Mental health: evidence and research., in: Dependence., D.o.M.H.a.S. (Ed.), Geneva.

World Health Organization, 2004. Prevention of mental disorders: Effective interventions and policy options: Summary report. World Health Organization, Geneva.