



King's Research Portal

DOI:

[10.1016/j.neuroimage.2016.03.075](https://doi.org/10.1016/j.neuroimage.2016.03.075)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gifford, G., Crossley, N., Fusar-Poli, P., Schnack, H. G., Kahn, R. S., Koutsouleris, N., Cannon, T. D., & McGuire, P. (2017). Using neuroimaging to help predict the onset of psychosis. *NeuroImage*, 145(Part B), 209-217. <https://doi.org/10.1016/j.neuroimage.2016.03.075>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript


Using neuroimaging to help predict the onset of psychosis

George Gifford, Nicolas Crossley, Paolo Fusar-Poli, Hugo G. Schnack, René S. Kahn, Nikolaos Koutsouleris, Tyrone D. Cannon, Philip McGuire

PII: S1053-8119(16)30019-2
DOI: doi: [10.1016/j.neuroimage.2016.03.075](https://doi.org/10.1016/j.neuroimage.2016.03.075)
Reference: YNIMG 13080

To appear in: *NeuroImage*

Received date: 12 October 2015
Revised date: 18 March 2016
Accepted date: 28 March 2016



Please cite this article as: Gifford, George, Crossley, Nicolas, Fusar-Poli, Paolo, Schnack, Hugo G., Kahn, René S., Koutsouleris, Nikolaos, Cannon, Tyrone D., McGuire, Philip, Using neuroimaging to help predict the onset of psychosis, *NeuroImage* (2016), doi: [10.1016/j.neuroimage.2016.03.075](https://doi.org/10.1016/j.neuroimage.2016.03.075)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Using neuroimaging to help predict the onset of psychosis

George Gifford ^a, Nicolas Crossley ^a, Paolo Fusar-Poli ^a, Hugo G. Schnack ^b, René S. Kahn ^b, Nikolaos Koutsouleris ^c, Tyrone D. Cannon ^{d,e}, Philip McGuire ^a

^a Department of Psychosis Studies, The Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

^b Department of Psychiatry, Rudolf Magnus Institute of Neuroscience, University Medical Centre Utrecht, Utrecht, The Netherlands

^c Department of Psychiatry and Psychotherapy, Ludwig Maximilian University, Munich, Germany

^d Department of Psychology, Yale University, New Haven, Connecticut, United States of America

^e Department of Psychiatry, Yale University, New Haven, Connecticut, United States of America

Abstract

The aim of this review is to assess the potential for neuroimaging measures to facilitate prediction of the onset of psychosis. Research in this field has mainly involved people at 'ultra-high risk' (UHR) of psychosis, who have a very high risk of developing a psychotic disorder within a few years of presentation to mental health services.

The review details the key findings and developments in this area to date, and examines the methodological and logistical challenges associated with making

predictions in an individual subject in a clinical setting.

Key words

Psychosis prediction; Ultra High-Risk of psychosis; machine learning; Support Vector Machines; multimodal neuroimaging; multicentre neuroimaging studies; graph analysis

Psychosis prediction and the ultra high-risk state

Psychosis describes a syndrome that includes symptoms such as hallucinations, delusions, disorganised thought, and catatonia^{1 2}. Psychotic disorders are disorders that include symptoms of psychosis. The most common psychotic disorder is schizophrenia, however psychotic symptoms can occur in other disorders, for example in mood disorders such as bipolar and depression³.

Before the onset of symptoms that are severe enough to meet the clinical thresholds of a psychotic disorder, those affected experience subclinical psychotic symptoms and a decline in social / occupational functioning⁴. Recognising the value of detecting and treating psychosis early⁵, researchers and clinicians have developed the Ultra High-Risk (UHR) of psychosis criteria⁶. UHR criteria are the most commonly used criteria for indicating risk of developing a psychotic disorder⁷.

Criteria for UHR status include attenuated psychotic symptoms and / or a brief limited intermittent psychotic episode and / or a genetically determined vulnerability, alongside deterioration in social and occupational functioning^{7 8}. UHR status comes

with the caveat that those who meet UHR status are selectively those who have come into contact with clinical services. This review uses the term UHR throughout.

The UHR population is strikingly heterogeneous in terms of clinical outcomes. Follow-up studies^{9 10} suggest that 7 years after clinical presentation, approximately a third of UHR subjects will have developed a psychotic disorder, with most transitions occurring in the first 2 years¹¹. Most of those who do not develop a psychotic disorder will have persistent attenuated symptoms and / or have developed another mental health disorder, whilst 14% will have recovered (see figure 1).

Clinical intervention in the UHR group may reduce the likelihood of the onset of a psychotic disorder¹². However, as most UHR subjects do not develop a psychotic disorder, providing preventative treatment to all of those at risk is clinically inefficient. Identifying biomarkers that could be used to stratify the UHR group according to clinical outcome would enable the selective delivery of preventative interventions to the subgroup that would benefit the most.

It is difficult to predict clinical outcomes in an UHR subject on the basis of their clinical features at presentation. Although the clinical assessment at presentation has good diagnostic validity for ruling out a future psychotic disorder (meta analytical sensitivity of UHR assessment = 0.96)¹³, it has only a modest ability to rule in a future psychotic disorder (meta-analytical specificity of UHR assessment = 0.47)¹³. There is thus a need to find other forms of assessment that can improve the specificity of psychosis prediction.




Figure 1. Colour intensifies as those with underlying vulnerability move closer towards disorder manifestation. Taken with permission from Fusar-Poli et al. (2015)¹⁴ and based on data by Lin et al. (2011)¹⁰. [Single column fitting image]

Neuroimaging psychosis risk in the UHR population

Neuroimaging techniques have provided a wealth of new information about the

pathophysiology of psychiatric disorders¹⁰. Neuroimaging studies of the UHR population have implicated increased functional abnormalities in the prefrontal cortex (PFC), midbrain and medial temporal lobe (MTL)¹⁵, structural abnormalities involving the PFC^{16 17} and MTL¹⁸, elevated dopaminergic activity in the striatum¹⁹ and midbrain¹⁵, and abnormalities in levels of glutamate²⁰ and GABA²¹. In general, these abnormalities are qualitatively similar but less severe than those seen in patients with psychotic disorders⁵.

These findings represent the difference between the UHR population as a whole and healthy controls. However, the comparison that is relevant to the prediction of psychosis onset is between the minority of UHR subjects that subsequently develops a psychotic disorder and the majority that does not. Cross-sectional neuroimaging studies comparing these two subgroups at presentation have reported differences in activation in the PFC, MTL, midbrain and caudate^{10 22}, in the volume of the MTL, PFC, and cingulate cortex^{23 24}, in the integrity of white matter pathways²⁵, and in glutamate levels in the caudate nuclei²⁶. Longitudinal studies with repeated measurements before and after the onset of a psychotic disorder have described progressive changes in MTL and PFC volume^{14 27 28}, in white matter volume²⁹ and integrity³⁰, and in striatal dopamine function³¹. These changes were not evident in UHR subjects who did not develop a psychotic disorder.

Neuroimaging findings associated with the transition to psychosis have understandably been the focus of most studies to date, and most studies have subdivided UHR samples into groups who did or did not develop a psychotic

disorder. However, this subdivision is potentially misleading, as the subgroup that does not develop a psychotic disorder is not homogeneous: in some subjects, their presenting features resolve, such that they no longer meet the UHR criteria, whereas in others their symptoms persist. A further subgroup develop mental health disorders other than psychosis, particularly depression and anxiety (Figure 1)^{9 10}. Identifying neuroimaging measures that predict these other outcomes is also of interest. For example, determining the factors that predict recovery may improve our understanding of what determines resilience to mental illness, and could allow clinicians to avoid giving unnecessary treatments to people who would be very likely to recover without any intervention. Recent studies have begun to examine predictors of recovery in the UHR population, but at present this literature is still relatively small^{32 33}.

Outcome can also be defined in terms of level of functioning, as opposed to clinical status. Long term follow up studies suggest that an UHR individual's level of social and vocational functioning can sometimes be more clinically meaningful than their diagnostic category³⁴. Studies have also recently begun to examine the relationship between neuroimaging measures and functional outcomes^{32 35 36}.

Multi-centre studies

Although the results have been of great interest, to date most neuroimaging studies in UHR subjects have involved small sample sizes³⁷. Traditionally, mental health services have not engaged people with psychotic symptoms until after development of a first episode of a psychotic disorder. The concept of clinical provision for UHR

individuals is relatively new, and specialised services for this group are still limited to a minority of psychiatric centres. As UHR subjects are usually ascertained through specialised clinical services, this constrains recruitment to research studies. This can lead to studies being underpowered, particularly if the hypotheses relate to a specific clinical outcome (such as the onset of a psychotic disorder) that is only evident in a subset of the total UHR sample. This issue can be addressed through the recruitment of UHR subjects from multiple different sites: a number of such multi-centre studies involving neuroimaging have recently been completed^{15 19} and several others are on-going^{38 39 40}.

Although multi-centre studies can provide greater statistical power, there is a potential danger of introducing bias from between-centre differences in image acquisition and processing. For example, differences in scanning protocols^{41 42} and scanning hardware⁴³ can cause significant variability in results, as can upgrades of software and hardware within centres⁴⁴. These effects may be reduced through the harmonisation of scanning parameters and quality control across centres, and by the development of acquisition protocols that are specifically designed to minimise between centre variance, such as those used in the Alzheimer's Disease Neuroimaging Initiative (ADNI)^{45 44}.

Nevertheless, some between-centre variance is inevitable. Some studies quantify this by directly comparing neuroimaging data acquired from a group of volunteers⁴²^{46 47 48} or identical phantoms at different sites⁴⁹. Researchers have also used methods of post hoc statistical correction to adjust for between scanner variance⁵⁰.

However, if a predictive tool is to be employed at a variety of sites in a clinical setting, it may not be feasible to use phantoms or travelling subjects to control for between-site effects. In this case, the use of methods for post hoc correction may be more realistic.

Machine learning for individual prediction of psychosis transition


In clinical practice, predictions about the risk of psychosis need to be made using data from a single subject. However, most studies that have identified neuroimaging features that are associated with the onset of a psychotic disorder in UHR samples have described *average* differences between groups, which although statistically significant, describe two highly overlapping populations. In this context, individual inference has a limited utility. This has led to research on the application of statistical approaches that use multivariate patterns rather than group level differences, which are more suited to making individual level inferences. These approaches include machine-learning methods⁵¹, such as Random Forests⁵², Support Vector Machines (SVMs)⁵³, Linear Discriminant Analyses⁵⁴, and K-Nearest Neighbour algorithms⁵⁵. Machine learning methods allow for individual classification and are sensitive to subtle effects that would go unnoticed using traditional univariate analyses⁵⁶.

The method that has been most widely applied to neuroimaging data is the SVM⁵⁷, which may reflect its comparatively better performance with highly dimensional data⁵⁸. One reason for this is that SVMs base classification decisions on the most useful data points (see figure 2), which helps to overcome some of the difficulties

associated with modelling highly dimensional data ⁵³.

The support vector machine (SVM) is a supervised classification algorithm that learns from an initial 'training' dataset to classify new cases into two or more groups ⁵⁶. For example, a SVM can be trained on a dataset of baseline neuroimaging measurements from UHR subjects defined by whether they subsequently developed a clinically defined outcome or not, and can then be applied to a new sample of UHR individuals ⁵⁶ (see figure 2).

Figure 2. A. A classification boundary is created based on the maximum margin space between the two distributions of data points. Only data points near the margin (the support vectors) affect the classification boundary, facilitating a good generalization of the classification model. B. If it is not possible to create a linear classification boundary, a kernel function can be used to transform the data into higher dimensional space where classes become linearly separable. C. Schematic of SVM training and testing with neuroimaging data. In training, information from the two groups, or classes, is used to make a classification algorithm based on the predictive differences of the two groups. In testing, the algorithm is applied to data from a new subject to classify them as belonging to either group. [1.5 column fitting image]



A SVM is validated by demonstrating that it can classify individuals in a sample independent of the dataset it was originally trained on. Ideally this is done using a second dataset that has been acquired separately from the original⁵⁶. Validation

can also be attempted without a second dataset, for example by splitting the original sample (repeatedly), with one portion being used as the training dataset and other portions being used for cross-validation (e.g. k-fold cross-validation⁵⁹). However, any overlap between the training and test datasets will result in the model overfitting⁶⁰. This issue can be addressed by using methods that ensure separation between the training and test data, such as repeated double cross-validation⁶¹. Nevertheless, splitting a single sample for training and validation will include sampling error in the classification, and there is a risk that the original sample is atypically easy or difficult to classify, particularly when the sample is small.

Preliminary results from the application of SVMs in UHR samples have been promising. A SVM trained on volumetric MRI data was able to distinguish between UHR subjects who later transitioned and a combined group of non-transitioning UHR subjects and healthy controls with an accuracy of 88%, and between UHR who did not transition and a combined group of healthy controls and UHR who did transition with an accuracy of 86%⁶². A subsequent study used voxel-level MRI measures of grey matter volume to distinguish between UHR participants who did or did not go on to develop a psychotic disorder with an accuracy of 84.2%⁶³.

To date, applications of SVMs to UHR samples have used cross-validation methods within a single sample^{63 64} rather than the ideal validation method of using a second independent dataset. This partly reflects the fact that it is logistically difficult to ascertain large samples of UHR subjects with neuroimaging data and initial studies did not have access to another large dataset with comparable neuroimaging

measures. One study has used a SVM on two small datasets from different centres, although these employed different UHR criteria and MRI protocols. This reported an accuracy of classification of UHR transition outcomes of 80%⁶⁵, and also suggested that between centre effects could be minimised by using a sample pooled across both centres for training and testing⁶⁵.

The use of larger UHR samples would reduce (though not completely eliminate⁶⁶) the risk of models over-fitting the data⁶⁷. A model that over-fits the data is one that has been influenced by random error and noise in the training data to the extent that it does not accurately reflect the underlying phenomenon being studied⁶⁸. Over-fitting is a particular concern in models where the number of dimensions greatly outweighs the sample size and in highly complex models.

In addition to ensuring a sufficient sample size, over-fitting can be reduced by limiting the analysis to the most predictive features, using dimensionality reduction or subset selection techniques⁶⁹. It is also possible to use the results of previous research to limit predictive models to regions of interest⁷⁰, although this relies on the assumption that findings from mass univariate approaches are applicable to a multi-variate analyses.

The size of the sample studied also impacts on the generalizability of the findings. Small samples may result in higher fluctuations of accuracy estimates when using neuroimaging data in SVMs, suggesting poor model generalizability⁷¹. In addition, larger sample sizes are less affected by sampling effects and capture more of the population's heterogeneity, making them more likely to be representative of the

population being studied⁷². The numerous advantages of studying large samples has led to a wave of multicentre neuroimaging studies that are recruiting UHR subjects from several different centres, often using standardised image acquisition protocols^{38 39 40}. A further development has been the HARMONY initiative, which is designed to standardise neuroimaging, clinical, and electrophysiological assessments across these different multi-centre UHR consortia⁷³. As well as the opportunity to pool data to create even larger datasets, this will allow a SVM developed using one large sample to be validated on large independent datasets collected by other consortia.

There is also scope for the modification and optimisation of the machine learning methods that have been used in UHR subjects. For instance, SVMs can be modified to incorporate different statistical components. This includes the application of Bayesian probability and ‘fuzzy logic’ to SVMs, which have been used to create Relevance vector machines (RVM)⁷⁴ and Fuzzy SVMs⁷⁵ respectively. Both of these methods can be used to minimise the effect of outliers on the training phase of a machine^{74 75}.

By introducing a different loss function⁷⁶ support / relevance vector algorithms can also be used in regression (Support Vector Regression: SVR⁷⁷; Relevance Vector Regression: RVR⁷⁴), which allows for classification in terms of a continuous, as opposed to a categorical outcome. Tognin et al. (2014) used this approach to show that symptom progression in UHR subjects could be predicted using a RVR applied to cortical thickness at presentation⁷⁸. Regression-based machine learning has also

been applied to neuroimaging data to predict 'brain age', whereby individual brains are characterised according to indices of maturity^{79 80}. Such examples show high accuracy in predicting age based on MRI-derived brain descriptors in large-scale cohorts of healthy adolescents and adults⁸¹. In patient populations, Koutsouleris et al. (2013⁷⁹, 2015⁸²) demonstrated 'accelerated aging effects' using volumetric MRI data in different patients groups, including UHR participants. In a longitudinal study, Schnack et al. (2016) showed that aging of the brain continues to accelerate during the first five years after onset of a psychotic disorder⁸³. Brain age estimation has also been conducted with resting state fMRI data in a large sample of children, suggesting delayed maturation to predict the presence of mental health problems⁸⁴.

These findings suggest that machine learning-based biomarkers for psychosis prediction could be improved by taking account of the potential impact of neurodevelopmental changes on neuroimaging data. For example, predictors trained on UHR samples that are limited to subjects of a narrow age range may not be generalizable to samples that include subjects outside that range, who are at different stages of development. This issue may be particularly relevant to prediction in UHR samples, as these can include subjects within a wide age window (14-40 years), but there can be marked differences in the age of those enrolled at different centres, reflecting variations in the way that UHR subjects are recruited. For example, the mean age in UHR subjects recruited through adolescent or youth mental health services is significantly lower than in subjects enrolled through adult early detection services, and age therefore varies across studies⁸⁵. Large-scale collaborative efforts that include samples with different age ranges such as

HARMONY⁷³ may help to address the issue of ‘neurodevelopmental heterogeneity’ in the UHR state.

Multimodal prediction of psychosis onset

A feature of machine learning methods that has yet to be fully exploited in the UHR field is that their use is not restricted to neuroimaging data: they can be applied to multiple data modalities⁸⁶. The integration of data from different modalities may be particularly useful in predicting the onset of psychosis, as it is the result of interactions between a diversity of genetic, environmental and neurobiological factors⁸⁷. Moreover, multimodal neuroimaging studies in UHR subjects indicate that the normal relationships between glutamate and grey matter volume (MRS and sMRI)⁸⁸, glutamate and functional activity (MRS and fMRI)^{89 90 91}, glutamate and dopamine (MRS and PET)⁹² and dopamine and functional activity (PET and fMRI)^{15 93} are perturbed. These observations are consistent with animal models of psychosis, which also implicate interactions between the medial temporal cortex and the striatum and midbrain, and dopaminergic, glutaminergic and GABAergic dysfunction⁹⁴.

To date, most SVM studies in UHR subjects have involved data from a single neuroimaging modality (MRI), although the impact of introducing additional data modalities has recently been explored⁸⁶. This partly reflects the logistical demands associated with acquiring multi-modal neuroimaging data in samples that are large enough to permit comparison of subgroups with different clinical outcomes⁹⁵. However, ongoing multi-centre studies in this field involve a range of different

neuroimaging modalities, as well as genetic, clinical and cognitive data, and will be able to recruit samples large enough to assess predictors of outcomes.

There is some evidence that combining modalities may improve the prediction accuracy of machine learning classifiers. In predicting the onset of Alzheimer's disease from mild cognitive impairment, combining neuroimaging and neurocognitive measures increased the accuracy of SVMs beyond that of a single modality SVM^{96 97}. Conversely, a study of UHR, FEP and healthy control subjects found that prediction in some comparisons was more accurate when using a single neuroimaging modality compared to multiple modalities⁸⁶. The authors suggest that this may have been a result of an absence of complementary information in the additional modalities⁸⁶.

A further consideration in the use of multiple data modalities is practicality and cost. Data collection with some imaging modalities, such as PET, is expensive, can involve long acquisition times or may require specialist equipment and technical support⁴⁴. On the other hand, the collection of volumetric MRI, blood samples, and behavioural measures is feasible in many centres, making them the most viable options for wider clinical use. However, if predictive findings identified using relatively sophisticated neuroimaging techniques were closely correlated with measures in other domains, the latter could then serve as more accessible and less expensive proxy measures for a neuroimaging gold standard. For example, in the prediction of the onset of Alzheimer's disease, the level of past education and occupational functioning are putative proxy measures for PET data on brain glucose

metabolism⁹⁸.

Consortium recruiting large samples of UHR participants will collect a range of imaging, blood-based (bio-marker and genetic), and questionnaire based data^{38 39}⁴⁰. This is a benefit both due to the range of data-types available for predictive modelling and because it allows for an understanding of the interactions between different biological systems, which may be crucial to predicting the onset of psychosis. For example, it has been suggested that the hypothalamic-pituitary-adrenal axis acts as the mediating system for the influence of stress on psychotic symptoms, a system involving both hormones / inflammatory markers, brain areas, and psychological stress^{99 100 101 102}.

Another possibility is that less expensive measures may provide predictive models without the need for imaging. Promising preliminary results have been found using machine learning methods on blood-based biomarkers (mainly hormones and inflammatory markers) for the prediction of a future psychotic disorder^{103 104}. These early studies used around 15-30 blood-based markers^{103 104}, which gives an appropriate number of degrees of freedom for a predictive model, a benefit over models based on highly dimensional neuroimaging or genetic data. Given that blood-based biomarkers would likely reflect psychosis risk due to their interface with the brain however, predictive models using both imaging and blood-based biomarkers could perform better than each modality alone.

Graph analysis

Psychosis can be conceptualised as disorder of brain dysconnectivity^{105 106 107}, and neuroimaging studies suggest that both functional and structural connectivity are altered in people at UHR. Moreover, differences in connectivity have also been identified between UHR subjects who later develop a psychotic disorder and those who do not^{108 109}, suggesting that measures of connectivity may be useful in predicting the onset of psychosis¹¹⁰.

There are several methods that can be used to assess connectivity, but one of the most promising involves the application of network modelling approaches such as graph theory (GT), which forms graphs representing patterns of connectivity between brain areas¹¹¹. Brain graphs are conceptualised as comprising nodes (neuroanatomical regions) that are connected by edges (structural/functional connections)¹¹². This allows for connectivity to be characterised in terms of functional segregation (e.g. modularity), functional integration (e.g. shortest path length (see figure 3), global efficiency), small world, network motifs, centrality (e.g. degree), and network resilience (e.g. degree distribution)¹¹³. Graph analysis thus provides information on the integrative features of brain networks.

Brain graphs in healthy participants suggest that the brain is normally organised in a way that maximises the integration of the whole network, while allowing for the segregation of specific brain regions. Brain graphs typically show a modular organization, such that the network can be broken down into smaller sub networks¹¹¹, with modules specialized for specific computational processes¹¹⁴

Within brain networks, certain nodes, termed hubs, are much more connected than others ¹¹¹, and form ‘rich clubs’ ¹¹⁵ (see figure 3). Hubs have more long distance connections and higher metabolic demands than regular nodes ^{116 117}, and because of their strategic position in the network are thought to have an important role in brain functioning ^{118 119}.




Figure 3. A. Visualisation of a brain graph with 90 defined nodes, parcellated according to the Automatic Anatomical Labelling atlas ¹²⁰. B. Simple example of shortest path length. Shortest path between the two blue nodes highlighted by blue edges. C. Example of a network that shows community structure (sub-networks highlighted in grey) with hubs (purple nodes) connected in a rich club (purple edges). [Single column fitting image]

Graph analysis of neuroimaging data in patients with schizophrenia suggests that the disorder is associated with decreased efficiency¹²¹, clustering and small worldness¹²², and MRI abnormalities in psychotic disorders appear to be preferentially concentrated in hub regions^{123 124}. Abnormal GT measures have also been associated with elevated psychotic symptomatology in a sample of UHR participants¹²⁵.

The mathematical functions of machine learning cannot easily be applied to graphs however¹²⁶. To date, studies that have used graph analysis metrics for machine learning have used whole brain graph theory metrics rather than representing the characteristics of individual nodes and / or edges^{127 128}. In answer to this, methods for transforming graph data in to real vector space have been proposed, allowing for the application of machine learning classification and pattern recognition methods to graph data^{126 129}. For instance, techniques such as graph embedding allow for vector representation of nodes or edges, which can additionally incorporate graph topological features¹³⁰. Additionally, graph kernels carry the possibility of applying graphs to kernel based machines such as SVMs¹²⁶. Developments in this area are on-going¹³¹ but these methods have not yet been validated in brain graphs, and it remains to be seen whether the complex and noisy graphs that are derived from neuroimaging data can be adequately represented¹³⁰.

Conclusions

Because it is difficult to predict whether a UHR subject will develop a psychotic disorder on the basis of a clinical assessment, there is a need for biomarkers that

can help to predict outcomes in this group. Neuroimaging measures have the potential to facilitate the prediction of outcomes in UHR subjects. Further progress in their application requires the development of methods that permit prediction using data from an individual subject, and their validation in large independent samples. The extent to which the accuracy of prediction is enhanced by using more than one type of imaging data, or incorporating non-imaging data remains to be determined. Many of these issues are currently being addressed in multi-centre neuroimaging studies in UHR subjects.

The on-going effort to use neuroimaging to facilitate prediction of the onset of psychosis provides a good example of a recent shift in the focus of psychiatric neuroimaging research, from descriptive studies to investigations with an explicitly translational objective.

Acknowledgements

George Gifford is supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care South London at King's College Hospital NHS Foundation Trust. The views expressed are those of the author[s] and not necessarily those of the NHS, the NIHR or the Department of Health.

Highlights

- Machine learning methods allow for prediction of psychosis outcomes in individuals.
- Current multicentre studies seek to recruit sufficient samples for ML prediction.
- Multiple modalities can be incorporated into ML predictive models.
- ML methods can be used to predict continuous outcomes.
- ML can incorporate graph analysis if data is transformed into vector space.

ACCEPTED MANUSCRIPT

References

-
- ¹ American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders: DSM-5. Washington, D.C: American Psychiatric Association.
- ² World Health Organization. (1992). *The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and diagnostic guidelines*. Geneva: World Health Organization.
- ³ Howes, O. D., & Murray, R. M. (2014). Schizophrenia: an integrated sociodevelopmental-cognitive model. *The Lancet*, 383(9929), 1677-1687.
- ⁴ Yung, A. R., & McGorry, P. D. (1996). The prodromal phase of first-episode psychosis: past and current conceptualizations. *Schizophrenia bulletin*, 22(2), 353-370.
- ⁵ McGorry, P. D., Killackey, E., & Yung, A. (2008). Early intervention in psychosis: concepts, evidence and future directions. *World Psychiatry*, 7(3), 148-156.
- ⁶ Yung, A. R., Stanford, C., Cosgrave, E., Killackey, E., Phillips, L., Nelson, B., & McGorry, P. D. (2006). Testing the Ultra High Risk (prodromal) criteria for the prediction of psychosis in a clinical sample of young people. *Schizophrenia research*, 84(1), 57-66. Doi: 10.1016/j.schres.2006.03.014
- ⁷ Fusar-Poli, P., Borgwardt, S., Bechdolf, A., Addington, J., Riecher-Rössler, A., Schultze-Lutter, F., ... & Valmaggia, L. (2013). The psychosis high-risk state: a comprehensive state-of-the-art review. *JAMA psychiatry*, 70(1), 107-120.

-
- ⁸ Fusar-Poli, P., McGuire, P., & Borgwardt, S. (2012). Mapping prodromal psychosis: a critical review of neuroimaging studies. *European Psychiatry*, 27(3), 181-191. Doi: 10.1016/j.eurpsy.2011.06.006
- ⁹ Nelson, B., Yuen, H. P., Wood, S. J., Lin, A., Spiliotacopoulos, D., Bruxner, A., ... & Yung, A. R. (2013). Long-term follow-up of a group at ultra high risk ("prodromal") for psychosis: the PACE 400 study. *JAMA psychiatry*, 70(8), 793-802. Doi: 10.1001/jamapsychiatry.2013.1270
- ¹⁰ Lin, A., Wood, S. J., Nelson, B., Brewer, W. J., Spiliotacopoulos, D., Bruxner, A., ... & Yung, A. R. (2011). Neurocognitive predictors of functional outcome two to 13 years after identification as ultra-high risk for psychosis. *Schizophrenia research*, 132(1), 1-7. Doi: 10.1016/j.schres.2011.06.014
- ¹¹ Kempton, M. J., Bonoldi, I., Valmaggia, L., McGuire, P., & Fusar-Poli, P. (2015). Speed of Psychosis Progression in People at Ultra-High Clinical Risk: A Complementary Meta-analysis. *JAMA psychiatry*, 72(6), 622-623. Doi: 10.1001/jamapsychiatry.2015.0094
- ¹² Stafford, M. R., Jackson, H., Mayo-Wilson, E., Morrison, A. P., & Kendall, T. (2013). Early interventions to prevent psychosis: systematic review and meta-analysis. *BMJ*, 346, f185. Doi: 10.1136/bmj.f185
- ¹³ Fusar-Poli, P., Cappucciati, M., Rutigliano, G., Schultze-Lutter, F., Bonoldi, I., Borgwardt, S., Riecher-Rossler, A., Addington, J., Perkins, D., Woods, S. W., McGlashan, T. H., Lee, J., Klosterkötter, J., Yung, A. R., & McGuire, P. (2015). At risk or not at risk? A meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. *World Psychiatry*, 14(3), 322-332.

-
- ¹⁴ Fusar-Poli, P. (2015). The Enduring Search for the Koplik Spots of Psychosis. *JAMA psychiatry*, 72(9), 863-864. Doi:10.1001/jamapsychiatry.2015.0611
- ¹⁵ Allen, P., Luigjes, J., Howes, O. D., Egerton, A., Hirao, K., Valli, I., ... & McGuire, P. (2012). Transition to psychosis associated with prefrontal and subcortical dysfunction in ultra high-risk individuals. *Schizophrenia bulletin*, 38(6), 1268-1276. Doi: 10.1093/schbul/sbr194
- ¹⁶ Jung, W. H., Kim, J. S., Jang, J. H., Choi, J. S., Jung, M. H., Park, J. Y., ... & Kwon, J. S. (2011). Cortical thickness reduction in individuals at ultra-high-risk for psychosis. *Schizophrenia bulletin*, 37(4), 839-849. Doi: 10.1093/schbul/sbp151
- ¹⁷ Mechelli, A., Riecher-Rössler, A., Meisenzahl, E. M., Tognin, S., Wood, S. J., Borgwardt, S. J., ... & McGuire, P. (2011). Neuroanatomical abnormalities that predate the onset of psychosis: a multicenter study. *Archives of general psychiatry*, 68(5), 489-495.
- ¹⁸ Buehlmann, E., Berger, G. E., Aston, J., Gschwandtner, U., Pflueger, M. O., Borgwardt, S. J., ... & Riecher-Rössler, A. (2010). Hippocampus abnormalities in at risk mental states for psychosis? A cross-sectional high resolution region of interest magnetic resonance imaging study. *Journal of psychiatric research*, 44(7), 447-453.
- ¹⁹ Egerton, A., Chaddock, C. A., Winton-Brown, T. T., Bloomfield, M. A., Bhattacharyya, S., Allen, P., ... & Howes, O. D. (2013). Presynaptic striatal dopamine dysfunction in people at ultra-high risk for psychosis: findings in a second cohort. *Biological psychiatry*, 74(2), 106-112. Doi: 10.1016/j.biopsych.2012.11.017

- ²⁰ Stone, J. M., Howes, O. D., Egerton, A., Kambeitz, J., Allen, P., Lythgoe, D. J., ... & McGuire, P. (2010). Altered relationship between hippocampal glutamate levels and striatal dopamine function in subjects at ultra high risk of psychosis. *Biological psychiatry*, *68*(7), 599-602. Doi: 10.1016/j.biopsych.2010.05.034
- ²¹ Kang, J. I., Park, H. J., Kim, S. J., Kim, K. R., Lee, S. Y., Lee, E., ... & Lee, J. D. (2013). Reduced binding potential of GABA-A/benzodiazepine receptors in individuals at ultra-high risk for psychosis: an [18F]-fluoroflumazenil positron emission tomography study. *Schizophrenia bulletin*, *30*(3), 548-557. Doi: 10.1093/schbul/sbt052
- ²² Sabb, F. W., van Erp, T. G., Hardt, M. E., Dapretto, M., Caplan, R., Cannon, T. D., & Bearden, C. E. (2010). Language network dysfunction as a predictor of outcome in youth at clinical high risk for psychosis. *Schizophrenia research*, *116*(2), 173-183. Doi: 10.1016/j.schres.2009.09.042
- ²³ Pantelis, C., Velakoulis, D., McGorry, P. D., Wood, S. J., Suckling, J., Phillips, L. J., ... & McGuire, P. K. (2003). Neuroanatomical abnormalities before and after onset of psychosis: a cross-sectional and longitudinal MRI comparison. *The Lancet*, *361*(9354), 281-288. Doi: 10.1016/S0140-6736(03)12323-9
- ²⁴ Mechelli, A., Riecher-Rössler, A., Meisenzahl, E. M., Tognin, S., Wood, S. J., Borgwardt, S. J., ... & McGuire, P. (2011). Neuroanatomical abnormalities that predate the onset of psychosis: a multicenter study. *Archives of general psychiatry*, *68*(5), 489-495. Doi: 10.1001/archgenpsychiatry.2011.42
- ²⁵ Bloemen, O. J. N., De Koning, M. B., Schmitz, N., Nieman, D. H., Becker, H. E., De Haan, L., ... & van Amelsvoort, T. A. M. J. (2010). White-matter markers

for psychosis in a prospective ultra-high-risk cohort. *Psychological medicine*, 40(08), 1297-1304. Doi: 10.1017/S0033291709991711

²⁶ de la Fuente-Sandoval, C., León-Ortiz, P., Azcárraga, M., Favila, R., Stephano, S., & Graff-Guerrero, A. (2013). Striatal glutamate and the conversion to psychosis: a prospective 1H-MRS imaging study. *International Journal of Neuropsychopharmacology*, 16(2), 471-475. Doi: 10.1017/S1461145712000314

²⁷ Borgwardt, S. J., McGuire, P. K., Aston, J., Gschwandtner, U., Pflüger, M. O., Stieglitz, R. D., ... & Riecher-Rössler, A. (2008). Reductions in frontal, temporal and parietal volume associated with the onset of psychosis. *Schizophrenia research*, 106(2), 108-114. Doi: 10.1016/j.schres.2008.08.007

²⁸ Cannon, T. D., Chung, Y., He, G., Sun, D., Jacobson, A., van Erp, T. G., ... & North American Prodrome Longitudinal Study Consortium. (2015). Progressive reduction in cortical thickness as psychosis develops: a multisite longitudinal neuroimaging study of youth at elevated clinical risk. *Biological psychiatry*, 77(2), 147-157. Doi: 10.1016/j.biopsych.2014.05.023

²⁹ Ziermans, T. B., Schothorst, P. F., Schnack, H. G., Koolschijn, P. C. M., Kahn, R. S., van Engeland, H., & Durston, S. (2012). Progressive structural brain changes during development of psychosis. *Schizophrenia bulletin*, 38(3), 519-530. doi: 10.1093/schbul/sbq113

³⁰ Peters, B. D., Dingemans, P. M., Dekker, N., Blaas, J., Akkerman, E., van Amelsvoort, T. A., ... & de Haan, L. (2010). White matter connectivity and psychosis in ultra-high-risk subjects: a diffusion tensor fiber tracking study. *Psychiatry Research: Neuroimaging*, 181(1), 44-50. Doi: 10.1016/j.psychres.2009.10.008

- ³¹ Howes, O. D., Bose, S. K., Turkheimer, F., Valli, I., Egerton, A., Valmaggia, L. R., ... & McGuire, P. (2011). Dopamine synthesis capacity before onset of psychosis: a prospective [18F]-DOPA PET imaging study. *American Journal of Psychiatry*, *168*(12), 1311-1317. Doi: 10.1176/appi.ajp.2011.11010160
- ³² Allen, P., Chaddock, C. A., Egerton, A., Howes, O. D., Barker, G., Bonoldi, I., ... & McGuire, P. (2015). Functional outcome in people at high risk for psychosis predicted by thalamic glutamate levels and prefronto-striatal activation. *Schizophrenia bulletin*, *41*(2), 429-439. Doi: 10.1093/schbul/sbu115
- ³³ Egerton, A., Stone, J. M., Chaddock, C. A., Barker, G. J., Bonoldi, I., Howard, R. M., ... & McGuire, P. K. (2014). Relationship between brain glutamate levels and clinical outcome in individuals at ultra high risk of psychosis. *Neuropsychopharmacology*, *39*(12), 2891-2899. Doi: 10.1038/npp.2014.143
- ³⁴ Lin, A., Wood, S. J., Nelson, B., Brewer, W. J., Spiliotacopoulos, D., Bruxner, A., ... & Yung, A. R. (2011). Neurocognitive predictors of functional outcome two to 13years after identification as ultra-high risk for psychosis. *Schizophrenia research*, *132*(1), 1-7.
- ³⁵ Karlsgodt, K. H., Niendam, T. A., Bearden, C. E., & Cannon, T. D. (2009). White matter integrity and prediction of social and role functioning in subjects at ultra-high risk for psychosis. *Biological psychiatry*, *66*(6), 562-569. Doi: 10.1016/j.biopsych.2009.03.013
- ³⁶ Kambeitz-Illankovic, L., Meisenzahl, E. M., Cabral, C., von Saldern, S., Kambeitz, J., Falkai, P., ... & Koutsouleris, N. (2015). Prediction of outcome in the psychosis prodrome using neuroanatomical pattern classification *Schizophrenia research*. Doi: 10.1016/j.schres.2015.03.005

- ³⁷ Fusar-Poli, P., Rocchetti, M., Sardella, A., Avila, A., Brandizzi, M., Caverzasi, E., Politi, P., Ruhrmann, S. & McGuire, P. Disorder, not just state of risk: meta-analysis of functioning and quality of life in people at high risk of psychosis. *The British Journal of Psychiatry*, 207(3), 198-206. Doi: 10.1192/bjp.bp.114.157115
- ³⁸ European Network of National Schizophrenia Networks Studying Gene-Environment Interactions. <http://www.eu-gei.eu> (accessed June 5th 2015)
- ³⁹ PSYSCAN - Translating neuroimaging findings from research into clinical practice. <http://www.psyscan.eu> (accessed June 5th 2015)
- ⁴⁰ Personalised Prognostic Tools for Early Psychosis Management. <http://www.pronia.eu> (accessed June 5th 2015)
- ⁴¹ Kempton, M. J., Underwood, T. S., Brunton, S., Stylios, F., Schmechtig, A., Ettinger, U., ... & Simmons, A. (2011). A comprehensive testing protocol for MRI neuroanatomical segmentation techniques: evaluation of a novel lateral ventricle segmentation method. *Neuroimage*, 58(4), 1051-1059. Doi: 0.1016/j.neuroimage.2011.06.080
- ⁴² Schnack, H. G., van Haren, N. E., Brouwer, R. M., van Baal, G. C. M., Picchioni, M., Weisbrod, M., ... & Hilleke, E. (2010). Mapping reliability in multicenter MRI: Voxel-based morphometry and cortical thickness. *Human brain mapping*, 31(12), 1967-1982. Doi: 10.1002/hbm.20991
- ⁴³ Pfefferbaum, A., Adalsteinsson, E., & Sullivan, E. V. (2003). Replicability of diffusion tensor imaging measurements of fractional anisotropy and trace in brain. *Journal of Magnetic Resonance Imaging*, 18(4), 427-433. Doi: 10.1002/jmri.10377

-
- ⁴⁴ Kempton, M. J., & McGuire, P. (2015). How can neuroimaging facilitate the diagnosis and stratification of patients with psychosis? *European Neuropsychopharmacology*, *25*(5), 725-732. Doi: 10.1016/j.euroneuro.2014.07.006
- ⁴⁵ Jack, C. R., Bernstein, M. A., Fox, N. C., Thompson, P., Alexander, G., Harvey, D., ... & Weiner, M. W. (2008). The Alzheimer's disease neuroimaging initiative (ADNI): MRI methods. *Journal of Magnetic Resonance Imaging*, *27*(4), 685-691. Doi: 10.1002/jmri.21049
- ⁴⁶ Suckling, J., Barnes, A., Job, D., Brennan, D., Lymer, K., Dazzan, P., ... & Lawrie, S. (2012). The neuro/PsyGRID calibration experiment. *Human brain mapping*, *33*(2), 373-386. Doi: 10.1002/hbm.21210
- ⁴⁷ Gradin, V., Gountouna, V. E., Waiter, G., Ahearn, T. S., Brennan, D., Condon, B., ... & Steele, J. D. (2010). Between-and within-scanner variability in the CaliBrain study n-back cognitive task. *Psychiatry Research: Neuroimaging*, *184*(2), 86-95. Doi: 10.1016/j.psychresns.2010.08.010
- ⁴⁸ Moorhead, T. W. J., Gountouna, V. E., Job, D. E., McIntosh, A. M., Romaniuk, L., Lymer, G. K. S., ... & Lawrie, S. M. (2009). Prospective multi-centre Voxel Based Morphometry study employing scanner specific segmentations: procedure development using CaliBrain structural MRI data. *BMC medical imaging*, *9*(1), 8. Doi: 10.1186/1471-2342-9-8
- ⁴⁹ Gunter, J. L., Bernstein, M. A., Borowski, B. J., Ward, C. P., Britson, P. J., Felmlee, J. P., ... & Jack, C. R. (2009). Measurement of MRI scanner performance with the ADNI phantom. *Medical physics*, *36*(6), 2193-2205.

- ⁵⁰ Koutsouleris, N., Riecher-Rössler, A., Meisenzahl, E. M., Smieskova, R., Studerus, E., Kambaitz-Ilankovic, L., ... & Borgwardt, S. (2014). Detecting the psychosis prodrome across high-risk populations using neuroanatomical biomarkers. *Schizophrenia bulletin*, sbu078. Doi: 10.1093/schbul/sbu078
- ⁵¹ Orrù, G., Pettersson-Yeo, W., Marquand, A. F., Sartori, G., & Mechelli, A. (2012). Using support vector machine to identify imaging biomarkers of neurological and psychiatric disease: a critical review. *Neuroscience & Biobehavioral Reviews*, 36(4), 1140-1152. Doi: 10.1016/j.neubiorev.2012.01.004
- ⁵² Breiman, L. (2001). Random forests. *Machine learning*, 45(1), 5-32.
- ⁵³ Cortes, C., & Vapnik, V. (1995). Support-vector networks. *Machine learning*, 20(3), 273-297.
- ⁵⁴ McLachlan, G. (2004). *Discriminant analysis and statistical pattern recognition* (Vol. 544). John Wiley & Sons.
- ⁵⁵ Dasarathy, B. V. (1991). Nearest neighbor (NN) norms: NN pattern classification techniques.
- ⁵⁶ Orrù, G., Pettersson-Yeo, W., Marquand, A. F., Sartori, G., & Mechelli, A. (2012). Using support vector machine to identify imaging biomarkers of neurological and psychiatric disease: a critical review. *Neuroscience & Biobehavioral Reviews*, 36(4), 1140-1152. Doi: 10.1016/j.neubiorev.2012.01.004
- ⁵⁷ Pettersson-Yeo, W., Benetti, S., Marquand, A. F., Dell'Acqua, F., Williams, S. C. R., Allen, P., ... & Mechelli, A. (2013). Using genetic, cognitive and multi-modal neuroimaging data to identify ultra-high-risk and first-episode psychosis at the individual level. *Psychological medicine*, 43(12), 2547-2562. Doi:

10.1017/S003329171300024X

⁵⁸ Khondoker, M., Dobson, R., Skirrow, C., Simmons, A., & Stahl, D. (2013). A comparison of machine learning methods for classification using simulation with multiple real data examples from mental health studies. *Statistical methods in medical research*. Doi: 10.1177/0962280213502437

⁵⁹ Anguita, D., Ghio, A., Ridella, S., & Sterpi, D. (2009). K-Fold Cross Validation for Error Rate Estimate in Support Vector Machines. In DMIN (pp. 291-297).

⁶⁰ Varma S, Simon R. (2006). Bias in error estimation when using cross-validation for model selection. *BMC Bioinformatics*, 7(1), 91.

⁶¹ Filzmoser, P., Liebmann, B., & Varmuza, K. (2009). *Repeated double cross validation*. *Journal of Chemometrics*, 23(4), 160-171. Doi: 10.1002/cem.1225

⁶² Koutsouleris, N., Meisenzahl, E. M., Davatzikos, C., Bottlender, R., Frodl, T., Scheuerecker, J., ... & Gaser, C. (2009). Use of neuroanatomical pattern classification to identify subjects in at-risk mental states of psychosis and predict disease transition. *Archives of general psychiatry*, 66(7), 700-712. Doi:10.1001/archgenpsychiatry.2009.62

⁶³ Koutsouleris, N., Borgwardt, S., Meisenzahl, E. M., Bottlender, R., Möller, H. J., & Riecher-Rössler, A. (2012). Disease prediction in the at-risk mental state for psychosis using neuroanatomical biomarkers: results from the FePsy study. *Schizophrenia bulletin*, 38(60), 1234-1246. Doi: 10.1093/schbul/sbr145

⁶⁴ Koutsouleris, N., Meisenzahl, E. M., Davatzikos, C., Bottlender, R., Frodl, T., Scheuerecker, J., ... & Gaser, C. (2009). Use of neuroanatomical pattern classification to identify subjects in at-risk mental states of psychosis and predict

disease transition. *Archives of general psychiatry*, 66(7), 700-712.

Doi:10.1001/archgenpsychiatry.2009.62

⁶⁵ Koutsouleris, N., Riecher-Rössler, A., Meisenzahl, E. M., Smieskova, R., Studerus, E., Kambitz-Illankovic, L., ... & Borgwardt, S. (2014). Detecting the psychosis prodrome across high-risk populations using neuroanatomical biomarkers. *Schizophrenia bulletin*, 41(2), 471-482.. Doi: 10.1093/schbul/sbu078

⁶⁶ Joachims, T. (1998). Text categorization with support vector machines: Learning with many relevant features (pp. 137-142). Springer Berlin Heidelberg.

⁶⁷ Guyon, I., Weston, J., Barnhill, S., & Vapnik, V. (2002). Gene selection for cancer classification using support vector machines. *Machine learning*, 46(1-3), 389-422.

⁶⁸ Dietterich, T. (1995). Overfitting and under computing in machine learning. *ACM computing surveys (CSUR)*, 27(3), 326-327.

⁶⁹ Mwangi, B., Tian, T. S., & Soares, J. C. (2014). A review of feature reduction techniques in neuroimaging. *Neuroinformatics*, 12(2), 229-244. Doi: 10.1007/s12021-013-9204-3

⁷⁰ Costafreda, S. G., Dinov, I. D., Tu, Z., Shi, Y., Liu, C. Y., Kloszewska, I., ... & Simmons, A. (2011). Automated hippocampal shape analysis predicts the onset of dementia in mild cognitive impairment. *Neuroimage*, 56(1), 212-219. Doi: 10.1016/j.neuroimage.2011.01.050

⁷¹ Nieuwenhuis, M., van Haren, N. E., Pol, H. E. H., Cahn, W., Kahn, R. S., & Schnack, H. G. (2012). Classification of schizophrenia patients and healthy controls from structural MRI scans in two large independent samples.

Neuroimage, 61(3), 606-612. Doi: 10.1016/j.neuroimage.2012.03.079

⁷² Schnack, H. G., & Kahn, R. (2016). Detecting Neuroimaging Biomarkers for Psychiatric Disorders: Sample Size Matters. *Frontiers in Psychiatry*, 7:50. Doi: 10.3389/fpsy.2016.00050

⁷³ Cannon, T.D. (March, 2015). *Biomarkers of Vulnerability and Progression in the Psychosis Prodrome*. Presented at the International Congress on Schizophrenia Research, Colorado Springs, Colorado.

⁷⁴ Tipping, M. E. (2001). Sparse Bayesian learning and the relevance vector machine. *The journal of machine learning research*, 1, 211-244.

⁷⁵ Lin, C. F., & Wang, S. D. (2002). Fuzzy support vector machines, *IEEE Transactions on Neural Networks*, 13(2), 464-471. Doi: 10.1109/72.991432

⁷⁶ Smola, A. J. (1996). Regression estimation with support vector learning machines. *Master's thesis, Technische Universit at Munchen*.

⁷⁷ Smola, A., & Vapnik, V. (1997). Support vector regression machines. *Advances in neural information processing systems*, 9, 155-161.

⁷⁸ Tognin, S., Pettersson-Yeo, W., Valli, I., Hutton, C., Woolley, J., Allen, P., ... & Mechelli, A. (2013). Using structural neuroimaging to make quantitative predictions of symptom progression in individuals at ultra-high risk for psychosis. *Frontiers in psychiatry*, 4(187). Doi: 10.3389/fpsy.2013.00187

⁷⁹ Koutsouleris, N., Davatzikos, C., Borgwardt, S., Gaser, C., Bottlender, R., Frodl, T., ... & Meisenzahl, E. (2013). Accelerated brain aging in schizophrenia

and beyond: a neuroanatomical marker of psychiatric disorders. *Schizophrenia bulletin*, 40(5), 1140-1153. Doi: 10.1093/schbul/sbt142

⁸⁰ Franke, K., Ziegler, G., Klöppel, S., Gaser, C., & Alzheimer's Disease Neuroimaging Initiative. (2010). Estimating the age of healthy subjects from T 1-weighted MRI scans using kernel methods: Exploring the influence of various parameters. *Neuroimage*, 50(3), 883-892. Doi: 10.1016/j.neuroimage.2010.01.005

⁸¹ Brown, T. T., Kuperman, J. M., Chung, Y., Erhart, M., McCabe, C., Hagler, D. J., ... & Dale, A. M. (2012). Neuroanatomical assessment of biological maturity. *Current Biology*, 22(18), 1693-1698. Doi: 10.1016/j.cub.2012.07.002

⁸² Koutsouleris, N., Meisenzahl, E. M., Borgwardt, S., Riecher-Rössler, A., Frodl, T., Kambitz, J., ... & Davatzikos, C. (2015). Individualized differential diagnosis of schizophrenia and mood disorders using neuroanatomical biomarkers. *Brain*, awv111. Doi: 10.1093/brain/awv111

⁸³ Schnack, H. G., van Haren, N. E., Nieuwenhuis, M., Pol, H. E. H., Cahn, W., & Kahn, R. S. (2016). Accelerated Brain Aging in Schizophrenia: A Longitudinal Pattern Recognition Study. *American Journal of Psychiatry*.

⁸⁴ Sato, J. R., Salum, G. A., Gadelha, A., Crossley, N., Vieira, G., Manfro, G. G., ... & Bressan, R. A. (2015). Default mode network maturation and psychopathology in children and adolescents. *Journal of Child Psychology and Psychiatry*, 51(1). Doi: 10.1111/jcpp.12444

⁸⁵ Fusar-Poli, P., Bonoldi, I., Yung, A. R., Borgwardt, S., Kempton, M. J., Valmaggia, L., ... & McGuire, P. (2012). Predicting psychosis: meta-analysis of

transition outcomes in individuals at high clinical risk. *Archives of general psychiatry*, 69(3), 220-229.

⁸⁶ Pettersson-Yeo, W., Benetti, S., Marquand, A. F., Joles, R., Catani, M., Williams, S. C., ... & Mechelli, A. (2014). An empirical comparison of different approaches for combining multimodal neuroimaging data with support vector machine. *Frontiers in neuroscience*, 8(189). Doi: 10.3389/fnins.2014.00189

⁸⁷ Modinos, G., Allen, P., Grace, A. A., & McGuire, P. (2015). Translating the MAM model of psychosis to humans. *Trends in neurosciences*, 38(3), 129-138. Doi: 10.1016/j.tins.2014.12.005

⁸⁸ Stone, J. M., Day, F., Tsagaraki, H., Valli, I., McLean, M. A., Lythgoe, D. J., ... & McGuire, P. K. (2009). Glutamate dysfunction in people with prodromal symptoms of psychosis: relationship to gray matter volume. *Biological psychiatry*, 66(6), 533-539. Doi: 10.1016/j.biopsych.2009.05.006

⁸⁹ Day, M., Langston, R., & Morris, R. G. (2003). Glutamate-receptor-mediated encoding and retrieval of paired-associate learning. *Nature*, 424(6945), 205-209. Doi:10.1038/nature01769

⁹⁰ Valli, I., Stone, J., Mechelli, A., Bhattacharyya, S., Raffin, M., Allen, P., ... & McGuire, P. (2011). Altered medial temporal activation related to local glutamate levels in subjects with prodromal signs of psychosis. *Biological psychiatry*, 69(1), 97-99. Doi: 10.1016/j.biopsych.2010.08.033

⁹¹ Fusar-Poli, P., Stone, J. M., Broome, M. R., Valli, I., Mechelli, A., McLean, M. A., ... & McGuire, P. K. (2011). Thalamic glutamate levels as a predictor of cortical response during executive functioning in subjects at high risk for

psychosis. *Archives of general psychiatry*, 68(9), 881-890. Doi: 10.1001/archgenpsychiatry.2011.46

⁹² Stone, J. M., Howes, O. D., Egerton, A., Kambeitz, J., Allen, P., Lythgoe, D. J., ... & McGuire, P. (2010). Altered relationship between hippocampal glutamate levels and striatal dopamine function in subjects at ultra high risk of psychosis. *Biological psychiatry*, 68(7), 599-602. Doi: 10.1016/j.biopsych.2010.05.034

⁹³ Fusar-Poli, P., Howes, O. D., Allen, P., Broome, M., Valli, I., Asselin, M. C., ... & McGuire, P. K. (2010). Abnormal frontostriatal interactions in people with prodromal signs of psychosis: a multimodal imaging study. *Archives of General Psychiatry*, 67(7), 683-691. Doi: doi:10.1001/archgenpsychiatry.2010.77

⁹⁴ Lodge, D. J., & Grace, A. A. (2011). Developmental pathology, dopamine, stress and schizophrenia. *International Journal of Developmental Neuroscience*, 29(3), 207-213. Doi: 10.1016/j.ijdevneu.2010.08.002

⁹⁵ Allen, P., Chaddock, C. A., Egerton, A., Howes, O. D., Barker, G., Bonoldi, I., ... & McGuire, P. (2015). Functional outcome in people at high risk for psychosis predicted by thalamic glutamate levels and prefronto-striatal activation. *Schizophrenia bulletin*, 41(2), 429-439. Doi: 10.1093/schbul/sbu115

⁹⁶ Zhang, D., Wang, Y., Zhou, L., Yuan, H., Shen, D., & Alzheimer's Disease Neuroimaging Initiative. (2011). Multimodal classification of Alzheimer's disease and mild cognitive impairment. *Neuroimage*, 55(3), 856-867. Doi: 10.1371/journal.pone.0021896

⁹⁷ Cui, Y., Liu, B., Luo, S., Zhen, X., Fan, M., Liu, T., ... & Jin, J. S. (2011). Identification of conversion from mild cognitive impairment to Alzheimer's disease

using multivariate predictors. *PloS one*, 6(7), e21896. Doi: 10.1371/journal.pone.0021896

⁹⁸ Garibotto, V., Borroni, B., Kalbe, E., Herholz, K., Salmon, E., Holtorf, V., ... & Perani, D. (2008). Education and occupation as proxies for reserve in aMCI converters and AD FDG-PET evidence. *Neurology*, 71(17), 1342-1349. Doi: 10.1212/01.wnl.0000327670.62378.c0

⁹⁹ Nordholm, D., Krogh, J., Mondelli, V., Dazzan, P., Pariante, C., & Nordentoft, M. (2013). Pituitary gland volume in patients with schizophrenia, subjects at ultra high-risk of developing psychosis and healthy controls: a systematic review and meta-analysis. *Psychoneuroendocrinology*, 38(11), 2394-2404.

¹⁰⁰ Shah, J. L., & Malla, A. K. (2015). Much ado about much: stress, dynamic biomarkers and HPA axis dysregulation along the trajectory to psychosis. *Schizophrenia research*, 162(1), 253-260.

¹⁰¹ Borges, S., Gayer-Anderson, C., & Mondelli, V. (2013). A systematic review of the activity of the hypothalamic–pituitary–adrenal axis in first episode psychosis. *Psychoneuroendocrinology*, 38(5), 603-611.

¹⁰² Thompson, K. N., Phillips, L. J., Komesaroff, P., Yuen, H. P., Wood, S. J., Pantelis, C., ... & McGorry, P. D. (2007). Stress and HPA-axis functioning in young people at ultra high risk for psychosis. *Journal of psychiatric research*, 41(7), 561-569.

¹⁰³ Perkins, D. O., Jeffries, C. D., Addington, J., Bearden, C. E., Cadenhead, K. S., Cannon, T. D., ... & Tsuang, M. T. (2014). Towards a psychosis risk blood diagnostic for persons experiencing high-risk symptoms: preliminary results from

the NAPLS project. *Schizophrenia bulletin*, 41(2), 419-428. Doi: 10.1093/schbul/sbu099.

¹⁰⁴ Chan, M. K., Krebs, M. O., Cox, D., Guest, P. C., Yolken, R. H., Rahmoune, H., ... & Niebuhr, D. W. (2015). Development of a blood-based molecular biomarker test for identification of schizophrenia before disease onset. *Translational Psychiatry*, 5, e601. Doi: 10.1038/tp.2015.91

¹⁰⁵ McGuire, P. K., & Frith, C. D. (1996). Disordered functional connectivity in schizophrenia. *Psychological medicine*, 26(04), 663-667.

¹⁰⁶ Stephan, K. E., Friston, K. J., & Frith, C. D. (2009). Dysconnection in schizophrenia: from abnormal synaptic plasticity to failures of self-monitoring. *Schizophrenia bulletin*, 35(3), 509-527. Doi: 10.1093/schbul/sbn176

¹⁰⁷ Pettersson-Yeo, W., Allen, P., Benetti, S., McGuire, P., & Mechelli, A. (2011). Dysconnectivity in schizophrenia: where are we now?. *Neuroscience & Biobehavioral Reviews*, 35(5), 1110-1124.

¹⁰⁸ Carletti, F., Woolley, J. B., Bhattacharyya, S., Perez-Iglesias, R., Poli, P. F., Valmaggia, L., ... & McGuire, P. K. (2012). Alterations in white matter evident before the onset of psychosis. *Schizophrenia bulletin*, 38(6), 1170-1179. Doi: 10.1093/schbul/sbs053

¹⁰⁹ Bloemen, O. J. N., De Koning, M. B., Schmitz, N., Nieman, D. H., Becker, H. E., De Haan, L., ... & van Amelsvoort, T. A. M. J. (2010). White-matter markers for psychosis in a prospective ultra-high-risk cohort. *Psychological medicine*, 40(08), 1297-1304. Doi: 10.1017/S0033291709991711

¹¹⁰ Schmidt, A., & Borgwardt, S. (2013). Abnormal effective connectivity in the psychosis high-risk state. *Neuroimage*, 81, 119-120. Doi: 10.1503/jpn.130102

-
- ¹¹¹ Bullmore, E. T., & Bassett, D. S. (2011). Brain graphs: graphical models of the human brain connectome. *Annual review of clinical psychology*, 7, 113-140.
- ¹¹² Schmidt, A., Diwadkar, V. A., Smieskova, R., Harrisberger, F., Lang, U. E., McGuire, P., ... & Borgwardt, S. (2014). Approaching a network connectivity-driven classification of the psychosis continuum: a selective review and suggestions for future research. *Frontiers in human neuroscience*, 8, 1047. Doi: 10.3389/fnhum.2014.01047
- ¹¹³ Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: uses and interpretations. *Neuroimage*, 52(3), 1059-1069. Doi: 10.1016/j.neuroimage.2009.10.003
- ¹¹⁴ Crossley, N. A., Mechelli, A., Fusar-Poli, P., Broome, M. R., Matthiasson, P., Johns, L. C., ... & McGuire, P. K. (2009). Superior temporal lobe dysfunction and frontotemporal dysconnectivity in subjects at risk of psychosis and in first-episode psychosis. *Human brain mapping*, 30(12), 4129-4137. Doi: 10.1002/hbm.20834
- ¹¹⁵ Stam, C. J. (2014). Modern network science of neurological disorders. *Nature Reviews Neuroscience*, 15(10), 683-695. Doi: 10.1038/nrn3801
- ¹¹⁶ Alexander-Bloch, A. F., Vértes, P. E., Stidd, R., Lalonde, F., Clasen, L., Rapoport, J., ... & Gogtay, N. (2012). The anatomical distance of functional connections predicts brain network topology in health and schizophrenia. *Cerebral cortex*, 23(1), 127-138. Doi: 10.1093/cercor/bhr388

-
- ¹¹⁷ Tomasi, D., Wang, G. J., & Volkow, N. D. (2013). Energetic cost of brain functional connectivity. *Proceedings of the National Academy of Sciences*, *110*(33), 13642-13647. Doi: 10.1073/pnas.1303346110
- ¹¹⁸ Crossley, N. A., Mechelli, A., Vértes, P. E., Winton-Brown, T. T., Patel, A. X., Ginestet, C. E., ... & Bullmore, E. T. (2013). Cognitive relevance of the community structure of the human brain functional coactivation network. *Proceedings of the National Academy of Sciences*, *110*(28), 11583-11588. Doi: 10.1073/pnas.1220826110
- ¹¹⁹ van den Heuvel, M. P., Kahn, R. S., Goñi, J., & Sporns, O. (2012). High-cost, high-capacity backbone for global brain communication. *Proceedings of the National Academy of Sciences*, *109*(28), 11372-11377. Doi: 10.1073/pnas.1203593109
- ¹²⁰ Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., ... & Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage*, *15*(1), 273-289.
- ¹²¹ Zhang, R., Wei, Q., Kang, Z., Zalesky, A., Li, M., Xu, Y., ... & Huang, R. (2014). Disrupted brain anatomical connectivity in medication-naïve patients with first-episode schizophrenia. *Brain Struct. Funct.* *220*(2), 11451159. Doi: 10.1007/s00429-014-0706-z
- ¹²² Anderson, A., & Cohen, M. S. (2013). Decreased small-world functional network connectivity and clustering across resting state networks in schizophrenia: an fMRI classification tutorial. *Frontiers in human neuroscience*, *7*, 520. Doi: 10.3389/fnhum.2013.00520

-
- ¹²³ Crossley, N. A., Mechelli, A., Scott, J., Carletti, F., Fox, P. T., McGuire, P., & Bullmore, E. T. (2014). The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain*, *137*(8), 2382-2395.
Doi:10.1093/brain/awu132
- ¹²⁴ Crossley, N. A., Mechelli, A., Ginestet, C., Rubinov, M., Bullmore, E. T., & McGuire, P. (in press). Altered hub functioning and compensatory activations in the connectome: a meta-analysis of functional neuroimaging studies in schizophrenia. *Schizophrenia Bulletin*. Doi: 10.1093/schbul/sbv146
- ¹²⁵ Lord, L. D., Allen, P., Expert, P., Howes, O., Lambiotte, R., McGuire, P., ... & Turkheimer, F. E. (2011). Characterization of the anterior cingulate's role in the at-risk mental state using graph theory. *Neuroimage*, *56*(3), 1531-1539.
Doi:10.1016/j.neuroimage.2011.02.012
- ¹²⁶ Foggia, P., Percannella, G., & Vento, M. (2014). Graph matching and learning in pattern recognition in the last 10 years. *International Journal of Pattern Recognition and Artificial Intelligence*, *28*(01), 1450001. Doi: 10.1142/S0218001414500013
- ¹²⁷ Sacchet, M. D., Prasad, G., Foland-Ross, L. C., Thompson, P. M., & Gotlib, I. H. (2015). Support vector machine classification of major depressive disorder using diffusion-weighted neuroimaging and graph theory. *Frontiers in psychiatry*, *6*. Doi: 10.3389/fpsy.2015.00021
- ¹²⁸ Joules, R., Doyle, O. M., Schwarz, A. J., O'Daly, O. G., Brammer, M., Williams, S. C., & Mehta, M. A. (2015). Ketamine induces a robust whole-brain connectivity pattern that can be differentially modulated by drugs of different

mechanism and clinical profile. *Psychopharmacology*, 232(21-22), 4205-4218.

Doi: 10.1007/s00213-015-3951-9

¹²⁹ Fu, Y., & Ma, Y. (Eds.). (2012). *Graph embedding for pattern analysis*.

Springer Science & Business Media.

¹³⁰ Richiardi, J., Achard, S., Bunke, H., & Van De Ville, D. (2013). Machine learning with brain graphs: predictive modeling approaches for functional imaging in systems neuroscience. *Signal Processing Magazine, IEEE*, 30(3), 58-70. Doi:

10.1109/MSP.2012.2233865

¹³¹ Vento, M. (2015). A long trip in the charming world of graphs for pattern recognition. *Pattern Recognition*, 48(2), 291-301. Doi:

10.1016/j.patcog.2014.01.002