King's Research Portal

DOI:

[10.5455/ijtrr.00000020](https://doi.org/10.5455/ijtrr.00000020)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Chattopadhyay, K., Douiri, A., Sheldenkar, A., Wolfe, C. D. A., Rudd, A. G., & Chen, R. (2013). Trends in Activities of Daily Living among Stroke Survivors: Analysis from the South London Stroke Register. *International Journal of Rehabilitation Research*, 2(2), 6-21. <https://doi.org/10.5455/ijtrr.00000020>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

International Journal of
Therapies and Rehabilitation Research
<http://www.scopemed.org/?jid=12>

Trends in Activities of Daily Living among Stroke Survivors: Analysis from the South London Stroke Register

Kaushik Chattopadhyay^{1*}, Abdel Douiri¹, Anita Sheldenkar¹, Anthony G. Rudd^{1 3},
Charles D.A. Wolfe^{1 2}, Ruoling Chen¹

¹Division of Health and Social Care Research, King's College London, London, UK.

²National Institute for Health Research (NIHR) Biomedical Research Centre, Guy's and St. Thomas' NHS Foundation Trust, London, UK.

³Department of Elderly Care, St. Thomas' Hospital, Guy's and St. Thomas' NHS Foundation Trust, London, UK.

Corresponding Author: Kaushik Chattopadhyay*, 7th Floor, Capital House, 42 Weston Street, Division of Health and Social Care Research, King's College London, London, UK, SE1 3QD, Email: kaushik.chattopadhyay@members.em-a.eu

ABSTRACT

Background National and international acute stroke care guidelines came into effect during the last decade to improve outcomes after stroke but their impact on activities of daily living (ADL) improvement over time is not known. The aim of the study was to examine post-stroke ADL trends over time in a multiethnic population in England, and to examine these trends in different socio-economic groups.

Methods Data from the South London Stroke Register were analysed from 1995 to 2011. At 3 months and 1 year post-stroke, basic and instrumental ADL were measured using Barthel Index (poor outcome- BI score<15) and Frenchay Activities Index (poor outcome- FAI score<=15), respectively. Simple and multiple logistic regression analyses were performed.

Results At 3 months post-stroke, the prevalence of poor basic ADL reduced significantly from 33.4% in 1995-1998 to 25.1% in 2008-2011 (trend $p<0.001$) and poor instrumental ADL declined significantly from 59.8% to 53.1% (trend $p=0.005$). The corresponding figures at 1 year were: from 27.8% to 24.3% (trend $p=0.001$) and from 51.6% to 42.8% (trend $p=0.004$). At 3 months, significant reduction in poor ADL was observed over time in the first (least deprived) and second Index of Multiple Deprivation (IMD) tertiles (trend $p=0.006$ and 0.001 , respectively in poor basic ADL; 0.019 and 0.047 , respectively in poor instrumental ADL). At 1 year, poor basic ADL declined significantly in the first and third IMD tertiles (trend $p=0.002$ and 0.043 , respectively), whereas poor instrumental ADL reduced significantly only in the first IMD tertile (trend $p=0.05$).

Conclusion ADL has improved over time among stroke survivors. This may reflect the effectiveness of acute stroke care. Disparities in ADL improvement still exist in different socio-economic groups, and health inequality needs to be tackled.

Key words Activities of daily living (ADL), Stroke, Time trends

INTRODUCTION

Stroke is a major chronic illness with high morbidity rates. During the last decade, improved survival after stroke has led to an increase in its prevalence in the United Kingdom (UK)¹. In England, stroke is the largest cause of major adult disability, leading to severe and moderate disabilities in around 300,000 people². Activities of daily living (ADL) can be classified into two: basic [primary level of activities which are necessary for daily living (self-care tasks)]³ and instrumental (not necessary for basic functioning, but allow a person to live independently within a community)⁴. Stroke can have a major negative impact on a patient's performance of ADL. The issue of poor ADL after stroke is of major importance, as this places a burden on the individual, family, community and health services⁵.

Stroke survivors have been reported as showing improvements in basic and instrumental ADL from stroke onset to 3 months and 1 year post-stroke⁶⁻¹⁰. A number of studies have reported predictors of poor ADL at 3 months and 1 year post-stroke such as gender, age of stroke onset, marital status, living conditions, pre-stroke basic ADL, hemiparesis, motor impairment, urinary incontinence, visuoperceptual deficit, communication problem, intelligence/memory impairment and total anterior circulation infarction^{8,10-13}. In general, socio-economic status is found to be associated with ADL¹⁴, and further research on ADL trends over time in different socio-economic groups has been recommended^{6,15}.

During the last decade, a number of national guidelines especially on acute stroke care (National Service Framework for Older People-2001, National Audit Office Report-2005 and 2010, National Stroke Strategy-2007, and National Clinical Guidelines for Stroke-2008) came into effect to improve outcomes after stroke^{2,16-19}. Other international guidelines also addressed this issue of improvement of outcomes after stroke²⁰⁻²¹. However, their

impact on ADL improvement over time is not known. Evidence suggests that no study has examined post-stroke ADL trends over time. The aim of the study was to examine post-stroke ADL trends over time in a multiethnic population in England, and to examine these trends in different socio-economic groups.

MATERIAL AND METHODS

Study population, patient notification and data collection

The South London Stroke Register (SLSR) is a prospective population-based stroke register set up in January 1995. The total source population of the SLSR area was 271,817 individuals, self-reported as 63% white, 28% black, and 9% other ethnic groups in the 2001 census²².

A stroke was defined as rapidly developing clinical signs of focal (at times global) disturbance of cerebral function, lasting more than 24 hours or leading to death with no apparent cause other than that of vascular origin²³. Patients with first-ever stroke in between 1st January 1995 and 31st December 2011, of all ages, residing in one of the 22 electoral wards of Lambeth and Southwark (an inner area of south London) and who gave written informed consent or assent to participate in the study were included. Exclusion criteria included patients with stroke recurrence, residing outside these south London wards and who refused to give written informed consent or assent. Multiple overlapping sources of notification were used to identify patients by specially trained doctors, nurses and field workers (hospital and community surveillance). After receiving notification of a stroke patient, he/she was contacted for registration within 48 hours where possible. The estimated completeness of case ascertainment in this population has been approximately 88% through a multinomial-logit capture recapture model²⁴. Stroke severity at the time of maximum impairment was: 15.8%, 11.4% and 68.9% of patients had severe, moderate and mild unconsciousness [Glasgow coma scale

(GCS) scores stratified into three: severe (3-8), moderate (9-12) and mild (13-15)]¹⁹, respectively; and 41.7% and 35.5% of patients had urinary incontinence and dysphagia, respectively.

Structured questionnaires were used for initial registration and for follow-ups. The follow-ups were performed at 3 months and annually thereafter through face-to-face or through postal and telephonic interviews, depending on the capability and availability of the patient to complete the questionnaire. At follow-ups, Barthel Index (BI) and Frenchay Activities Index (FAI) were completed. BI measures basic ADL (self care tasks) through 10 questions, namely feeding, bathing, grooming, dressing, bowels, bladder, toilet, transfers (bed to chair and back), mobility (on level surfaces) and stairs. BI is a valid and reliable stroke research tool with scores ranging from 0 to 20^{25,26}. Participants were categorised into one of the two groups: poor basic ADL (BI<15: severely/moderately inactive) and good basic ADL (BI≥15: mildly inactive/independent)²⁷. FAI measures instrumental ADL, that is, social activities and more complex ADL (such as outdoor mobility, domestic chores, leisure and gainful work) through 15 questions. This is a valid and reliable tool with scores ranging from 0 to 45²⁸⁻³¹. Participants were categorised into one of the two groups: poor instrumental ADL (FAI≤15: inactive) and good instrumental ADL (FAI>15: moderately active/independent)³¹. All cut-off points were pre-defined in order to determine poor outcomes.

The initial assessment included information on socio-demographics: age of stroke onset, sex, self-defined ethnicity, postcode [to determine Index of Multiple Deprivation (IMD) scores-categorised into tertiles (first tertile=least socio-economically deprived and third tertile=most socio-economically deprived)], pre-stroke living conditions, pre-stroke basic ADL; pre-stroke risk factors: smoking, alcohol drinking, hypertension, migraine, myocardial infarction, transient

ischemic attack, diabetes mellitus, atrial fibrillation, peripheral vascular disease; pathological stroke subtype classification; stroke severity at the time of maximum impairment: consciousness level using GCS, dysphagia, urinary incontinence; and processes of acute stroke care: hospital admission, admission/transfer to a stroke unit, more than 50% of hospital admission spent in a stroke unit, brain imaging [computed tomography (CT) or magnetic resonance imaging (MRI)]. These indicators of the processes of acute stroke care are some of the useful proxy measures for the overall quality of stroke care¹⁹.

Ethics

The study was ethically approved by the research ethics committees of Guy's and St. Thomas' National Health Service (NHS) Foundation Trust, King's College Hospital NHS Foundation Trust, National Hospital for Neurology and Neurosurgery (University College London Hospitals NHS Foundation Trust), St. George's Healthcare NHS Trust, and Chelsea and Westminster Hospital NHS Foundation Trust. Information sheets were distributed among all the eligible patients or their relatives, and written informed consents or assents were taken from those interested in participating in the research.

Statistical analyses

The chi-square test was used to investigate the association between ADL and categorical variables (socio-demographics, pre-stroke risk factors, stroke subtype, stroke severity and processes of acute stroke care). Seventeen years of SLSR data were divided into five year groups: 1995-1998, 1999-2001, 2002-2004, 2005-2007, and 2008-2011. Simple and multiple logistic regression analyses were performed to examine trends over time in the prevalence of poor ADL among stroke survivors in these five year groups, and to determine these trends in different socio-economic groups (IMD tertiles).

In multiple regression models, adjustments were performed for all the categorical variables. Sensitivity analyses were carried out using the 1st four year groups (1995-2007) and excluding the 5th year group (2008-2011) to examine these trends before relevant guidelines came into effect. Multiple regression models included a sample with missing values for these adjusted variables. Odds ratios (ORs), 95% confidence intervals (CIs) and P values were reported. All data were analyzed using STATA-12 for Windows software³².

RESULTS

Data collected on 4,413 first-ever stroke patients between 1st January 1995 and 31st December 2011 were analysed. Figure 1 showed the number of stroke survivors with basic and instrumental ADL measured at 3 months and 1 year post-stroke. 1,107 (25.1%) patients died between stroke onset and 3 months follow-up, and 329 (7.5%) patients died between 3 months and 1 year follow-ups. At 3 months, 639 (30.5%) of the 2,094 survivors measured with BI had poor basic ADL, and 1,101 (56.5%) of the 1,948 survivors measured with FAI had poor instrumental ADL. At 1 year, 497 (25.2%) of the 1,971 survivors had poor basic ADL, and 888 (47.3%) of the 1,878 survivors had poor instrumental ADL.

Table 1 reported the characteristics of stroke survivors with poor and good basic and instrumental ADL measured at 3 months and 1 year post-stroke. Age of stroke onset, sex, ethnicity, IMD, pre-stroke living conditions, pre-stroke basic ADL, smoking, alcohol drinking, migraine, atrial fibrillation, stroke subtype, GCS, dysphagia, urinary incontinence, hospital admission and more than 50% of hospital admission spent in a stroke unit were associated with basic ADL at 3 months. All these factors, hypertension, transient ischemic attack, diabetes mellitus and admission/transfer to a stroke unit were associated with instrumental ADL at 3 months. At 1 year, all the 3 months

basic ADL factors (except for IMD and more than 50% of hospital admission spent in a stroke unit), transient ischemic attack and diabetes mellitus were associated with basic ADL. Factors associated with instrumental ADL at 1 year were similar to the 3 months instrumental ADL factors (except for sex, IMD, hypertension and admission/transfer to a stroke unit).

Figure 2 showed the crude poor basic and instrumental ADL measured at 3 months and 1 year post-stroke over seventeen years (1995-2011). At 3 months, the prevalence of poor basic ADL among stroke survivors reduced significantly from 33.4% in 1995-1998 to 25.1% in 2008-2011 (trend $p<0.001$), and poor instrumental ADL declined significantly from 59.8% to 53.1% (trend $p=0.002$). The corresponding figures at 1 year were: from 27.8% to 24.3% (trend $p=0.064$) and from 51.6% to 42.8% (trend $p=0.006$). At 1 year, the prevalence of poor basic ADL among stroke survivors reduced significantly from 1995 to 2007 (trend $p=0.033$).

Figure 3 showed the crude poor basic and instrumental ADL measured at 3 months and 1 year post-stroke in three socio-economic groups over seventeen years (1995-2011). In simple regression analyses, significant reduction in poor ADL at 3 months and 1 year was observed over time only in the first IMD tertile.

Table 2 reported the adjusted poor basic and instrumental ADL measured at 3 months and 1 year post-stroke over seventeen years and in three socio-economic groups (1995-2011). The risk of poor basic ADL at 3 months and 1 year reduced significantly from 1995 to 2011. At 3 months, adjusted ORs (95% CIs) were 1.15 (0.69-1.94) in 1999-2001, 0.59 (0.39-0.88) in 2002-2004, 0.72 (0.46-1.12) in 2005-2007 and 0.62 (0.39-0.98) in 2008-2011 as compared to 1995-1998 (trend $p<0.001$). Similarly at 1 year, the corresponding figures were 0.93 (0.58-1.51), 0.47 (0.31-0.71), 0.59 (0.38-0.91) and 0.64 (0.40-

Table 1- Characteristics of stroke survivors with poor and good basic and instrumental ADL measured at 3 months and 1 year post-stroke: SLSR 1995-2011												
	3 months						1 year					
	BI (n=2094)			FAI (n=1948)			BI (n=1971)			FAI (n=1878)		
	BI<15 (n=639)	BI>=15 (n=1455)	P value	FAI<=15 (n=1101)	FAI>15 (n=847)	P value	BI<15 (n=497)	BI>=15 (n=1474)	P value	FAI<=15 (n=888)	FAI>15 (n=990)	P value
Socio-demographics												
Age of stroke onset (years)			<0.001			<0.001			<0.001			<0.001
0-64	132 (20.7)	564 (38.8)		284 (25.8)	368 (43.4)		113 (22.7)	602 (40.8)		227 (25.6)	458 (46.3)	
65-74	173 (27.1)	421 (28.9)		300 (27.2)	247 (29.2)		137 (27.6)	432 (29.3)		264 (29.7)	286 (28.9)	
75-84	201 (31.5)	358 (24.6)		327 (29.7)	192 (22.7)		158 (31.8)	358 (24.3)		277 (31.2)	206 (20.8)	
85+	133 (20.8)	112 (7.7)		190 (17.3)	40 (4.7)		89 (17.9)	82 (5.6)		120 (13.5)	40 (4.0)	
Sex			<0.001			0.006			<0.001			0.915
Male	280 (43.8)	823 (56.6)		549 (49.9)	476 (56.2)		224 (45.1)	828 (56.2)		475 (53.5)	532 (53.7)	
Female	359 (56.2)	632 (43.4)		552 (50.1)	371 (43.8)		273 (54.9)	646 (43.8)		413 (46.5)	458 (46.3)	
Ethnicity			0.048			<0.001			0.005			<0.001
White	442 (69.2)	1023 (70.3)		736 (66.8)	629 (74.3)		349 (70.2)	1006 (68.2)		596 (67.1)	701 (70.8)	
Black	142 (22.2)	340 (23.4)		276 (25.1)	167 (19.7)		100 (20.1)	367 (24.9)		202 (22.7)	235 (23.7)	
Other	53 (8.3)	79 (5.4)		85 (7.7)	41 (4.8)		46 (9.3)	85 (5.8)		84 (9.5)	42 (4.2)	
Unknown	2 (0.3)	13 (0.9)		4 (0.4)	10 (1.2)		2 (0.4)	16 (1.1)		6 (0.7)	12 (1.2)	
IMD			0.044			0.021			0.3			0.06
1 st tertile	190 (29.7)	514 (35.3)		345 (31.3)	313 (37.0)		155 (31.2)	512 (34.7)		276 (31.1)	359 (36.3)	
2 nd tertile	223 (34.9)	477 (32.8)		364 (33.1)	275 (32.5)		177 (35.6)	481 (32.6)		306 (34.5)	316 (31.9)	
3 rd tertile	224 (35.1)	461 (31.7)		389 (35.3)	259 (30.6)		165 (33.2)	481 (32.6)		306 (34.5)	315 (31.8)	
Unknown	2 (0.3)	3 (0.2)		3 (0.3)	0		0	0		0	0	
Pre-stroke living conditions			<0.001			<0.001			<0.001			<0.001
Home alone	181 (28.3)	471 (32.4)		307 (27.9)	297 (35.1)		142 (28.6)	464 (31.5)		246 (27.7)	325 (32.8)	
Home with someone	300 (46.9)	751 (51.6)		563 (51.1)	425 (50.2)		236 (47.5)	773 (52.4)		456 (51.4)	507 (51.2)	
Institution	73 (11.4)	70 (4.8)		107 (9.7)	27 (3.2)		56 (11.3)	62 (4.2)		77 (8.7)	34 (3.4)	
Unknown	85 (13.3)	163 (11.2)		124 (11.3)	98 (11.6)		63 (12.7)	175 (11.9)		109 (12.3)	124 (12.5)	
Pre-stroke basic ADL			<0.001			<0.001			<0.001			<0.001

Good (BI $\geq$ 15)	557 (87.2)	1444 (99.2)		1021 (92.7)	842 (99.4)		440 (88.5)	1463 (99.3)		827 (93.1)	987 (99.7)	
Poor (BI<15)	82 (12.8)	11 (0.8)		80 (7.3)	5 (0.6)		57 (11.5)	11 (0.7)		61 (6.9)	3 (0.3)	
Pre-stroke risk factors												
Smoking			0.001			0.002			0.002			0.047
Non-smoker	442 (69.2)	918 (63.1)		748 (67.9)	524 (61.9)		336 (67.6)	904 (61.3)		577 (65.0)	605 (61.1)	
Current smoker	175 (27.4)	516 (35.5)		327 (29.7)	311 (36.7)		144 (29.0)	546 (37.0)		290 (32.7)	369 (37.3)	
Unknown	22 (3.4)	21 (1.4)		26 (2.4)	12 (1.4)		17 (3.4)	24 (1.6)		21 (2.4)	16 (1.6)	
Alcohol drinking			<0.001			<0.001			0.001			<0.001
No	283 (44.3)	488 (33.5)		444 (40.3)	271 (32.0)		198 (39.8)	485 (32.9)		349 (39.3)	303 (30.6)	
Yes	304 (47.6)	894 (61.4)		587 (53.3)	536 (63.3)		257 (51.7)	898 (60.9)		485 (54.6)	617 (62.3)	
Unknown	52 (8.1)	73 (5.0)		70 (6.4)	40 (4.7)		42 (8.5)	91 (6.2)		54 (6.1)	70 (7.1)	
Hypertension			0.614			0.001			0.371			0.147
No	205 (32.1)	486 (33.4)		322 (29.2)	307 (36.2)		155 (31.2)	488 (33.1)		277 (31.2)	337 (34.0)	
Yes	413 (64.6)	930 (63.9)		750 (68.1)	515 (60.8)		328 (66.0)	934 (63.4)		585 (65.9)	616 (62.2)	
Unknown	21 (3.3)	39 (2.7)		29 (2.6)	25 (3.0)		14 (2.8)	52 (3.5)		26 (2.9)	37 (3.7)	
Migraine			0.014			0.011			0.004			0.001
No	567 (88.7)	1272 (87.4)		981 (89.1)	732 (86.4)		452 (90.9)	1283 (87.0)		803 (90.4)	854 (86.3)	
Yes	29 (4.5)	110 (7.6)		60 (5.4)	71 (8.4)		18 (3.6)	107 (7.3)		39 (4.4)	80 (8.1)	
Unknown	43 (6.7)	73 (5.0)		60 (5.4)	44 (5.2)		27 (5.4)	84 (5.7)		46 (5.2)	56 (5.7)	
Myocardial infarction			0.185			0.102			0.473			0.063
No	541 (84.7)	1271 (87.4)		944 (85.7)	747 (88.2)		429 (86.3)	1288 (87.4)		766 (86.3)	869 (87.8)	
Yes	73 (11.4)	140 (9.6)		122 (11.1)	75 (8.9)		52 (10.5)	138 (9.4)		99 (11.1)	84 (8.5)	
Unknown	25 (3.9)	44 (3.0)		35 (3.2)	25 (3.0)		16 (3.2)	48 (3.3)		23 (2.6)	37 (3.7)	
Transient ischemic attack			0.103			0.005			0.004			<0.001
No	521 (81.5)	1233 (84.7)		900 (81.7)	728 (86.0)		397 (79.9)	1253 (85.0)		717 (80.7)	859 (86.8)	
Yes	97 (15.2)	184 (12.6)		172 (15.6)	95 (11.2)		83 (16.7)	172 (11.7)		145 (16.3)	94 (9.5)	
Unknown	21 (3.3)	38 (2.6)		29 (2.6)	24 (2.8)		17 (3.4)	49 (3.3)		26 (2.9)	37 (3.7)	
Diabetes mellitus			0.189			<0.001			0.006			<0.001
No	490 (76.7)	1156 (79.5)		835 (75.8)	697 (82.3)		373 (75.1)	1175 (79.7)		668 (75.2)	802 (81.0)	
Yes	130 (20.3)	262 (18.0)		239 (21.7)	126 (14.9)		113 (22.7)	250 (17.0)		198 (22.3)	152 (15.4)	

Unknown	19 (3.0)	37 (2.5)		27 (2.5)	24 (2.8)		11 (2.2)	49 (3.3)		22 (2.5)	36 (3.6)	
Atrial fibrillation			0.004			0.001			<0.001			0.002
No	511 (80.0)	1248 (85.8)		900 (81.7)	737 (87.0)		395 (79.5)	1281 (86.9)		739 (83.2)	862 (87.1)	
Yes	101 (15.8)	167 (11.5)		167 (15.2)	86 (10.2)		83 (16.7)	146 (9.9)		122 (13.7)	91 (9.2)	
Unknown	27 (4.2)	40 (2.7)		34 (3.1)	24 (2.8)		19 (3.8)	47 (3.2)		27 (3.0)	37 (3.7)	
Peripheral vascular disease			0.944			0.413			0.835			0.517
No	574 (89.8)	1337 (91.9)		1007 (91.5)	772 (91.1)		457 (92.0)	1345 (91.2)		819 (92.2)	897 (90.6)	
Yes	21 (3.3)	48 (3.3)		34 (3.1)	32 (3.8)		15 (3.0)	47 (3.2)		27 (3.0)	35 (3.5)	
Unknown	44 (6.9)	70 (4.8)		60 (5.4)	43 (5.1)		25 (5.0)	82 (5.6)		42 (4.7)	58 (5.9)	
Stroke subtype			0.006			<0.001			0.005			<0.001
Ischemic	536 (83.9)	1176 (80.8)		911 (82.7)	687 (81.1)		413 (83.1)	1200 (81.4)		743 (83.7)	799 (80.7)	
Primary intra-cerebral haemorrhage	70 (11.0)	140 (9.6)		125 (11.4)	69 (8.1)		57 (11.5)	138 (9.4)		103 (11.6)	86 (8.7)	
Subarachnoid haemorrhage	10 (1.6)	55 (3.8)		23 (2.1)	37 (4.4)		6 (1.2)	66 (4.5)		14 (1.6)	51 (5.2)	
Undefined	23 (3.6)	84 (5.8)		42 (3.8)	54 (6.4)		21 (4.2)	70 (4.7)		28 (3.2)	54 (5.5)	
Stroke severity												
GCS			<0.001			<0.001			<0.001			<0.001
Severe (≤ 8)	42 (6.6)	35 (2.4)		57 (5.2)	13 (1.5)		29 (5.8)	47 (3.2)		47 (5.3)	26 (2.6)	
Moderate (9–12)	109 (17.1)	88 (6.0)		137 (12.4)	39 (4.6)		75 (15.1)	92 (6.2)		109 (12.3)	55 (5.6)	
Mild (≥ 13)	463 (72.5)	1305 (89.7)		874 (79.4)	780 (92.1)		381 (76.7)	1304 (88.5)		709 (79.8)	889 (89.8)	
Unknown	25 (3.9)	27 (1.9)		33 (3.0)	15 (1.8)		12 (2.4)	31 (2.1)		23 (2.6)	20 (2.0)	
Dysphagia			<0.001			<0.001			<0.001			<0.001
No	301 (47.1)	1140 (78.4)		652 (59.2)	690 (81.5)		265 (53.3)	1092 (74.1)		526 (59.2)	761 (76.9)	
Yes	307 (48.0)	224 (15.4)		401 (36.4)	93 (11.0)		198 (39.8)	259 (17.6)		309 (34.8)	132 (13.3)	
Unknown	31 (4.9)	91 (6.3)		48 (4.4)	64 (7.6)		34 (6.8)	123 (8.3)		53 (6.0)	97 (9.8)	
Urinary incontinence			<0.001			<0.001			<0.001			<0.001
No	223 (34.9)	1147 (78.8)		557 (50.6)	724 (85.5)		216 (43.5)	1129 (76.6)		465 (52.4)	811 (81.9)	
Yes	389 (60.9)	278 (19.1)		509 (46.2)	105 (12.4)		264 (53.1)	302 (20.5)		392 (44.1)	151 (15.3)	
Unknown	27 (4.2)	30 (2.1)		35 (3.2)	18 (2.1)		17 (3.4)	43 (2.9)		31 (3.5)	28 (2.8)	
Processes of acute stroke care												

Hospital admission			<0.001			<0.001			<0.001			<0.001
No	38 (5.9)	203 (14.0)		75 (6.8)	144 (17.0)		43 (8.7)	236 (16.0)		87 (9.8)	181 (18.3)	
Yes	601 (94.1)	1252 (86.0)		1026 (93.2)	703 (83.0)		454 (91.3)	1238 (84.0)		801 (90.2)	809 (81.7)	
Admission/transfer to a stroke unit			0.052			0.012			0.446			0.744
No	243 (38.0)	449 (30.9)		407 (37.0)	237 (28.0)		168 (33.8)	484 (32.8)		310 (34.9)	306 (30.9)	
Yes	349 (54.6)	787 (54.1)		605 (55.0)	456 (53.8)		279 (56.1)	737 (50.0)		480 (54.1)	490 (49.5)	
Unknown	47 (7.4)	219 (15.1)		89 (8.1)	154 (18.2)		50 (10.1)	253 (17.2)		98 (11.0)	194 (19.6)	
>50% of hospital admission spent in a stroke unit			<0.001			<0.001			0.301			0.009
No	306 (47.9)	516 (35.5)		489 (44.4)	272 (32.1)		203 (40.8)	550 (37.3)		374 (42.1)	374 (42.1)	
Yes	219 (34.3)	625 (43.0)		433 (39.3)	366 (43.2)		192 (38.6)	587 (39.8)		336 (37.8)	336 (37.8)	
Unknown	114 (17.8)	314 (21.6)		179 (16.3)	209 (24.7)		102 (20.5)	337 (22.9)		178 (20.0)	178 (20.0)	
Brain imaging (CT/MRI)			0.977			0.952			0.46			0.647
No	14 (2.2)	31 (2.1)		22 (2.0)	17 (2.0)		11 (2.2)	25 (1.7)		18 (2.0)	17 (1.7)	
Yes	613 (95.9)	1370 (94.2)		1053 (95.6)	798 (94.2)		474 (95.4)	1410 (95.7)		853 (96.1)	942 (95.2)	
Unknown	12 (1.9)	54 (3.7)		26 (2.4)	32 (3.8)		12 (2.4)	39 (2.6)		17 (1.9)	31 (3.1)	
n (%), chi-square test, P value excludes unknown												

Table 2- Adjusted poor basic and instrumental ADL measured at 3 months and 1 year post-stroke over seventeen years and in three socio-economic groups: SLSR 1995-2011										
		1995-1998	1999-2001	2002-2004	2005-2007	2008-2011	Trend P	IMD group	Adjusted OR (95% CI)	Trend P
3 months										
BI<15	OR (95% CI), P	1	1.15 (0.69-1.94), 0.588	0.59 (0.39-0.88), 0.009	0.72 (0.46-1.12), 0.147	0.62 (0.39-0.98), 0.039	<0.001	1 st tertile	0.79 (0.66-0.93)	0.006
								2 nd tertile	0.76 (0.64-0.90)	0.001
								3 rd tertile	0.93 (0.80-1.09)	0.37
FAI<=15	OR (95% CI), P	1	1.15 (0.69-1.93), 0.586	0.93 (0.64-1.34), 0.684	0.64 (0.42-0.96), 0.033	0.78 (0.51-1.20), 0.261	0.005	1 st tertile	0.83 (0.71-0.97)	0.019
								2 nd tertile	0.86 (0.74-1.00)	0.047
								3 rd tertile	0.98 (0.85-1.14)	0.815
1 year										
BI<15	OR (95% CI), P	1	0.93 (0.58-1.51), 0.775	0.47 (0.31-0.71), <0.001	0.59 (0.38-0.91), 0.018	0.64 (0.40-1.01), 0.055	0.001	1 st tertile	0.76 (0.63-0.90)	0.002
								2 nd tertile	0.93 (0.79-1.09)	0.377
								3 rd tertile	0.83 (0.70-0.99)	0.043
FAI<=15	OR (95% CI), P	1	0.73 (0.47-1.15), 0.174	0.76 (0.53-1.09), 0.142	0.78 (0.53-1.16), 0.219	0.79 (0.52-1.20), 0.274	0.004	1 st tertile	0.86 (0.73-1.00)	0.05
								2 nd tertile	0.91 (0.79-1.05)	0.208
								3 rd tertile	0.90 (0.78-1.04)	0.161

Figure 1- Number of stroke survivors with basic and instrumental ADL measured at 3 months and 1 year post-stroke

Figure 2- Crude poor basic and instrumental ADL measured at 3 months and 1 year post-stroke over seventeen years: SLSR 1995-2011

*[Unadjusted OR (95% CI); Trend P]. ^Significant trend for 1995-2007.

Figure 3- Crude poor basic and instrumental ADL measured at 3 months and 1 year post-stroke in three socio-economic groups over seventeen years: SLSR 1995-2011

*[Unadjusted OR (95% CI); Trend P].

1.01) with trend $p=0.001$. At 3 months and 1 year, the risk of poor instrumental ADL declined significantly from 1995 to 2011. At 3 months, adjusted ORs (95% CIs) were 1.15 (0.69-1.93) in 1999-2001, 0.93 (0.64-1.34) in 2002-2004, 0.64 (0.42-0.96) in 2005-2007 and 0.78 (0.51-1.20) in 2008-2011 as compared to 1995-1998 (trend $p=0.005$). Similarly at 1 year, the corresponding figures were 0.73 (0.47-1.15), 0.76 (0.53-1.09), 0.78 (0.53-1.16) and 0.79 (0.52-1.20) with trend $p=0.004$. In multiple regression analyses, significant reduction in poor ADL at 3 months was observed over time among stroke survivors in the first and second IMD tertiles (adjusted trend $p=0.006$ and 0.001 , respectively in poor basic ADL; 0.019 and 0.047, respectively in poor instrumental ADL). At 1 year, poor basic ADL declined significantly among stroke survivors in the first and third IMD tertiles (adjusted trend $p=0.002$ and 0.043 , respectively), and poor instrumental ADL reduced significantly only in the first IMD tertile (adjusted trend $p=0.05$).

DISCUSSION

ADL has improved over time among stroke survivors in this population-based cohort study. However, disparities in ADL improvement still exist in different socio-economic groups. To the best of knowledge, this is the first study to report ADL trends over time among stroke survivors. A number of ADL time trend studies have been conducted especially among older people (general studies but not specific to stroke) using different methodologies (such as different case-definitions, measuring tools and cut-off points)³³⁻³⁹.

The risk of poor ADL at 3 months and 1 year post-stroke decreased significantly from 1995 to 2011. The prevalence of poor ADL among older people declined over time in the UK and in the United States of America (USA)³³⁻³⁸. However, another research conducted among older people in England documented an increase in the crude poor ADL over time³³, but also reported a decline in the moderate ADL

limitation over the same time period³⁹. The post-stroke ADL could be improved through effective stroke management^{15,40}. The receipt of acute stroke care has improved over time in this population⁴¹, and this could explain the post-stroke ADL improvement over time. However, ADL is also improving in the general population over time, and further research is needed to explore all the possible reasons behind this ADL improvement.

Poor ADL at 3 months reduced significantly over time among stroke survivors in the least and second least deprived groups, but not in the most deprived group. At 1 year, poor basic and instrumental ADL declined significantly in the least and most deprived groups, and only in the least deprived group, respectively. In England, the age-standardised poor ADL prevalence declined over time in social classes I, II and V whereas it increased in social class IV³⁹. In the USA, the ADL improved over time among older people in the highest income quintile group, but not in the lowest one³⁶. Non-manual stroke patients in this population were more likely to be admitted to hospitals and stroke units⁴² and receive brain imaging⁴¹ as compared to manual stroke patients, and this could explain the post-stroke ADL improvement over time in the least deprived group. Poor basic ADL at 1 year also decreased significantly over time in the most deprived group, which could be due to other confounding factors (such as stroke recurrence) that were not adjusted for in this study, and needs further research.

This study has a number of strengths and weaknesses. The SLSR is a multiethnic population-based stroke register with a large number of registered patients (hospitalised and non-hospitalised patients), long-term follow-ups and includes a vast number of baseline variables which have never been explored before. Multiple overlapping sources of notification were used to identify patients. A standardised protocol was used for data collection by specially trained doctors, nurses and field

workers. Perhaps, no dataset worldwide would be representative to allow such ADL time trend analyses. The study findings could be generalized to other similar populations with similar healthcare provisions.

ADL measurement was subjective to participants and thus, valid and reliable tools were used to measure ADL. However, objectively measured ADL trends over time could be different, and further studies need to be carried out. The focus of this study was at 3 months and 1 year after stroke (1995-2011), and further research that focuses on more than a year is needed. However, the outcomes remain comparatively stable after 1 year of stroke⁶. This study provided a better picture of the after-effects of acute stroke care on ADL improvement at 3 months and 1 year post-stroke, and included both severe and milder stroke survivors.

All the structured questionnaires had been amended regularly to reflect changes in evidence, and some of the processes of acute stroke care variables were added in 2005. These variables could not be used, as the available data were limited for analyses. However, four indicators of the processes of acute stroke care were included, which are some of the useful proxy measures for the overall quality of stroke care¹⁹. Moreover, if a patient has been admitted/transferred to a stroke unit in London, it can be assumed that the patient will receive adequate care from the stroke-skilled multidisciplinary team (which includes specialist stroke physician, physiotherapist, occupational therapist, speech and language therapist, dietician, psychologist and social worker) and appropriate secondary prevention drugs.

Missing data could lead to bias, but it was generally low in this study. Multiple regression analyses included a sample with missing values for the adjusted variables. The loss to follow-up could lead to bias, and this could be an issue in some socio-demographic groups. People who

are healthier or wealthier may be more likely to participate in research follow-ups⁴³. However, any such group in the analyses were not found. Unlike some other cohort and stroke register studies, loss to follow-up was reported. Some patients were registered retrospectively and thus, 3 months and 1 year follow-ups could not be performed. This rate was higher at 3 months as compared to 1 year. The south London population is quite mobile, with a large number of migrants. This increases the chances of loss to follow-up, however every attempt was made to keep a record of the latest contact details of the patient (through general practitioners, hospitals or relatives). In case the patient migrated to an area outside London, postal follow-up was completed.

CONCLUSION

ADL has improved over time among stroke survivors. This may reflect the effectiveness of acute stroke care. However, disparities in ADL improvement still exist in different socio-economic groups, and health inequality needs to be tackled.

ACKNOWLEDGEMENTS

The study was funded by the NIHR Programme grant (RP-PG-0407-10184). The funders had no role in the study design, data collection and analysis or preparation of the manuscript. We wish to thank our patients, their relatives and healthcare/research professionals involved.

CONFLICT OF INTEREST

None

REFERENCES

1. Lee S, Shafe AC, Cowie MR. UK stroke incidence, mortality and cardiovascular risk management 1999–2008: time-trend analysis from the General Practice Research Database. *BMJ Open*. 2011;1(2):e000269.

2. National Audit Office. Department of Health: Progress in Improving Stroke Care. London: The Stationery Office. 2010.
3. McDowell I and Newell C. Measuring Health: A Guide to Rating Scales and Questionnaires. 2nd Edition. New York: Oxford University Press. 1996.
4. Bookman A, Harrington M, Pass L, Reisner E. The Family Caregiver Handbook: Finding Eldercare Resources in Massachusetts. Cambridge, MA: Massachusetts Institute of Technology. 2007.
5. Power KM. Creating a Culture for Rehabilitation in the Acute Stroke Unit (dissertation). Canada: University of Victoria. 2008.
6. Wolfe CD, Crichton SL, Heuschmann PU, McKeivitt CJ, Toschke AM, Grieve AP, Rudd AG. Estimates of outcomes up to ten years after stroke: analysis from the prospective South London Stroke Register. PLoS Med. 2011;8(5):e1001033.
7. Hofgren C, Björkdahl A, Esbjörnsson E, Sunnerhagen KS. Recovery after stroke: cognition, ADL function and return to work. Acta Neurol Scand. 2007;115(2):73-80.
8. Kotila M, Waltimo O, Niemi ML, Laaksonen R, Lempinen M. The profile of recovery from stroke and factors influencing outcome. Stroke. 1984;15(6):1039-44.
9. Patel MD, Tilling K, Lawrence E, Rudd AG, Wolfe CD, McKeivitt C. Relationships between long-term stroke disability, handicap and health-related quality of life. Age Ageing. 2006;35(3):273-9.
10. Schepers VP, Visser-Meily AM, Ketelaar M, Lindeman E. Prediction of social activity 1 year poststroke. Arch Phys Med Rehabil. 2005;86(7):1472-6.
11. Veerbeek JM, Kwakkel G, Van Wegen EE, Ket JC, Heymans MW. Early prediction of outcome of activities of daily living after stroke: a systematic review. Stroke. 2011;42(5):1482-8.
12. Patel M, Coshall C, Lawrence E, Rudd AG, Wolfe CD. Recovery from poststroke urinary incontinence: associated factors and impact on outcome. J Am Geriatr Soc. 2001;49(9):1229-33.
13. Sturm JW, Dewey HM, Donnan GA, Macdonell RA, McNeil JJ, Thrift AG. Handicap after stroke: how does it relate to disability, perception of recovery, and stroke subtype?: the North East Melbourne Stroke Incidence Study (NEMESIS). Stroke. 2002;33(3):762-8.
14. Fuller-Thomson E, Gadalla T. Income inequality and limitations in activities of daily living: a multilevel analysis of the 2003 American Community Survey. Public Health. 2008;122(3):221-8.
15. Manton KG. Recent declines in chronic disability in the elderly U.S. population: risk factors and future dynamics. Annu Rev Public Health. 2008;29:91-113.
16. Department of Health. National Service Framework for Older People. London: Department of Health. 2001.
17. National Audit Office. Department of Health: Reducing Brain Damage- Faster Access to Better Stroke Care. London: The Stationery Office. 2005.
18. Department of Health. National Stroke Strategy. London: Department of Health. 2007.
19. Royal College of Physicians Intercollegiate Stroke Working Party. National Clinical Guidelines for Stroke. 3rd Edition. London: Royal College of Physicians. 2008.
20. Kjellstrom T, Norrving B, Shatchkute A. Helsingborg declaration 2006 on European stroke strategies. Cerebrovasc Dis. 2007;23(2-3):231-41.
21. European Stroke Organisation (ESO) Executive Committee, ESO Writing Committee. Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. Cerebrovasc Dis. 2008;25(5):457-507.
22. Heuschmann PU, Grieve AP, Toschke AM, Rudd AG, Wolfe CD. Ethnic group disparities in 10-year trends in stroke incidence and vascular risk factors: the South London Stroke Register (SLSR). Stroke. 2008;39(8):2204-10.
23. Hatano S. Experience from a multicentre stroke register: a preliminary report. Bull World Health Organ. 1976;54(5):541-53.
24. Tilling K, Sterne JA, Wolfe CD. Estimation of the incidence of stroke using a capture-

- recapture model including covariates. *Int J Epidemiol.* 2001;30(6):1351-9.
25. Mahoney FI, Barthel DW. Functional evaluation: the Barthel Index. *Md State Med J.* 1965;14:61-5.
 26. Collin C, Wade D, Davies S, Horne V. The Barthel ADL Index: a reliability study. *Int Disabil Stud.* 1988;10(2):61-3.
 27. Wolfe CD, Taub NA, Woodrow EJ, Burney PG. Assessment of scales of disability and handicap for stroke patients. *Stroke.* 1991;22(10):1242-4.
 28. Wade DT, Legh-Smith J, Langton Hewer R. Social activities after stroke: measurement and natural history using Frenchay Activities Index. *Int Rehabil Med.* 1985;7(4):176-81.
 29. Schuling J, de Haan R, Limburg M, Groenier KH. The Frenchay Activities Index: assessment of functional status in stroke patients. *Stroke.* 1993;24(8):1173-7.
 30. Tooth LR, McKenna KT, Smith M, O'Rourke P. Further evidence for the agreement between patients with stroke and their proxies on the Frenchay Activities Index. *Clin Rehabil.* 2003;17(6):656-65.
 31. Anderson CS, Jamrozik KD, Broadhurst RJ, Stewart-Wynne EG. Predicting survival for 1 year among different subtypes of stroke. Results from the Perth Community Stroke Study. *Stroke.* 1994;25(10):1935-44.
 32. STATACorp. STATA Statistical Software: Release 12. College Station, TX: STATACorp LP. 2011.
 33. Lafortune G, Balestat G and the Disability Study Expert Group Members. Trends in Severe Disability among Older People: Assessing the Evidence in 12 OECD Countries and the Future Implications. OECD Health Working Papers No 26. Paris: OECD Publications Service. 2007.
 34. Donald IP, Foy C, Jagger C. Trends in disability prevalence over 10 years in older people living in Gloucestershire. *Age Ageing.* 2010;39(3):337-42.
 35. Freedman VA, Martin LG, Schoeni RF. Recent trends in disability and functioning among older adults in the United States: a systematic review. *JAMA.* 2002;288(24):3137-46.
 36. Schoeni RF, Martin LG, Andreski PM, Freedman VA. Persistent and growing socioeconomic disparities in disability among the elderly: 1982-2002. *Am J Public Health.* 2005;95(11):2065-70.
 37. Manton KG, Gu X, Lamb VL. Change in chronic disability from 1982 to 2004/2005 as measured by long-term changes in function and health in the U.S. elderly population. *Proc Natl Acad Sci USA.* 2006;103(48):18374-9.
 38. Martin LG, Schoeni RF, Freedman VA, Andreski P. Feeling better? Trends in general health status. *J Gerontol B Psychol Sci Soc Sci.* 2007;62(1):S11-21.
 39. Department of Health. Health Survey for England 2001: Disability. London: The Stationery Office. 2003.
 40. Jagger C, Matthews R, Matthews F, Robinson T, Robine JM, Brayne C, Medical Research Council Cognitive Function and Ageing Study Investigators. The burden of diseases on disability-free life expectancy in later life. *J Gerontol A Biol Sci Med Sci.* 2007;62(4):408-14.
 41. Addo J, Bhalla A, Crichton S, Rudd AG, McKeivitt C, Wolfe CD. Provision of acute stroke care and associated factors in a multiethnic population: prospective study with the South London Stroke Register. *BMJ.* 2011;342:d744.
 42. McKeivitt C, Coshall C, Tilling K, Wolfe CD. Are there inequalities in the provision of stroke care? Analysis of an inner-city stroke register. *Stroke.* 2005;36(2):315-20.
 43. Bailey L, Vardulaki K, Langham J, Chandramohan D. Introduction to Epidemiology. Maidenhead, England: Open University Press. 2005.