King's Research Portal

DOI:

[10.1136/eb-2016-102461](https://doi.org/10.1136/eb-2016-102461)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Banaschewski, T., Buitelaar, J., Chui, C., Cortese, S., Simonoff, E., & Wong, I. C. K. (Accepted/In press). Methylphenidate for ADHD in children and adolescents: throwing the baby out with the bathwater. *Evidence-Based BMJ*. <https://doi.org/10.1136/eb-2016-102461>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

PERSPECTIVE ARTICLE

Methylphenidate for ADHD in children and adolescents: throwing the baby out with the bathwater

Tobias Banaschewski,¹ Jan Buitelaar,² Celine SL Chui,³ David Coghill,⁴ Samuele Cortese,⁵

Emily Simonoff,^{6*} Ian CK Wong^{3,7} on behalf of the European ADHD Guidelines Group

¹Department of Child Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Mannheim, Germany

²Department of Cognitive Neuroscience, Donders Institute of Brain, Cognition and Behavior, Radboud University Medical Center, Nijmegen, The Netherlands, and Karakter Child and Adolescent Psychiatry University Center, Nijmegen, The Netherlands

³Department of Pharmacology and Pharmacy, University of Hong Kong, Hong Kong, China

⁴Departments of Paediatrics and Psychiatry, Faculty of Medicine, Dentistry and Health Sciences, University of Melbourne, Victoria, Australia

⁵Department of Psychology, University of Southampton, Southampton, UK and Clinical and Experimental Sciences (CNS and Psychiatry), Faculty of Medicine, University of Southampton, UK

⁶Department of Child & Adolescent Psychiatry, King's College London, Institute of Psychiatry Psychology & Neuroscience, London, UK

⁷Research Department of Practice and Policy, UCL School of Pharmacy, London, UK

Word count: 1992

*Correspondence to:

Emily Simonoff

Department of Child and Adolescent Psychiatry

King's College London, Institute of Psychiatry, Psychology and Neuroscience

Box 85, DeCrespigny Park, London SE5 8LF

Phone (00 44) (0) 207 8848 5313

Fax (00 44) (0) 207 508 5600

E-mail: emily.simonoff@kcl.ac.uk

ABSTRACT

A recent Cochrane review assessed the efficacy of methylphenidate for attention-deficit/hyperactivity disorder (ADHD) in children and adolescents. Notwithstanding the moderate-to-large effect sizes for ADHD symptom reduction found in the meta-analysis, the authors concluded that the quality of the evidence is low and therefore the true magnitude of these effects remain uncertain. We identified a number of major concerns with the review, in the domains of study inclusion, approaches to quality assessment, and interpretation of data relating to serious adverse events as well as of the clinical implications of the reported effects. We also found errors in the extraction of data used to estimate the effect size of the primary outcome. Considering all the shortcomings, the conclusion in the Cochrane review that the status of the evidence is uncertain is misplaced. Professionals, parents and patients should refer to previous reviews and existing guidelines, which include methylphenidate as one of the safe and efficacious treatment strategies for ADHD.

Keywords: Attention-deficit/hyperactivity disorder; methylphenidate; randomized controlled trials; meta-analysis; systematic review

Attention-deficit/hyperactivity disorder (ADHD) is a common disorder commencing in childhood and frequently persisting across the lifespan. Current treatment guidelines, including those of the National Institute for Health and Care Excellence (NICE)¹, identify methylphenidate (MPH) as a first-line treatment for ADHD. Furthermore, previous systematic reviews and meta-analyses (e.g.^{2 3}) have reported large effect sizes for the efficacy of MPH in the treatment of ADHD, at least in the short term.

In November 2015, Storebø et al.⁴ published a Cochrane review on the efficacy and tolerability of MPH for the treatment of ADHD in children and adolescents. This review challenged the conclusions of previous reviews and guidelines. Whilst the meta-analysis found similar effect sizes in relation to efficacy to those previously reported, the authors concluded that the magnitude of this effect is uncertain due to the very low quality of the evidence. As such, the conclusions of Storebø et al.⁴ could raise questions about the role of MPH as a core component of ADHD treatment. As an international group committed to the provision of evidence-based clinical guidance on the management of ADHD, we do not agree with the conclusions of Storebø et al.⁴ Rather, we argue that the Storebø et al. review⁴ is flawed in a number of ways that lead to these incorrect conclusions:

1. Inappropriate selection of studies for inclusion;
2. Internal inconsistencies and idiosyncratic procedures in the risk of bias and overall study quality assessment;
3. Misinterpretations of evidence in relation to serious adverse events;
4. A misunderstanding of the meaning of effect sizes, and their clinical implications for individual patients.

Additionally, there are a number of errors in the calculation of standardized mean differences and meta-analytic weights.

Inappropriate study inclusion. As per their protocol, Storebø et al.⁴ aimed to include randomized controlled trials (RCTs) comparing MPH with placebo or no intervention, allowing co-interventions provided that the compared intervention groups received the co-intervention similarly. In fact, Storebø et al.⁴ included three studies, amongst them the large MTA study, where there was no placebo/no treatment arm^{5 6} and/or where MPH was used as an add-on intervention⁵⁻⁷, and a study (included only in secondary analyses) that was not randomized.⁸ Removing these studies increases the effect size for the primary outcome of teacher-reported ADHD symptom ratings from -0.77 (95% CIs -0.90, -0.64) to -0.83 (95% CIs -0.96, - 0.70). The inclusion of the large MTA study⁵ has important implications for all subgroup analyses of long-term (>6 months) versus short-term MPH administration. Storebø et al.⁴ reported a smaller effect for long-term administration according to teacher (but not parent or observer) reports. However, as the MTA is the *only* study meeting their definition of long-term administration these analyses are misleading. We agree with Storebø et al.⁴ that there is an absence of long-term data, but disagree that long-term placebo-controlled RCTs provide the solution. There are serious ethical problems in extending placebo-controlled RCTs into the longer-term where there is evidence of strong benefit. In our view a more appropriate design would be “randomized discontinuation trials” as proposed in the NICE guideline¹. These, along with longer-term safety studies, are now a requirement of the regulatory development programme of the European Medicines Agency for new ADHD medications.

Assessment of study quality. Storebø et al.⁴ adopted the GRADE approach (http://handbook.cochrane.org/chapter_12/12_2_1_the_grade_approach.htm), which includes the Cochrane risk of bias (RoB) tool (http://handbook.cochrane.org/chapter_8/8_assessing_risk_of_bias_in_included_studies.htm). Using GRADE with regards to the main outcome (teacher-rated ADHD symptoms), they

downgraded the quality of evidence by one point for inconsistency of effects (heterogeneity) and by two points for risk of bias. Both these decisions are questionable. In relation to heterogeneity, I^2 for the meta-analysis of the primary outcome was 37%. The Cochrane Handbook suggests that heterogeneity up to 40% “might not be important”. This downgrading is not appropriate, especially considering that exclusion of the MTA study reduces heterogeneity to 25%.

With regard to the RoB assessment, we identified two major issues. First, Storebø et al.⁴ added an additional domain, ‘vested interests’, which is not included in the current Cochrane RoB. The authors support their choice, citing a single work⁹ and the Assessing the Methodological Quality of Systematic Reviews approach (AMSTAR) (<http://amstar.ca/>). While there remains controversy over the best method to account for vested interests,^{10 11} the Cochrane Handbook suggests that it should be reported in the ‘characteristics of included studies’ table rather than in the RoB. Moreover, it is unclear how Storebø and colleagues⁴ handled the assessment of vested interests in specific cases. They defined a trial as being at “high risk of bias” when it was “funded by parties that might have had a conflict of interest (e.g., a manufacturer of methylphenidate) or where potential conflicts of interest were reported by trial authors”. However there appears to be no further specification, for example, the number or role of authors with supposed conflict of interest, the time frame of the putative conflicts or the type of industry support that would lead to a study being rated as biased. This may partially explain why this domain was inconsistently rated across the included trials (see Appendix, Table 1 for examples). In a BMJ online reply to the concerns of others about their use of this domain, the authors stated “There were no trials with only the ‘vested interest bias’ domain assessed as ‘unclear risk of bias’ or ‘high risk of bias’.” This is incorrect; there are actually seven studies in which this was the only RoB domain rated as unclear or high (see Appendix, Table 2).

Storebø et al.⁴ rated studies as being at high risk of bias if any domain (including their additional domain of vested interests) was rated as high *or* unclear. This approach is problematic because domains were rated as *unclear* due to lack of detailed information. However, there is no evidence that the authors routinely attempted to contact authors for clarification regarding RoB domains (including vested interest) rated as ‘unclear’ (see Appendix, page 3), although they did contact some authors for missing quantitative data. For example, in the study by Ashare et al.¹² five RoB domains were rated as “unclear” due to a lack of information. Whilst Storebø et al.⁴ wrote that they contacted the authors for quantitative data, there is no evidence they requested RoB information.

Storebø et al.⁴ argue that the low quality of studies (their assessment) casts doubt on the accuracy of effect sizes. A more scientific approach is to test their opinion with the available data on RoB and study quality. The authors compared RCTs at high versus low risk of *all* bias (their ratings) and found no significant difference in effect size ($\chi^2_1=2.43$, $p=.12$). Unfortunately, this finding is not included in their adjunct publications in the BMJ¹³ and JAMA¹⁴.

Storebø et al.⁴ went on to suggest that even those studies where no item of the RoB was rated as high/unclear were likely to be biased, due to unblinding. They assumed that “people in the trial might know which treatments the children were taking” because of adverse effects associated with MPH. We deem this unlikely, at least with their primary outcome ratings from teachers. The most common adverse effects reported are sleep difficulties and appetite reduction. There is no evidence to support the claim that teachers would be aware of these symptoms in their pupils and this is not our clinical experience.

Serious adverse effects (SAEs) of MPH. The authors used data from nine parallel group trials involving 1,532 participants to explore SAEs. They reported a risk ratio of 0.98 (95% CIs 0.44, 2.22), which did not change significantly with the inclusion of 1,712

participants from both arms of crossover trials (RR=1.50, 95% CIs 0.34, 7.71). They point to the findings from trial sequential analysis to show that the sample size is underpowered to detect SAEs of the frequency reported for MPH versus placebo. It is common knowledge that short-term RCTs are neither intended nor powered to evaluate rare adverse effects and long-term safety. While this is a limitation of the available evidence, a more balanced interpretation of the current evidence is that SAEs *due to* MPH are rare in clinical trials. The vast majority of published observational studies also suggest that SAEs are rare in clinical practice and that the causal associations between these SAE and MPH (rather than to ADHD itself or other associated conditions) remain to be confirmed.

Effect sizes and clinical effectiveness. In a comparison of meta-analyses for common treatments in medicine and psychiatry, Leucht et al.¹⁶ showed that the effect size for MPH compares favourably both with medications for other psychiatric conditions and also with other commonly used interventions in medical conditions. Notable were the moderate effects for corticosteroids in asthma (0.54) and antihypertensives in high blood pressure (0.54, 0.56 for systolic and diastolic blood pressure), with a small effect of the latter for long-term cardiovascular outcome/mortality (0.11). In this context, the efficacy of MPH should be viewed favourably.

A further difficulty with the interpretation of Storebø et al.⁴ is their translation of group-based effect sizes into mean levels of symptom reduction in an attempt to quantify individual improvement. This is an uninformative and inaccurate measure of individual response as it fails to account for baseline symptom severity and the proportion of treated individuals showing clinical benefit, due to inter-individual variability in response. Clinical impact for individuals is better described using the number needed to treat and to harm metrics, which were not presented.

In addition to these major conceptual and methodological issues, we checked Storebø et al.'s⁴ primary analysis of teacher-reported ADHD symptoms. Among the 19 studies reported (three of which we have argued should not have been included), compared to the published data, we found errors in the imputation of data and/or sample size in seven. These errors included a systematic doubling of the sample sizes for *all* crossover trials (by counting each arm as if they were independent) (Appendix, Table 3). Because these errors occurred in both directions, correction of the data had little impact on the aggregate SMD for this outcome. Our limited check raises concerns about the accuracy of the other quantitative findings and the overall quality and conduct of the review.

In summary, we think that the analysis undertaken by Storebø et al.⁴ is flawed and the conclusions misplaced. The additional benefits of MPH in improving both general behaviour (SMD -0.68, 95% CIs -0.78, -0.60) and quality of life (SMD 0.61, 95% CIs 0.48, 0.80) further support the value of MPH for youth with ADHD. We are therefore perplexed why the authors refer to their own findings as “*apparent effects*” (our emphasis). While we agree that some of the trials included in this review are at risk of bias, this aspect of the evidence has not been assessed well. This has led to an overly negative interpretation of the evidence.

It is unfortunate that the authors focus on problems with the quality of the evidence when their own analyses do not support a significant impact of study risk of bias on the effect of MPH. Reviewers are becoming more transparent about the methods to assess study design and data analysis. However, it is still much more difficult to objectively address the introduction of bias through *interpretation* that appears in the discussion section of a paper. In this case, we believe that the interpretation given by Storebø et al.⁴ is not warranted, given their errors and deviations from currently recommended methods.

The beneficial effects of MPH (and other medications for ADHD) need to be placed in the context of other, non-pharmacological, interventions for ADHD. In a systematic review

and meta-analysis of RCTs, we found a relative lack of efficacy on blinded outcomes in reducing ADHD symptoms for many of these.¹⁷⁻²⁰ Where these interventions lead to beneficial reductions in ADHD symptoms, the magnitude of these effects is considerably less than that reported for MPH, and indeed other stimulant and non-stimulant ADHD medications. We therefore believe that there is continued support for the use of MPH as an effective and safe treatment for ADHD.

References

1. National Institute of Health and Clinical Excellence. Attention deficit hyperactivity disorder: Diagnosis and management of ADHD in children, young people and adults. In: Excellence NIfHaC, ed. London, 2008.
2. Faraone SV. Using meta-analysis to compare the efficacy of medications for Attention-Deficit/Hyperactivity Disorder in youths. *Pharmacy & Therapeutics* 2009;34(12):678-94. [published Online First: 2010/02/09]
3. Van der Oord S, Prins PJ, Oosterlaan J, et al. Efficacy of methylphenidate, psychosocial treatments and their combination in school-aged children with ADHD: a meta-analysis. *Clinical Psychology Review* 2008;28(5):783-800.
4. Storebø OJ, Ramstad E, Krogh HB, et al. Methylphenidate for children and adolescents with attention deficit hyperactivity disorder (ADHD). *Cochrane Database of Systematic Reviews* 2015(12) doi: 10.1002/14651858.CD009885
5. The MTA Group. A 14-month randomized clinical trial of treatment strategies for attention-deficit hyperactivity disorder. *Archives of General Psychiatry* 1999;56:1073-86.
6. Firestone P, Kelly MJ, Goodman JT, et al. Differential effects of parent training and stimulant medication with hyperactives: A progress report. *J Am Acad Child Psychiatry* 1981;20(1):135-47. [published Online First: 1981/01/01]
7. Brown RT, Wynne ME, Medenis R. Methylphenidate and cognitive therapy: a comparison of treatment approaches with hyperactive boys. *Journal of Abnormal Child Psychology* 1985;13(1):69-87. [published Online First: 1985/03/01]

8. Lufi D, Parish-Plass J, Gai E. The effect of methylphenidate on the cognitive and personality functioning of ADHD children. *Israel Journal of Psychiatry and Related Science* 1997;34(3):200-09.
9. Lundh A, Sismondo S, Lexchin J, et al. Industry sponsorship and research outcome (Review). *The Cochrane Collaboration* 2012;12:1=88.
10. Sterne JA. Why the Cochrane risk of bias tool should not include funding source as a standard item *Cochrane Database of Systematic Reviews* 2013;12:ED000076.
11. Bero LA. Why the Cochrane risk of bias tool should include funding source as a standard item. *Cochrane Database of Systematic Reviews* 2013;12:ED000075.
12. Ashare R, L., Hawk L, W.,, Shiels K, et al. Methylphenidate enhances prepulse inhibition during processing of task-relevant stimuli in attention-deficit/hyperactivity disorder. *Psychophysiology* 2010;47(5):838-45.
13. Storebø OJ, Krogh HB, Ramstad E, et al. Methylphenidate for attention-deficit/hyperactivity disorder in children and adolescents: Cochrane systematic review with meta-analyses and trial sequential analyses of randomised clinical trials. *British Medical Journal* 2015;351 doi: 10.1136/bmj.h5203
14. Storebø OJ, Simonsen E, Gluud C. Methylphenidate for attention-deficit/hyperactivity disorder in children and adolescents. *The Journal of the American Medical Association* 2016;315(18):2009-10. doi: 10.1001/jama.2016.3611
15. Cortese S, Holtmann M, Banaschewski T, et al. Practitioner review: Current best practice in the management of adverse events during treatment with ADHD medications in children and adolescents. *Journal of Child Psychiatry and Psychology* 2013;54(3):227-46. . doi: doi: 10.1111/jcpp.12036. [published Online First: Epub 2013 Jan 7]

16. Leucht S, Hierl S, Kissling W, et al. Putting the efficacy of psychiatric and general medicine medication into perspective: review of meta-analyses. *British Journal of Psychiatry* 2012;200(2):97-106. doi: 10.1192/bjp.bp.111.096594 [published Online First: 2012/02/03]
17. Sonuga-Barke E, Brandeis D, Cortese S, et al. Non-pharmacological interventions for attention-deficit/hyperactivity disorder: systematic review and meta-analyses of randomised controlled trials of dietary and psychological treatments. *The American Journal of Psychiatry* 2013;170(3):275-89. [published Online First: January 2013]
18. Daley D, van der Oord S, Ferrin M, et al. Behavioral Interventions in Attention-Deficit/Hyperactivity Disorder: A Meta-Analysis of Randomized Controlled Trials Across Multiple Outcome Domains. *Journal of the American Academy of Child and Adolescent Psychiatry* 2014;53(8):835-47 e5. doi: 10.1016/j.jaac.2014.05.013 [published Online First: 2014/07/27]
19. Cortese S, Ferrin M, Brandeis D, et al. Cognitive training for attention-deficit/hyperactivity disorder: meta-analysis of clinical and neuropsychological outcomes from randomized controlled trials. *Journal of the American Academy of Child and Adolescent Psychiatry* 2015;54(3):164-74. doi: 10.1016/j.jaac.2014.12.010 [published Online First: 2015/02/28]
20. Cortese S, Ferrin M, Brandeis D, et al. Neurofeedback for attention-deficit/hyperactivity disorder: meta-analysis of clinical and neuropsychological outcomes from randomized controlled trials. *Journal of the American Academy of Child and Adolescent Psychiatry* 2016;55(6):444-55.

Acknowledgements

This article is written on behalf of the European ADHD Guidelines Group, whose members are: Philip Asherson, Tobias Banaschewski, Daniel Brandeis, Jan Buitelaar , David Coghill, Samuele Cortese , David Daley, Marina Danckaerts , Ralf Dittmann , Manfred Doepfner , Maite Ferrin, Chris Hollis, Martin Holtmann, Eric Konofal, Michel Lecendreux , Aribert Rothenberger, Paramala Santosh, Emily Simonoff, Edmund Sonuga-Barke, Cesar Soutullo, Hans-Christoph Steinhausen, Argyris Stringaris, Eric Taylor, Saskia Van der Oord , Ian CK Wong, Alessandro Zuddas

Contribution authorship statement

All authors contributed to

- the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work;
- Drafting the work or revising it critically for important intellectual content;
- Final approval of the version to be published;
- Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.