



King's Research Portal

DOI:

[10.1111/all.13718](https://doi.org/10.1111/all.13718)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Untersmayr, E., Bax, H. J., Bergmann, C., Bianchini, R., Cozen, W., Gould, H. J., Hartmann, K., Josephs, D. H., Levi-Schaffer, F., Penichet, M. L., O'Mahony, L., Poli, A., Redegeld, F. A., Roth-Walter, F., Turner, M. C., Vangelista, L., Karagiannis, S. N., & Jensen-Jarolim, E. (2019). AllergoOncology: Microbiota in allergy and cancer: a European Academy for Allergy and Clinical Immunology position paper. *Allergy*, 74(6), 1037-1051. <https://doi.org/10.1111/all.13718>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

DR. EVA UNTERSMAJR (Orcid ID : 0000-0002-1963-499X)

DR. RODOLFO BIANCHINI (Orcid ID : 0000-0003-0351-6937)

PROF. LIAM O'MAHONY (Orcid ID : 0000-0003-4705-3583)

DR. SOPHIA N KARAGIANNIS (Orcid ID : 0000-0002-4100-7810)

PROF. ERIKA JENSEN-JAROLIM (Orcid ID : 0000-0003-4019-5765)

Article type : Position Paper

AllergoOncology: microbiota in allergy and cancer - an EAACI position paper

Untersmayr E¹ (<https://orcid.org/0000-0002-1963-499X>), Bax HJ^{2,3} (<http://orcid.org/0000-0003-0432-4160>), Bergmann C⁴ (<https://orcid.org/0000-0002-2076-8813>), Bianchini R⁵ (<https://orcid.org/0000-0003-0351-6937>); Cozen W^{6,7,8} (NO ORCID), Gould HJ^{9,10} (<https://orchid.org/0000-0003-0411-688>), Hartmann K¹¹, Josephs DH^{2,3} (<https://orcid.org/0000-0002-5610-9123>) Levi-Schaffer F¹² (<https://orcid.org/0000-0003-0620-2810>), Penichet ML^{13,14,15,16,17} (No ORCID), O'Mahony L¹⁸ (<https://orcid.org/0000-0003-4705-3583>), Poli A¹⁹ (No ORCID), Redegeld F²⁰ (<https://orcid.org/0000-0001-8830-7960>), Roth-Walter F⁵ (<https://orcid.org/0000-0001-5005-9228>), Turner MC^{21,22,23,24} (<https://orcid.org/0000-0002-6431-1997>), Vangelista L²⁵, Karagiannis SN^{*2} (<https://orcid.org/0000-0002-4100-7810>) and Jensen-Jarolim E^{*1,5} (<https://orcid.org/0000-0003-4019-5765>) (*shared last and co-corresponding)

¹ Institute of Pathophysiology and Allergy Research, Center of Pathophysiology, Infectiology and Immunology, Medical University Vienna, Vienna, Austria.

² St. John's Institute of Dermatology, School of Basic & Medical Biosciences, King's College London, Guy's Hospital, London, United Kingdom.

³ School of Cancer and Pharmaceutical Sciences, King's College London, Guy's Hospital, London, United Kingdom.

⁴ ENT Research Institute for Clinical Studies, Essen, Germany.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/all.13718

This article is protected by copyright. All rights reserved.

- ⁵ Department of Comparative Medicine, The Interuniversity Messerli Research Institute, University of Veterinary Medicine Vienna, Medical University Vienna, University Vienna, Austria.
- ⁶ Department of Preventive Medicine, Keck School of Medicine of University of Southern California, Los Angeles, CA, United States.
- ⁷ Department of Pathology, Keck School of Medicine of University of Southern California, Los Angeles, CA, United States.
- ⁸ Norris Comprehensive Cancer Center, Keck School of Medicine of Los Angeles, CA, United States.
- ⁹ Randall Centre for Cell and Molecular Biophysics, School of Basic & Medical Biosciences, King's College London, New Hunt's House, London, SE1 1UL, United Kingdom.
- ¹⁰ Medical Research Council & Asthma UK Centre in Allergic Mechanisms of Asthma, London, United Kingdom.
- ¹¹ Department of Dermatology, University of Luebeck, Luebeck, Germany.
- ¹² Pharmacology and Experimental Therapeutics Unit, The Institute for Drug Research, School of Pharmacy, Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel.
- ¹³ Division of Surgical Oncology, Department of Surgery, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, United States.
- ¹⁴ Department of Microbiology, Immunology and Molecular Genetics, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, United States.
- ¹⁵ Jonsson Comprehensive Cancer Center, University of California, Los Angeles, CA, United States.
- ¹⁶ The Molecular Biology Institute, University of California, Los Angeles, CA, United States.
- ¹⁷ UCLA AIDS Institute, Los Angeles, CA, United States.
- ¹⁸ Departments of Medicine and Microbiology, APC Microbiome Ireland, National University of Ireland, Cork, Ireland.
- ¹⁹ Luxembourg Institute of Health, Department of Infection and Immunity, Esch-sur-Alzette, Luxembourg.
- ²⁰ Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University, Utrecht, The Netherlands.
- ²¹ Barcelona Institute for Global Health (ISGlobal), Barcelona, Spain.
- ²² Universitat Pompeu Fabra (UPF), Barcelona, Spain.
- ²³ CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain.
- ²⁴ McLaughlin Centre for Population Health Risk Assessment, University of Ottawa, Ottawa, Canada.

²⁵ Department of Biomedical Sciences, Nazarbayev University School of Medicine, Astana, Kazakhstan.

Acknowledgements:

The AllergoOncology Task Force was financed by the European Academy for Allergy and Clinical Immunology (EAACI). The authors would like to thank EAACI for their financial support in the development of this Task Force position paper.

The authors acknowledge support by the Austrian Science Fund FWF grants P23398-B11, F4606-B28 (JJE), KLI284 (UE); by the Israel Cancer Association #20161131, the Israel Science Foundation #213/05 the Aimwell Charitable Trust and the Emalie Gutterman Memorial Endowed Fund (LSF); by the National Institutes of Health (NIH) grants R01CA196266, R01CA196266-01A1:S1 and R01 CA181115 (PM); by NIH/NCI grant R03CA173842 and American Society of Hematology Bridge Grant (CW); German Research Council (DFG; HA 2393/6-1 and Excellence Cluster „Inflammation at Interfaces“) (HK); by IGEM Therapeutics Ltd (BHJ; KSN); by Cancer Research UK (C30122/A11527; C30122/A15774) (KSN, BHJ); The Academy of Medical Sciences (JDH, KSN); the Medical Research Council (MR/L023091/1); Breast Cancer Now - working in partnership with Walk the Walk (KSN); CR UK/NIHR in England/DoH for Scotland, Wales and Northern Ireland Experimental Cancer Medicine Centre (C10355/A15587) (KSN); The research was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London (IS-BRC-1215-20006) (KSN). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the UK Department of Health; TMC was supported by the Departament de Salut, Generalitat de Catalunya (SLT002/16/00232). ISGlobal is a member of the CERCA Programme, Generalitat de Catalunya. The authors thank BCAM van Esch for her contribution to the section “Non-digestible Oligosaccharides” and to Table 4.

Co-Correspondence:

1) Erika Jensen-Jarolim, MD

Institute of Pathophysiology and Allergy Research, Center of Pathophysiology, Infectiology and Immunology, Medical University Vienna, Spitalgasse 23, 1090 Vienna, Austria
erika.jensen-jarolim@meduniwien.ac.at

2) Sophia Karagiannis, PhD

St. John's Institute of Dermatology, School of Basic & Medical Biosciences, King's College London, 9th Floor, Guy's Tower, Guy's Hospital, London SE1 9RT, United Kingdom
sophia.karagiannis@kcl.ac.uk

Contributions (according to authorship series)

UE drafted the manuscript, contributed “Cells of the epithelial barrier”, “Oncoimmunology and allergy” and Table 1, compiled all contributions, and contributed to the Abstract and Introduction; BHJ and KSN contributed “B cells and Bregs”, and with GH “antibodies”; BC “Redox regulation in allergy and cancer”; BR “DC”, “Macrophages” and “ILCs”; CW and PM “Microbial translocation”; RF composed “Non-digestible Oligosaccharides” and Table 4; HK and LSF “Eosinophils”; RF with HK “mast cells”; OML contributed chapters “Microbiota metabolites” and “Tregs”; RWF wrote “lipocalins”, “Dietary micronutrients” and Tables 2 and 3; JDH “Hygiene hypothesis in cancer” and Suppl. Table 1; VL “Microbiota engineering” and designed Figure 2, JJE and TMC wrote “Hygiene hypothesis in allergy” with Suppl. Table 1. PA designed Figure 1; PA, BHJ and JDH completed the graphical abstract based on the paper contents; KSN and JJE orchestrated the whole process, wrote the Conclusion and Abstract, and conducted the final editing.

Abstract

The microbiota can play important roles in the development of human immunity and the establishment of immune homeostasis. Lifestyle factors including diet, hygiene, and exposure to viruses or bacteria, and medical interventions with antibiotics or anti-ulcer medications, regulate phylogenetic variability and the quality of cross-talk between innate and adaptive immune cells via mucosal and skin epithelia. More recently, microbiota and their composition have been linked to protective effects for health. Imbalance, however, has been linked to immune-related diseases such as allergy and cancer, characterized by impaired, or exaggerated immune tolerance, respectively. In this AllergoOncology position paper, we focus on the increasing evidence defining the microbiota composition as a key determinant of immunity and immune tolerance, linked to the risk for development of allergic and malignant diseases. We discuss novel insights into the role of microbiota in disease and patient responses to treatments in cancer and in allergy. These may highlight opportunities to improve patient outcomes with medical interventions supported through a restored microbiome.

Keywords:

Allergy; AllergoOncology; Breg; Treg; cancer; checkpoint; hygiene hypothesis; diet, lipocalin; microbiome; microbiota; immune tolerance;

1. Introduction

Why studying microbiota is important for the field of AllergoOncology

The collective genome of *all* microorganisms living in and on the surfaces of the human body is defined as the *microbiome* and contains 150 times more genes than the 23,000 protein-coding genes of human origin (**Table 1**). The human microbiome project (1) has contributed to understanding of the composition, function and diversity of the human *microbiota*, i.e. all microorganisms populating the inner and outer surfaces of the human body, including viruses, fungi, protozoa, archaea, and bacteria (**Table 2**).

The human microbiota show remarkable variability and have a mutualistic relationship with the human host. The microbiome profoundly affects the epithelium and the mucosal immune system and *vice versa*. There is growing evidence that microbiota play a paramount role in the control of immune mediated diseases such as allergy and cancer, the two complementary diseases in the frame of AllergoOncology (2).

Hygiene hypothesis in allergy: From epidemiology to mechanisms

The allergy epidemics correlate with improved hygiene practices associated with urban lifestyles (**Suppl. Table 1**). Milestone studies revealed that this epidemic can be counter-regulated in part by exposure to a traditional farming environment. The cornerstones of the allergy protective farm effect are: i) farm activities during pregnancy, ii) early life farm exposure, and iii) raw milk consumption, but farms are also protective against viral infections. Protection depends on the farm dust levels in an area, and on the type of farming.

To give an example, children from Amish traditional pre-industrial farming communities were less prone to allergy development than Hutterites who use modern industrial farming methods. Amish farm dust initiated innate immune pathways associated with protection from allergy development in a mouse model. Additionally, rich microbial exposure by cohabitation with wild mice improves the immune response in laboratory mice. In line with this, dog keeping is reported to protect children from asthma. Also helminth infections can support the establishment of protective microbiota (for summary and references see **Suppl. Table 1**).

Molecular mechanisms are increasingly understood. Viral antigens due to molecular mimicry counteract specific allergen sensitization and induce specific T effector memory responses. Exposure to N-Glycolylneuraminic acid (Neu5Gc), a sialic acid compound in farm dust, induces regulatory pathways. Endotoxin exposure of bronchial epithelial cells counteracts allergen-induced Th2 responses. Prenatal farm exposure supports Treg and Th17 cell differentiation. The protective immunity associated with the consumption of raw milk induces FOXP3 demethylation and T regulatory cells. Exposure of bronchial epithelia to farm dust, for instance containing CpG-DNA, or of gut epithelia to farm milk enhances epithelial barrier integrity, resulting in protective innate immunity to allergens and viruses (**Suppl. Table 1**).

In accordance, immature gut microbial composition at age 1-year was positively associated with asthma risk at 5-years in children with asthmatic mothers (3). There were inverse associations with relative abundances of genera *Faecalibacterium*, *Bifidobacterium*, *Roseburia*, *Alistipes*, *Lachnospiraceae incertae sedis*, *Ruminococcus*, and *Dialister*, and a positive association with *Veillonella*. Inverse associations of relative abundances of genera *Lachnospira*, *Veillonella*, *Faecalibacterium*, and *Rothia* at age 3-months and atopy and wheeze at age 1-year were observed in another study (4), as was amelioration of lung inflammation in adult offspring of germ-free mice inoculated with these taxa. The group at highest risk for multisensitized atopy (2-years) and asthma (4-years) was characterized by lower abundance of gut bacteria *Bifidobacterium*, *Akkermansia*, *Faecalibacterium*, and *Lactobacillus* and higher abundance of fungi *Candida* and *Rhodotorula* at 1-11 months (5). Decreased gut microbiome diversity was correlated with CD4+ T cell decline in immune-

deficient patients (6). Increasing use of antibiotics has been linked with dysbiosis and enhanced prevalence of allergies and asthma (7). Also, the pharmacologic impairment of gastric digestion is associated with gut dysbiosis and has been correlated with allergy in mice and humans (8, 9). However, study results of probiotic supplementation in childhood asthma or wheeze are inconclusive (10).

Hygiene hypothesis in oncology: From epidemiology to mechanisms

Increased incidence of certain cancers in Westernized countries (11) may be associated with under-exposure to certain microbial species, modern lifestyle, and consumption of sterilized food (12). Higher socioeconomic status is associated with increased incidence of Hodgkin lymphoma (13), while daycare attendance and higher number of childhood infections are linked with lower risk of acute lymphoblastic leukemia (14) and adult chronic lymphoid leukemia (15). Increased cancer risk is observed in patients with autoimmune diseases and chronic allergic disorders (16). Mice with rapid melanoma growth and poor immunosurveillance, exhibited relatively low levels of *Bifidobacterium* species which when restored by oral feeding and co-housing, resulted in enhanced tumour antigen presentation and reduced malignant growth (17).

Epidemiological studies have also examined specific bacteria, viruses, periodontal disease, and circulating antibodies to selected pathogens in relation to cancer risk (18). Carriage of *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans* in pre-diagnostic oral wash samples was positively associated with pancreatic cancer risk and *Fusobacteria* and its genus *Leptotrichia* were inversely associated (19). *Tannerella forsythia* was positively associated with esophageal adenocarcinoma and *Porphyromonas gingivalis* with squamous cell carcinoma (20). Greater abundance of genera *Corynebacterium* and *Kingella* were inversely associated with head and neck squamous cell cancer (21).

An omnivore, but not a vegetarian, diet promotes butyrate-producing *Lachnospiraceae* (Firmicutes/Clostridiales). *Bifidobacterium spp* (Actinobacteria/ Bifidobacteriales/ Bifidobacteriaceae) are strongly stimulated by dietary intake of oligosaccharides as present

in milk, while *Erysipelotrichi* (Firmicutes/Erysipelotrichales/Erysiopelotrichaceae) are stimulated by high fat, Western diets (22). A prudent diet (rich in whole grains and fiber) was associated with a lower risk of *Fusobacterium nucleatum*-positive colorectal cancer incidence (23). African Americans fed a high-fibre, low-fat diet and rural Africans a high-fat, low-fibre diet, demonstrated reciprocal changes in colonic mucosal inflammation and biomarkers of cancer risk and in colonic microbiota and metabolome including in saccharolytic fermentation and butyrogenesis, and secondary bile acid synthesis (24).

It has also been postulated that the use of antibiotics may support cancer progression through subversion of immunosurveillance. In a transgenic mouse model of spontaneous mammary carcinoma development, treatment with metronidazole and ciprofloxacin resulted in accelerated development of these tumours (25). Repeated antibiotic use may also increase risk of certain human malignancies (26).

2. Metabolites critically shape the microbiome

Dietary micronutrients

Micronutrients being essential food for microbes substantially influence microbiota composition. Most bacteria require iron and have evolved multiple strategies for sequestration including the production of hemophores and low molecular iron chelators named siderophores. Iron supplementation promotes the establishment of bacteria relying on this metal (**Table 3**). In atopy a poor iron status is associated with allergy, whereas an improved status abrogates or seems to prevent the onset of allergy. In cancer, iron contributes to tumor growth with high serum iron increasing the risk of several cancers (**Table 4**) (27).

Vitamin A and D (**Tables 3 and 4**) are important for mucosal immunity (28). Retinol supplementation does not change the risk of allergy (29) or cancer (30), though increased serum retinol levels are associated with prostate cancer (31). Vitamin D supplementation

slightly decreases respiratory infections and the incidence of asthma in off-spring (32), prevents hematologic malignancies in elderly women and advanced colorectal adenomas (33), but seem not to impact allergy risk (34). Vitamin D supplementation decreases the abundance of *Bifidobacterium* and *Coprococcus* (35).

Serum folate levels are associated with serum IgE (36), high plasma folate is associated with decreased risk of wheezing (37), and folic acid supplementation has been linked with an increased risk for lung cancer (38). Folate can be produced by most Bacteroidetes, Fusobacteria and Proteobacteria, but rarely from Actinobacteria and Firmicutes (39). In summary, dietary micronutrients shape the microbiome which contributes products essential for an effective immune defence.

Non-digestible Oligosaccharides

Non-digestible Oligosaccharides (NDO) are potential substrates for bacterial metabolism in the colon and a declining microbial diversity in the gut is implicated in the rising incidence of allergic disorders in early life (40). Manipulation of the gut microbiota with NDO from natural sources or supplemented galacto-oligosaccharides (GOS), lactose and fructo-oligosaccharides (FOS) holds great promise for the treatment of inflammatory and allergic diseases (**Table 4**). NDO feeding reduced the development of atopic dermatitis (AD) in infants at risk of allergy which was associated with increased *Bifidobacterium* and *Lactobacilli* (41, 42), re-balancing the immune response from a predominant Th2-type allergic profile at birth to a more Th1-type and regulatory T cell profile. NDO protect against allergic manifestations in experimental models of food allergy and allergic asthma (43, 44), and reshape the gut microbiota with increased levels of short chain fatty acids (SCFAs) (45). Less is known concerning the anti-cancer effects of NDO (46). Experimental models show encouraging effects of GOS, FOS, agaro-oligosaccharides (47) and butyrate (46) in influencing microbiota diversity (e.g. increased *Fusobacterium*) and the prevention and progression of colorectal cancer (48). NDO showed potential in the protection against the development of cancer in healthy subjects (49). Since Western diets are typically low on

NDO, increasing the intake of NDO may benefit human health via an improved microbiome and/or via direct effects on the immune system.

Microbiota metabolites

Dairy products such as yogurt and butter contain short chain fatty acids (SCFAs), while SCFAs are also produced by intestinal bacteria following their fermentation of fibers present in vegetables and fruits.(50) The major SCFAs are acetate, propionate and butyrate. SCFAs are an important energy source for intestinal epithelial cells and regulate the assembly and organization of epithelial tight junctions. Abnormalities in the production of these metabolites (due to dietary factors and/or microbial dysbiosis) might play a role in the pathogenesis of type-2 diabetes, obesity, inflammatory bowel disease, colorectal cancer and allergies (51, 52). Butyrate influences the activity of histone deacetylases (HDAC), responsible for decreasing dendritic cell IL-12 and IL-6 secretion, and allows dendritic cells to promote Tregs. Propionate can also contribute to the induction of T cell Foxp3 expression by dendritic cells, while acetate does not have this activity, possibly due to the lack of HDAC activity (53). G-Protein coupled receptors GPR109a and GPR43 have been described to contribute to these effects, while GPR43 expression on colonic inducible Treg cells is associated with their expansion and IL-10 secretion (54, 55).

3. Microbiota regulating cellular players in Innate and adaptive responses

Cells of the epithelial barrier

Epithelial cells have a very close interaction with compartment-specific microbiota (**Figure 1**). These cells are able to sense conserved microbial associated molecular patterns by innate pattern recognition receptors (PRR) such as TLRs and Nod-like receptors. These interactions can control epithelial cell proliferation and barrier function (56). Not only direct

Accepted Article
interaction, but also microbiota derived metabolites such as SCFAs and bacterial quorum sensing (QS) can have an impact on epithelial cells. In the intestine, SCFAs stimulate the inflammasome pathway via binding to GPRs on the surface of epithelial cells. Associated with enhanced epithelial IL-18 secretion, barrier integrity is enhanced and epithelial cells secrete more antimicrobial peptides (57, 58). Both, SCFA and QS signals have immunomodulatory potential by stimulating survival pathways and contributing to intestinal homeostasis (59-61). Thus, microbiota and metabolites have a paramount influence on epithelial cell function.

Dendritic cells

As professional antigen presenting cells (APC), dendritic cells (DC) are in contact with both invading pathogens and commensal microbiota and maintain the balance between inflammatory and tolerogenic immune responses. Human monocyte-derived dendritic cells (moDCs), when matured in the presence of the SCFA butyrate, increase IL-10, but decrease IL-6 and IL-12 expression (62). Similarly, treatment of mice with the SCFA propionate generate new bone marrow DC precursors with high phagocytic capacity but an impaired ability to promote T helper type 2 (Th2) responses in the lung. This effect is dependent on G protein-coupled receptor 41 (GPR41, also called free fatty acid receptor 3 or FFAR3), but not GPR43 (also called free fatty acid receptor 2 or FFAR2) (55). DCs can also exert roles in adoptive T cell therapy (ACT) in cancer, and the composition of the gut microbiome or treatment with antibiotics could lead to an increase in CD8 α^+ DCs, and consequent IL-12 release that sustains the anti-tumor ACT (63).

Macrophages

The microbiota and its metabolites such as SCFAs strongly influence myelopoiesis and the tissue-resident macrophages. In the intestine, microbial SCFA butyrate activates its endogenous receptor GPR109a promoting anti-inflammatory properties in colonic macrophages that induce differentiation of Treg cells and IL-10-producing T cells. Such

responses may be beneficial in food tolerance, but perhaps not in cancer. Moreover, butyrate exerts an anti-inflammatory effect on lamina propria macrophages by inhibition of IL-6, IL-12 and NO via inhibition of HDACs (64). Noteworthy, this mechanism did not affect primary LPS response genes such as *Tnf α* and *Ccl2*. Microbiota can also affect macrophage phenotype and response if they are not in contact with the innate immune system. In the CNS of germ-free mice, macrophages (microglia) display an altered phenotype. In the lungs of mice treated with antibiotics, macrophages are polarized towards a M2 pro-allergic phenotype by prostaglandin E2 (PGE₂) enhanced by commensal fungi (65).

Mast cells

Mast cells (MC) are crucial effector cells in allergy and other inflammatory diseases (66). They also play important roles in regulating the microenvironment of various tumors (66). There is increasing evidence that MC function can be modulated by commensal, symbiotic and pathogenic microorganisms (67). Microorganisms may influence MC activation via direct interaction or via secreted metabolites. For example, co-culture with *Lactobacillus rhamnosus* downregulates gene expression for the high affinity IgE receptor and histamine H4 receptor in human MC, while increasing IL-8, IL-10, CCL2 and TNF α (67). Stabilization of MC by *L. rhamnosus* is possibly induced indirectly via targeting the KCa3.1 channel (68). Some *Lactobacillus* strains, such as *L. paracasei*, inhibited IgE-mediated activation of murine MC with a possible involvement of TLR2. Inhibition by *L. casei* is cell contact-dependent, but TLR- or Nod1/2-independent (67). Inhibition of MC activation by microbiota-derived metabolites such as SCFA may be mediated via the surface receptors GPR41, GPR43 and GPR109A (69, 70) nuclear receptor family peroxisome proliferator-activated receptors (PPAR) or direct inhibition of histone deacetylase activity (71). The bioactive milk peptide glycomacropeptide (GMP) increased intestinal *Lactobacillus* and *Bifidobacterium* levels in rats and decreased allergen-induced MC activation (72).

Eosinophils

Eosinophils are granulocytic leukocytes that exert important functions in protective immune responses against helminths and other pathogens (73). They also serve as key effector cells in allergies and other inflammatory diseases. Recent studies demonstrated that eosinophils crucially participate in maintaining the specific tissue resident microbiome (74). Conversely, functions of eosinophils are regulated by pathogenic as well as probiotic microorganisms; e.g. human eosinophils were found to ingest *Clostridium difficile*, which subsequently stimulated the release of eosinophil-derived neurotoxin (75). In contrast, ingestion of the probiotic strain *Bifidobacterium bifidum* resulted only in a minor neurotoxin release (75). Comparably, mouse eosinophils were found to take up the probiotic *Lactobacillus reuteri* (76). Using various mouse models of asthma and AD, probiotics like *Lactobacillus fermentum* and *Lactobacillus rhamnosus* were found to improve allergic inflammation associated with decreased eosinophil infiltration, although a direct effect of probiotic bacterial strains on eosinophils was not demonstrated in these studies (77, 78).

ILCs

Innate lymphoid cells (ILCs) develop normally in the absence of the microbiota, however, signals from commensal microorganisms influence the maturation and acquisition of the tissue-specific functions of ILCs. ILC3 cells maintain tolerance to commensal-specific T cell responses and their depletion and subsequent abrogation of IL-22 production, results in loss of intestinal bacteria (65). Moreover, ILC3s release GM-CSF and induce tolerance when intestinal macrophages release IL-1 β in response to microbial sensing. A recent study demonstrates that TNF- β production by ILC3s is crucial for the production of IgA and for microbiota homeostasis in the intestine. Another ILC subtype, ILC2s, are activated by IL-25 produced in a microbiota-dependent manner by epithelial tuft cells. Deletion of the ILC1-lineage transcription factor T-bet in the innate immune system results in ILC-dependent and *Helicobacter typhlonius*-driven inflammation of the intestine, connected to cancer initiation

and promotion (65). Taken together, the microbiota and their metabolites shape the spectrum of different immunoregulatory ILCs and modulate their ability to prevent tumour immune evasion or allergic responses.

Tregs

Tregs are potent immune regulating cells that play a central role in controlling immune responses. Tregs can support the reduction of allergic diseases, and, on the other hand, the progression of cancer (79). Multiple mechanisms are employed by Treg cells and include production of inhibitory cytokines (IL-10, TGF- β and IL-35), cytotoxicity of effector T cells and APCs (via granzymes A and B), direct inhibition of DCs (e.g. via PD-1 and CTLA4) and metabolic disruption of effector cells (CD25, cAMP, adenosine, CD39, and CD73) (80). Germ-free mice do not fully develop Tregs, similarly to mice treated with antibiotics or mice lacking Toll-like receptors (TLRs). However, not all bacterial strains are equally effective in inducing Tregs. *Bifidobacterium longum* 35624, Clostridia and *Bacterioides fragilis* have been shown to induce intestinal Treg cells, while other bacterial strains do not induce Tregs (81, 82). Pattern-recognition receptor activation on DCs seems to be an important mechanism by which intestinal microbes may promote Treg cell differentiation (83).

B cells and B regs

B cells are known to promote allergy through antigen presentation and class-switching to IgE, whereas under certain conditions, tumour-infiltrating B cells can be associated with improved patient survival in cancer (79, 84). In contrast, Breg-associated immune tolerance can lead to control of allergy or tumour progression in cancer (79).

The microbiome may have a role in mediating these multifaceted and opposing B cell effects in allergy and cancer. Microbiota regulate activation and differentiation of B cells (85). Gut microbiota antigens directly trigger B cell activation by binding BCRs, and microbial products activate TLR-expressing B cells, increasing B cell survival, antigen presentation and antibody production (86). Microbiota also mediate the release of epithelial cell and

eosinophil-derived cytokines/chemokines that activate and recruit B cells, and may promote T follicular helper cell-mediated differentiation of plasma cells (85). Conversely, the gut microbiome reportedly induces DCs to produce cytokines, such as IL-1 β and IL-6, which promote B cell differentiation to Bregs (87).

These findings point to a functional cross-talk between humoral immunity and the microbiome.

4. Molecular crosstalk of microbiota with innate and specific immune defence

Redox regulation in allergy and cancer

Recent reports demonstrated that certain microbes can stimulate intracellular signaling, involving PTEN, MAPK and PTP via the generation of reactive oxygen species (ROS) in epithelia, when harboring regulatory redox-sensitive thiolates (88). Commensal bacteria alter the epithelial redox environment by production of oxygen radicals, cause epithelial cell DNA damage and may harbor carcinogenic properties, for example in colon carcinoma development (89). Furthermore, allergic asthma is associated with an increase in endogenous ROS formation, leading to oxidative stress-induced damage to the respiratory system and weakened antioxidant defences. These may result in abnormal physiologic function of DNA, proteins, and lipids that can augment bronchial hyperresponsiveness and inflammation (90).

Overall, the microbiome is implicated in redox regulating pathways, which are relevant for both chronic inflammation and cancer.

Lipocalins

Human lipocalins, such as tear lipocalin 1 and lipocalin 2 present in the nasal mucosa, can interfere with bacterial iron sequestration (27), and may directly modulate site-specific microbial composition. The immunomodulatory and apoptosis-regulating properties of LCN2

Accepted Article

have been linked to its ability to shuttle iron (91). In line with the hygiene hypothesis, the limited “microbial exposure” of allergic individuals may also result in deficiencies of the immune regulatory machinery and this can lead to hyper-reactive responses (92). Interestingly, nearly all major mammalian allergens belong to the lipocalin family (93), and are similarly able to bind to bacterial and plant-derived iron chelators, pointing to a potential role for interference of LCN2 function. In several cancer types and in line with a perturbed iron regulation, LCN2 plays an important role in oncogenesis and cancer progression and may serve as a disease biomarker (94).

Antibodies

Microbiota associated antigens can induce IgA-producing plasma cells in the gut; CD40-ligand and IL-21 from T follicular helper cells, APRIL and BAFF from DCs, induce Activation-Induced Cytidine Deaminase (AID) expression by B cells, promoting IgA class switch recombination. As the composition of the microbiota changes, so do the IgA repertoires (95). In a recent mouse study, IL-33 deficient mice, with significantly-lower levels of intestinal IgA and colon-residing IgA+ B cells, had increased DNA damage-induced tumours; observations that were ameliorated upon microbiome restoration by co-housing deficient animals with wildtype mice (96). In humans, altered gut microbiota diversity and low total IgA levels may be associated with development of allergies and asthma (97), and IgA-deficient individuals have a moderately increased risk of cancer, with higher risks in gastrointestinal cancer (98). This disease risk may be a result of impaired mucosal barrier function. However, the associations between microbiota-driven antibody production with allergies and cancer are not yet sufficiently understood.

5. Translational implications of microbiota

Oncoimmunology and allergy

Specific microbes and the microbiota in general can be considered as important drivers of immunomodulation and can contribute to establishing immune tolerance, with differential impact on the many diverse immune-mediated diseases. Due to the close interaction and the bidirectional influence of intestinal microbiota on the mucosal immune system, the gut can be considered as an essential site of immune cross-talk in the human host. When it comes to allergy or cancer, however, regulation of tolerance has an opposing impact on disease development and treatment (79).

In cancer the impact of microbiota was for a long time primarily considered in the context of dysbiosis, increased epithelial translocation and carcinogenic effector mechanisms (99).

However, in recent years, emerging knowledge points to the role of microbiota, especially in the gut, in anti-cancer immune mechanisms (100). This is due to studies describing a reduced anti-cancer efficacy of chemotherapeutics such as cyclophosphamide and platinum salts in germ-free and antibiotic-treated mice. Moreover, the redox equilibrium of myeloid cells contained in the tumor microenvironment is influenced by intestinal microbiota and due to the importance of the gut in immune fitness, intestinal microbes are essential for the availability of immunomodulators.

In cancer immunotherapy, major advances have been made in the past years to understand the contribution of microbiota composition to successful treatment. Studies are now starting to evaluate the role of the gut microbiome in anti-PD-1 immunotherapy (101, 102), including the first clinical study focusing on PD-1 checkpoint inhibitor response in metastatic melanoma. In a recent ground-breaking study, the authors were able to demonstrate a correlation between response to PD-1 checkpoint inhibitor treatment and diversity of microbial strains in the intestine. Moreover, high abundance of *Faecalibacterium* and low

abundance of *Bacteroidales* was associated with good prognosis and longer progression-free survival after checkpoint inhibitor treatment (101). Moreover, toxicity-related side effects and a dysfunctional intestinal epithelial barrier seem to be attenuated by a beneficial microbiota composition, potentially acting not only via modulation of the host immune response but also via modulation of cancer metabolism (103, 104).

Even though immunotherapy has been the treatment of choice for allergic diseases for more than a century, it seems that the cancer field is more advanced than allergy research with regards to evaluating the role of microbiota. There is a growing number of studies proving the major immunomodulating effects of intestinal microbiota and dietary supplementation with probiotic strains, as well as growing evidence demonstrating the role of intestinal microbial colonization in prevention and onset of allergic disease (9, 105). However, only a limited number of studies have so far evaluated the influence of beneficial bacterial strains in the context of successful allergen specific immunotherapy (AIT). Beneficial immunomodulatory and clinical effects of probiotic supplementation was observed when using different probiotic strains such as *L. rhamnosus* in a randomized clinical trial of peanut oral immunotherapy or in grass-pollen sublingual immunotherapy (SLIT) or by combining SCIT with *Clostridium butyricum* supplementation (106-108). Thus, major research efforts will be essential in the future to close this current knowledge gap and to ensure optimized AIT formulations for efficient treatment of allergic patients.

Microbial translocation in cancer and allergy

Under certain conditions, the normal gut epithelial barrier can become leaky, permitting the passage of microbes and microbial molecules into the systemic circulation, known as microbial translocation. Microbial products, especially cell walls of gram-negative bacteria (lipopolysaccharide [LPS] or endotoxin) are powerful stimulators of innate immunity and of B cell activation, resulting in inflammation that can lead to disease states, and even septic shock if present in very high doses (109). Microbial translocation can be assessed by

measuring the immune molecules stimulated by LPS, including soluble CD14 (sCD14), LPS-binding protein (LBP), and antibodies recognizing the core LPS antigen (Endocab) (109).

Gut barrier permeability is well-established in the setting of HIV infection. Two different prospective cohort studies reported that immune markers reflecting microbial translocation are associated with an elevated risk of AIDS-related non-Hodgkin lymphoma (NHL) (110, 111), presumably due to increased B cell activation resulting from the triggered innate inflammation signals. Both studies found that sCD14, measured years prior to diagnosis, was associated with a 2-4-fold increase in NHL risk, but the findings were inconsistent with respect to the other measures. Besides implications of microbial translocation in food allergy and asthma, it also strongly influences AD (112). A randomized trial among adult AD patients, given either selected probiotics or placebo, resulted in clinical improvement, decreased T-helper type 2 (Th2) immune activation and reduced plasma LPS as a measure of microbial translocation (112). Thus, microbial translocation may play a role in the etiology of both AIDS-related lymphoma and AD. It may therefore be possible to decrease risk by reducing microbial translocation through the selective use of probiotics.

Outlook: microbiota engineering

The beneficial effect of a balanced microbiota on human health can be restored or potentiated by external/medical intervention. There are three main routes to microbiota engineering (**Figure 2**). Firstly, in pathological conditions commensal bacteria could be administered to move towards a species composition more representative of a healthy microbiota. Examples are fecal transplantation and vaginal swab, but other types of more specific interventions are being studied. Secondly, bacterial biosynthetic pathways could be engineered for *de novo* or enhanced production of compounds that can promote healthier mucosal environments and homeostasis. Thirdly, selected commensal bacterial strains (especially lactic acid bacteria) could be engineered for the live delivery of recombinant therapeutic proteins either to prevent or to combat disease (113, 114). To this end, live recombinant protein delivery could occur via secretion, membrane-anchorage or intracellular

load. Engineering biosynthetic pathways or exogenous recombinant protein production implies the genetic modification of commensal bacteria, therefore sophisticated strategies to control their survival in the host and in the local environment are being devised (115). Cancer and allergy are widely the focus of many of these efforts (**Figure 2**), with a number of studies showing how microbiota composition and its engineering may ameliorate or support the appropriate immune response to contribute to clinical benefits for each condition (116).

6. Conclusion

The biodiversity hypothesis is a cornerstone in the understanding of the allergy epidemics. Subsequently, the era of microbiota research has opened up novel perspectives on allergy pathogenesis (117), but also on cancer due to the immunomodulatory properties of the mutualistic microbes. The impact of the microbiota can be direct or indirect by cellular cross-talk with and among innate or specific immune cells, by translocation of microbiota via epithelia into tissues, or by the exchange of molecules which may stimulate inflammatory or regulatory cells. The subsequent result can be inflammation or reconstitution, with opposing consequences in allergy and cancer. Importantly, the composition of the human microbiota can be disturbed by environmental factors, composition of the diet, and especially by medical interventions such as antibiotics and anti-ulcer medications. Phylogenetic variability and stable composition of microbiota seem to be determined in early life, defining this as a critical period for establishing health and homeostasis. However, it also offers a potential window of opportunity for interventions aiming at the establishment of a healthy microbiota to support long-term human health.

Abbreviations

APRIL	A proliferation-inducing ligand
AD	atopic dermatitis
BAFF	B-cell activating factor
ACT	adoptive T cell therapy
APC	antigen presenting cells
CNS	central nervous system
CCL2	gene of C-C motif chemokine 2
CTLA4	Cytotoxic T-Lymphocyte Associated Protein 4
DC	Dendritic cells
FFAR3	free fatty acid receptor 3
FOS	fructo-oligosaccharides
HDAC	histone deacetylases
GOS	galacto-oligosaccharides
GMP	glycomacropeptide
GM-CSF	Granulocyte-macrophage colony-stimulating factor
IL	interleukin
ILC	Innate lymphoid cells
LPS	Lipopolysaccharide
LBP	LPS-binding protein
NDO	Non-digestible Oligosaccharides
Nod-like	nucleotide-binding oligomerization domain-like
SCFAs	short chain fatty acids
HDAC	histone deacetylases

PD-1	Programmed cell death protein 1
PPAR	peroxisome proliferator-activated receptors
PRR	pattern recognition receptors
Th2	T-helper type 2
Treg	T-regulatory cell
GPR	G-Protein coupled receptor
HIV	human immunodeficiency virus
MAPK	Mitogen-activated protein kinase 1
NHL	non-Hodgkin lymphoma
PTEN	Phosphatidylinositol 3,4,5-trisphosphate
PTP	protein tyrosine phosphatase
QS	quorum sensing
ROS	reactive oxygen species
sCD14	soluble CD14
TLR	toll like receptor
TNF α	tumor necrosis factor

References:

1. Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI. The human microbiome project. *Nature* 2007;449(7164):804-10.
2. Jensen-Jarolim E, Bax HJ, Bianchini R, Capron M, Corrigan C, Castells M, et al. AllergoOncology - the impact of allergy in oncology: EAACI position paper. *Allergy* 2017;72(6):866-887.
3. Stokholm J, Blaser MJ, Thorsen J, Rasmussen MA, Waage J, Vinding RK, et al. Maturation of the gut microbiome and risk of asthma in childhood. *Nat Commun* 2018;9(1):141.
4. Arrieta MC, Stiemsma LT, Dimitriu PA, Thorson L, Russell S, Yurist-Doutsch S, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci Transl Med* 2015;7(307):307ra152.
5. Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosch D, et al. Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nat Med* 2016;22(10):1187-1191.
6. Ribeiro A, Heimesaat MM, Bereswill S. Changes of the Intestinal Microbiome-Host Homeostasis in HIV-Infected Individuals - A Focus on the Bacterial Gut Microbiome. *Eur J Microbiol Immunol (Bp)* 2017;7(3):158-167.
7. Wypych TP, Marsland BJ. Antibiotics as Instigators of Microbial Dysbiosis: Implications for Asthma and Allergy. *Trends Immunol* 2018.
8. Mitre E, Susi A, Kropp LE, Schwartz DJ, Gorman GH, Nylund CM. Association Between Use of Acid-Suppressive Medications and Antibiotics During Infancy and Allergic Diseases in Early Childhood. *JAMA Pediatr* 2018;172(6):e180315.
9. Diesner SC, Bergmayr C, Pfitzner B, Assmann V, Krishnamurthy D, Starkl P, et al. A distinct microbiota composition is associated with protection from food allergy in an oral mouse immunization model. *Clin Immunol* 2016;173:10-18.
10. Azad MB, Coneys JG, Kozyrskyj AL, Field CJ, Ramsey CD, Becker AB, et al. Probiotic supplementation during pregnancy or infancy for the prevention of asthma and wheeze: systematic review and meta-analysis. *BMJ* 2013;347:f6471.
11. Rastogi T, Devesa S, Mangtani P, Mathew A, Cooper N, Kao R, et al. Cancer incidence rates among South Asians in four geographic regions: India, Singapore, UK and US. *Int J Epidemiol* 2008;37(1):147-60.
12. Oikonomopoulou K, Brinc D, Kyriacou K, Diamandis EP. Infection and cancer: reevaluation of the hygiene hypothesis. *Clin Cancer Res* 2013;19(11):2834-41.
13. Gutensohn N, Cole P. Childhood social environment and Hodgkin's disease. *N Engl J Med* 1981;304(3):135-40.
14. Gilham C, Peto J, Simpson J, Roman E, Eden TO, Greaves MF, et al. Day care in infancy and risk of childhood acute lymphoblastic leukaemia: findings from UK case-control study. *BMJ* 2005;330(7503):1294.
15. Parodi S, Crosignani P, Miligi L, Nanni O, Ramazzotti V, Rodella S, et al. Childhood infectious diseases and risk of leukaemia in an adult population. *Int J Cancer* 2013;133(8):1892-9.
16. Josephs DH, Spicer JF, Corrigan CJ, Gould HJ, Karagiannis SN. Epidemiological associations of allergy, IgE and cancer. *Clin Exp Allergy* 2013;43(10):1110-23.
17. Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, Earley ZM, et al. Commensal *Bifidobacterium* promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* 2015;350(6264):1084-9.
18. Vogtmann E, Goedert JJ. Epidemiologic studies of the human microbiome and cancer. *Br J Cancer* 2016;114(3):237-42.
19. Fan X, Alekseyenko AV, Wu J, Peters BA, Jacobs EJ, Gapstur SM, et al. Human oral microbiome and prospective risk for pancreatic cancer: a population-based nested case-control study. *Gut* 2018;67(1):120-127.

20. Peters BA, Wu J, Pei Z, Yang L, Purdue MP, Freedman ND, et al. Oral Microbiome Composition Reflects Prospective Risk for Esophageal Cancers. *Cancer Res* 2017;77(23):6777-6787.
21. Hayes RB, Ahn J, Fan X, Peters BA, Ma Y, Yang L, et al. Association of Oral Microbiome With Risk for Incident Head and Neck Squamous Cell Cancer. *JAMA Oncol* 2018;4(3):358-365.
22. Rajilic-Stojanovic M, de Vos WM. The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS Microbiol Rev* 2014;38(5):996-1047.
23. Mehta RS, Nishihara R, Cao Y, Song M, Mima K, Qian ZR, et al. Association of Dietary Patterns With Risk of Colorectal Cancer Subtypes Classified by *Fusobacterium nucleatum* in Tumor Tissue. *JAMA Oncol* 2017;3(7):921-927.
24. O'Keefe SJ, Li JV, Lahti L, Ou J, Carbonero F, Mohammed K, et al. Fat, fibre and cancer risk in African Americans and rural Africans. *Nat Commun* 2015;6:6342.
25. Rossini A, Rumio C, Sfondrini L, Tagliabue E, Morelli D, Miceli R, et al. Influence of antibiotic treatment on breast carcinoma development in proto-neu transgenic mice. *Cancer Res* 2006;66(12):6219-24.
26. Boursi B, Mamtani R, Haynes K, Yang YX. Recurrent antibiotic exposure may promote cancer formation--Another step in understanding the role of the human microbiota? *Eur J Cancer* 2015;51(17):2655-64.
27. Roth-Walter F, Pacios LF, Bianchini R, Jensen-Jarolim E. Linking iron-deficiency with allergy: role of molecular allergens and the microbiome. *Metallomics* 2017;9(12):1676-1692.
28. Mandal S, Godfrey KM, McDonald D, Treuren WV, Bjornholt JV, Midtvedt T, et al. Fat and vitamin intakes during pregnancy have stronger relations with a pro-inflammatory maternal microbiota than does carbohydrate intake. *Microbiome* 2016;4(1):55.
29. Kull I, Bergstrom A, Melen E, Lilja G, van Hage M, Pershagen G, et al. Early-life supplementation of vitamins A and D, in water-soluble form or in peanut oil, and allergic diseases during childhood. *J Allergy Clin Immunol* 2006;118(6):1299-304.
30. Bhatia AK, Lee JW, Pinto HA, Jacobs CD, Limburg PJ, Rubin P, et al. Double-blind, randomized phase 3 trial of low-dose 13-cis retinoic acid in the prevention of second primaries in head and neck cancer: Long-term follow-up of a trial of the Eastern Cooperative Oncology Group-ACRIN Cancer Research Group (C0590). *Cancer* 2017;123(23):4653-4662.
31. Mondul AM, Watters JL, Mannisto S, Weinstein SJ, Snyder K, Virtamo J, et al. Serum retinol and risk of prostate cancer. *Am J Epidemiol* 2011;173(7):813-21.
32. Wolsk HM, Harshfield BJ, Laranjo N, Carey VJ, O'Connor G, Sandel M, et al. Vitamin D supplementation in pregnancy, prenatal 25(OH)D levels, race, and subsequent asthma or recurrent wheeze in offspring: Secondary analyses from the Vitamin D Antenatal Asthma Reduction Trial. *J Allergy Clin Immunol* 2017;140(5):1423-1429 e5.
33. Barry EL, Peacock JL, Rees JR, Bostick RM, Robertson DJ, Bresalier RS, et al. Vitamin D Receptor Genotype, Vitamin D3 Supplementation, and Risk of Colorectal Adenomas: A Randomized Clinical Trial. *JAMA Oncol* 2017;3(5):628-635.
34. Pacheco-Gonzalez RM, Garcia-Marcos L, Morales E. Prenatal vitamin D status and respiratory and allergic outcomes in childhood: A meta-analysis of observational studies. *Pediatr Allergy Immunol* 2018.
35. Luthold RV, Fernandes GR, Franco-de-Moraes AC, Folchetti LG, Ferreira SR. Gut microbiota interactions with the immunomodulatory role of vitamin D in normal individuals. *Metabolism* 2017;69:76-86.
36. Skaaby T, Taylor AE, Jacobsen RK, Mollehave LT, Friedrich N, Thuesen BH, et al. Associations of genetic determinants of serum vitamin B12 and folate concentrations with hay fever and asthma: a Mendelian randomization meta-analysis. *Eur J Clin Nutr* 2018;72(2):264-271.
37. Roy A, Kocak M, Hartman TJ, Vereen S, Adgent M, Piyathilake C, et al. Association of prenatal folate status with early childhood wheeze and atopic dermatitis. *Pediatr Allergy Immunol* 2018;29(2):144-150.

38. Brasky TM, White E, Chen CL. Long-Term, Supplemental, One-Carbon Metabolism-Related Vitamin B Use in Relation to Lung Cancer Risk in the Vitamins and Lifestyle (VITAL) Cohort. *J Clin Oncol* 2017;35(30):3440-3448.
39. Magnusdottir S, Ravcheev D, de Crecy-Lagard V, Thiele I. Systematic genome assessment of B-vitamin biosynthesis suggests co-operation among gut microbes. *Front Genet* 2015;6:148.
40. Maslowski KM, Mackay CR. Diet, gut microbiota and immune responses. *Nat Immunol* 2011;12(1):5-9.
41. Moro G, Arslanoglu S, Stahl B, Jelinek J, Wahn U, Boehm G. A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child* 2006;91(10):814-9.
42. van der Aa LB, Heymans HS, van Aalderen WM, Sprikkelman AB. Probiotics and prebiotics in atopic dermatitis: review of the theoretical background and clinical evidence. *Pediatr Allergy Immunol* 2010;21(2 Pt 2):e355-67.
43. Hogenkamp A, Knippels LM, Garssen J, van Esch BC. Supplementation of Mice with Specific Nondigestible Oligosaccharides during Pregnancy or Lactation Leads to Diminished Sensitization and Allergy in the Female Offspring. *J Nutr* 2015;145(5):996-1002.
44. Verheijden KA, Willemsen LE, Braber S, Leusink-Muis T, Delsing DJ, Garssen J, et al. Dietary galacto-oligosaccharides prevent airway eosinophilia and hyperresponsiveness in a murine house dust mite-induced asthma model. *Respir Res* 2015;16:17.
45. Tan J, McKenzie C, Vuillermin PJ, Govere G, Vinuesa CG, Mebius RE, et al. Dietary Fiber and Bacterial SCFA Enhance Oral Tolerance and Protect against Food Allergy through Diverse Cellular Pathways. *Cell Rep* 2016;15(12):2809-24.
46. Garrett WS. Cancer and the microbiota. *Science* 2015;348(6230):80-6.
47. Bhattarai Y, Kashyap PC. Agaro-oligosaccharides: a new frontier in the fight against colon cancer? *Am J Physiol Gastrointest Liver Physiol* 2016;310(6):G335-6.
48. Macfarlane GT, Steed H, Macfarlane S. Bacterial metabolism and health-related effects of galacto-oligosaccharides and other prebiotics. *J Appl Microbiol* 2008;104(2):305-44.
49. van Dokkum W, Wezendonk B, Srikumar TS, van den Heuvel EG. Effect of nondigestible oligosaccharides on large-bowel functions, blood lipid concentrations and glucose absorption in young healthy male subjects. *Eur J Clin Nutr* 1999;53(1):1-7.
50. Parodi PW. Cows' milk fat components as potential anticarcinogenic agents. *J Nutr* 1997;127(6):1055-60.
51. Frei R, Lauener RP, Cramer R, O'Mahony L. Microbiota and dietary interactions: an update to the hygiene hypothesis? *Allergy* 2012;67(4):451-61.
52. Thorburn AN, Macia L, Mackay CR. Diet, metabolites, and "western-lifestyle" inflammatory diseases. *Immunity* 2014;40(6):833-42.
53. Arpaia N, Campbell C, Fan X, Dikiy S, van der Veeken J, deRoos P, et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature* 2013;504(7480):451-5.
54. Arpaia N, Rudensky AY. Microbial metabolites control gut inflammatory responses. *Proc Natl Acad Sci U S A* 2014;111(6):2058-9.
55. Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat Med* 2014;20(2):159-66.
56. Lin L, Zhang J. Role of intestinal microbiota and metabolites on gut homeostasis and human diseases. *BMC Immunol* 2017;18(1):2.
57. Chinthrajah RS, Hernandez JD, Boyd SD, Galli SJ, Nadeau KC. Molecular and cellular mechanisms of food allergy and food tolerance. *J Allergy Clin Immunol* 2016;137(4):984-97.
58. Levy M, Thaiss CA, Zeevi D, Dohnalova L, Zilberman-Schapira G, Mahdi JA, et al. Microbiota-Modulated Metabolites Shape the Intestinal Microenvironment by Regulating NLRP6 Inflammasome Signaling. *Cell* 2015;163(6):1428-43.
59. Fujiya M, Musch MW, Nakagawa Y, Hu S, Alverdy J, Kohgo Y, et al. The *Bacillus subtilis* quorum-sensing molecule CSF contributes to intestinal homeostasis via OCTN2, a host cell membrane transporter. *Cell Host Microbe* 2007;1(4):299-308.

60. Okamoto K, Fujiya M, Nata T, Ueno N, Inaba Y, Ishikawa C, et al. Competence and sporulation factor derived from *Bacillus subtilis* improves epithelial cell injury in intestinal inflammation via immunomodulation and cytoprotection. *Int J Colorectal Dis* 2012;27(8):1039-46.
61. Segain JP, Raingeard de la Bletiere D, Bourreille A, Leray V, Gervois N, Rosales C, et al. Butyrate inhibits inflammatory responses through NF κ B inhibition: implications for Crohn's disease. *Gut* 2000;47(3):397-403.
62. Nastasi C, Candela M, Bonefeld CM, Geisler C, Hansen M, Krejsgaard T, et al. The effect of short-chain fatty acids on human monocyte-derived dendritic cells. *Sci Rep* 2015;5:16148.
63. Uribe-Herranz M, Bittinger K, Rafail S, Guedan S, Pierini S, Tanes C, et al. Gut microbiota modulates adoptive cell therapy via CD8 α dendritic cells and IL-12. *JCI Insight* 2018;3(4).
64. Chang PV, Hao L, Offermanns S, Medzhitov R. The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition. *Proc Natl Acad Sci U S A* 2014;111(6):2247-52.
65. Thaiss CA, Zmora N, Levy M, Elinav E. The microbiome and innate immunity. *Nature* 2016;535(7610):65-74.
66. Siebenhaar F, Redegeld FA, Bischoff SC, Gibbs BF, Maurer M. Mast Cells as Drivers of Disease and Therapeutic Targets. *Trends Immunol* 2018;39(2):151-162.
67. Forsythe P. Microbes taming mast cells: Implications for allergic inflammation and beyond. *Eur J Pharmacol* 2016;778:169-75.
68. Forsythe P, Wang B, Khambati I, Kunze WA. Systemic effects of ingested *Lactobacillus rhamnosus*: inhibition of mast cell membrane potassium (IKCa) current and degranulation. *PLoS One* 2012;7(7):e41234.
69. Karaki S, Mitsui R, Hayashi H, Kato I, Sugiya H, Iwanaga T, et al. Short-chain fatty acid receptor, GPR43, is expressed by enteroendocrine cells and mucosal mast cells in rat intestine. *Cell Tissue Res* 2006;324(3):353-60.
70. Kim MH, Kang SG, Park JH, Yanagisawa M, Kim CH. Short-chain fatty acids activate GPR41 and GPR43 on intestinal epithelial cells to promote inflammatory responses in mice. *Gastroenterology* 2013;145(2):396-406 e1-10.
71. Zhang H, Du M, Yang Q, Zhu MJ. Butyrate suppresses murine mast cell proliferation and cytokine production through inhibiting histone deacetylase. *J Nutr Biochem* 2016;27:299-306.
72. Schilderink R, Verseijden C, de Jonge WJ. Dietary inhibitors of histone deacetylases in intestinal immunity and homeostasis. *Front Immunol* 2013;4:226.
73. Bochner BS. The Eosinophil: for better or worse, in sickness and in health. *Ann Allergy Asthma Immunol* 2018.
74. Rosenberg HF, Masterson JC, Furuta GT. Eosinophils, probiotics, and the microbiome. *J Leukoc Biol* 2016;100(5):881-888.
75. Hosoki K, Nakamura A, Nagao M, Hiraguchi Y, Tokuda R, Wada H, et al. Differential activation of eosinophils by 'probiotic' *Bifidobacterium bifidum* and 'pathogenic' *Clostridium difficile*. *Int Arch Allergy Immunol* 2010;152 Suppl 1:83-9.
76. Kraemer LS, Brenner TA, Krumholz JO, Rosenberg HF. A flow-cytometric method to evaluate eosinophil-mediated uptake of probiotic *Lactobacillus reuteri*. *J Microbiol Methods* 2017;137:19-24.
77. Choi CY, Kim YH, Oh S, Lee HJ, Kim JH, Park SH, et al. Anti-inflammatory potential of a heat-killed *Lactobacillus* strain isolated from Kimchi on house dust mite-induced atopic dermatitis in NC/Nga mice. *J Appl Microbiol* 2017;123(2):535-543.
78. Wang X, Hui Y, Zhao L, Hao Y, Guo H, Ren F. Oral administration of *Lactobacillus paracasei* L9 attenuates PM2.5-induced enhancement of airway hyperresponsiveness and allergic airway response in murine model of asthma. *PLoS One* 2017;12(2):e0171721.
79. Jensen-Jarolim E, Bax HJ, Bianchini R, Crescioli S, Daniels-Wells TR, Dombrowicz D, et al. AllergoOncology: Opposite outcomes of immune tolerance in allergy and cancer. *Allergy* 2018;73(2):328-340.
80. Wawrzyniak M, O'Mahony L, Akdis M. Role of Regulatory Cells in Oral Tolerance. *Allergy Asthma Immunol Res* 2017;9(2):107-115.

81. Konieczna P, Ferstl R, Ziegler M, Frei R, Nehrass D, Lauener RP, et al. Immunomodulation by *Bifidobacterium infantis* 35624 in the murine lamina propria requires retinoic acid-dependent and independent mechanisms. *PLoS One* 2013;8(5):e62617.
82. Lyons A, O'Mahony D, O'Brien F, MacSharry J, Sheil B, Ceddia M, et al. Bacterial strain-specific induction of Foxp3+ T regulatory cells is protective in murine allergy models. *Clin Exp Allergy* 2010;40(5):811-9.
83. Konieczna P, Groeger D, Ziegler M, Frei R, Ferstl R, Shanahan F, et al. *Bifidobacterium infantis* 35624 administration induces Foxp3 T regulatory cells in human peripheral blood: potential role for myeloid and plasmacytoid dendritic cells. *Gut* 2012;61(3):354-66.
84. DeFalco J, Harbell M, Manning-Bog A, Baia G, Scholz A, Millare B, et al. Non-progressing cancer patients have persistent B cell responses expressing shared antibody paratopes that target public tumor antigens. *Clin Immunol* 2018;187:37-45.
85. Kim M, Kim CH. Regulation of humoral immunity by gut microbial products. *Gut Microbes* 2017;8(4):392-399.
86. Buchta CM, Bishop GA. Toll-like receptors and B cells: functions and mechanisms. *Immunol Res* 2014;59(1-3):12-22.
87. Rosser EC, Oleinika K, Tonon S, Doyle R, Bosma A, Carter NA, et al. Regulatory B cells are induced by gut microbiota-driven interleukin-1beta and interleukin-6 production. *Nat Med* 2014;20(11):1334-9.
88. Swanson PA, 2nd, Kumar A, Samarin S, Vijay-Kumar M, Kundu K, Murthy N, et al. Enteric commensal bacteria potentiate epithelial restitution via reactive oxygen species-mediated inactivation of focal adhesion kinase phosphatases. *Proc Natl Acad Sci U S A* 2011;108(21):8803-8.
89. Sun J, Kato I. Gut microbiota, inflammation and colorectal cancer. *Genes Dis* 2016;3(2):130-143.
90. Jiang L, Diaz PT, Best TM, Stimpfl JN, He F, Zuo L. Molecular characterization of redox mechanisms in allergic asthma. *Ann Allergy Asthma Immunol* 2014;113(2):137-42.
91. Nairz M, Schroll A, Haschka D, Dichtl S, Sonnweber T, Theurl I, et al. Lipocalin-2 ensures host defense against *Salmonella Typhimurium* by controlling macrophage iron homeostasis and immune response. *Eur J Immunol* 2015;45(11):3073-86.
92. Roth-Walter F, Schmutz R, Mothes-Luksch N, Zieglmayer P, Lemell P, Zieglmayer R, et al. Clinical efficacy of sublingual immunotherapy is associated with restoration of steady-state serum lipocalin 2 after SLIT: a pilot study. *World Allergy Organ J.* 2018;in press.
93. Jensen-Jarolim E, Pacios LF, Bianchini R, Hofstetter G, Roth-Walter F. Structural similarities of human and mammalian lipocalins, and their function in innate immunity and allergy. *Allergy* 2016;71(3):286-94.
94. Lippi G, Meschi T, Nouvenne A, Mattiuzzi C, Borghi L. Neutrophil gelatinase-associated lipocalin in cancer. *Adv Clin Chem* 2014;64:179-219.
95. Honda K, Littman DR. The microbiota in adaptive immune homeostasis and disease. *Nature* 2016;535(7610):75-84.
96. Malik A, Sharma D, Zhu Q, Karki R, Guy CS, Vogel P, et al. IL-33 regulates the IgA-microbiota axis to restrain IL-1alpha-dependent colitis and tumorigenesis. *J Clin Invest* 2016;126(12):4469-4481.
97. Dzidic M, Abrahamsson TR, Artacho A, Bjorksten B, Collado MC, Mira A, et al. Aberrant IgA responses to the gut microbiota during infancy precede asthma and allergy development. *J Allergy Clin Immunol* 2017;139(3):1017-1025 e14.
98. Ludvigsson JF, Neovius M, Ye W, Hammarstrom L. IgA deficiency and risk of cancer: a population-based matched cohort study. *J Clin Immunol* 2015;35(2):182-8.
99. Schwabe RF, Jobin C. The microbiome and cancer. *Nat Rev Cancer* 2013;13(11):800-12.
100. Viaud S, Daillere R, Boneca IG, Lepage P, Langella P, Chamaillard M, et al. Gut microbiome and anticancer immune response: really hot Sh*t! *Cell Death Differ* 2015;22(2):199-214.
101. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets TV, et al. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* 2018;359(6371):97-103.

102. Routy B, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillere R, et al. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 2018;359(6371):91-97.
103. Botticelli A, Zizzari I, Mazzuca F, Ascierto PA, Putignani L, Marchetti L, et al. Cross-talk between microbiota and immune fitness to steer and control response to anti PD-1/PDL-1 treatment. *Oncotarget* 2017;8(5):8890-8899.
104. Johnson CH, Spilker ME, Goetz L, Peterson SN, Siuzdak G. Metabolite and Microbiome Interplay in Cancer Immunotherapy. *Cancer Res* 2016;76(21):6146-6152.
105. Smolinska S, Groeger D, O'Mahony L. Biology of the Microbiome 1: Interactions with the Host Immune Response. *Gastroenterol Clin North Am* 2017;46(1):19-35.
106. Jerzynska J, Stelmach W, Balcerak J, Woicka-Kolejwa K, Rychlik B, Blauz A, et al. Effect of *Lactobacillus rhamnosus* GG and vitamin D supplementation on the immunologic effectiveness of grass-specific sublingual immunotherapy in children with allergy. *Allergy Asthma Proc* 2016;37(4):324-34.
107. Liu J, Chen FH, Qiu SQ, Yang LT, Zhang HP, Liu JQ, et al. Probiotics enhance the effect of allergy immunotherapy on regulating antigen specific B cell activity in asthma patients. *Am J Transl Res* 2016;8(12):5256-5270.
108. Tang ML, Ponsonby AL, Orsini F, Tey D, Robinson M, Su EL, et al. Administration of a probiotic with peanut oral immunotherapy: A randomized trial. *J Allergy Clin Immunol* 2015;135(3):737-44 e8.
109. Brenchley JM, Douek DC. Microbial translocation across the GI tract. *Annu Rev Immunol* 2012;30:149-73.
110. Epeldegui M, Magpantay L, Guo Y, Halec G, Cumberland WG, Yen PK, et al. A prospective study of serum microbial translocation biomarkers and risk of AIDS-related non-Hodgkin lymphoma. *AIDS* 2018.
111. Marks MA, Rabkin CS, Engels EA, Busch E, Kopp W, Rager H, et al. Markers of microbial translocation and risk of AIDS-related lymphoma. *AIDS* 2013;27(3):469-74.
112. Iemoli E, Trabattoni D, Parisotto S, Borgonovo L, Toscano M, Rizzardini G, et al. Probiotics reduce gut microbial translocation and improve adult atopic dermatitis. *J Clin Gastroenterol* 2012;46 Suppl:S33-40.
113. Wells JM, Mercenier A. Mucosal delivery of therapeutic and prophylactic molecules using lactic acid bacteria. *Nat Rev Microbiol* 2008;6(5):349-62.
114. Kumar M, Yadav AK, Verma V, Singh B, Mal G, Nagpal R, et al. Bioengineered probiotics as a new hope for health and diseases: an overview of potential and prospects. *Future Microbiol* 2016;11(4):585-600.
115. Rovner AJ, Haimovich AD, Katz SR, Li Z, Grome MW, Gassaway BM, et al. Recoded organisms engineered to depend on synthetic amino acids. *Nature* 2015;518(7537):89-93.
116. Cano-Garrido O, Seras-Franzoso J, Garcia-Fruitos E. Lactic acid bacteria: reviewing the potential of a promising delivery live vector for biomedical purposes. *Microb Cell Fact* 2015;14:137.
117. Lunjani N, Satitsuksanoa P, Lukasik Z, Sokolowska M, Eiwegger T, O'Mahony L. Recent Developments and Highlights in Mechanisms of Allergic Diseases: Microbiome. *Allergy* 2018; Epub ahead of print.

Figure legends

Figure 1. Interaction of microbiota with immune cells in health and disease. (A)

Bidirectional interaction between gut microbiome and players of the immune system. Cells from the epithelial barrier (epithelial cells and dendritic cells) sense the microbiome through the expression of a large panel of receptors. In health, such interactions are essential for the maintenance of the local microbial homeostasis and the integrity of epithelial barriers. This mechanism involves a variety of regulatory immune cells resident in the gut mucosa as well as molecules secreted by the epithelial layer. (B) Human eosinophils were shown to secrete, among others, high amounts of neurotoxin following phagocytosis of the pathogenic bacteria *Clostridium difficile* that were limited following the engulfment of *Bifidobacterium bifidum*. (C) In vitro interaction of human mast cells with *Lactobacillus rhamnosus* and SCFAs results in their modified functions and phenotypes with the implication of the KCa3.1 channel as well as Gpr41, 43 and 109a, respectively. (D) The differentiation of human monocytes into moDC in the presence of SCFAs induces modified secretory capacities compared to controls. (E) The efficacy of adoptive T cell therapy was proven to be associated with the microbiome. The success of adoptive T cell therapy correlates with a peripheral increase and a more abundant tumor infiltration of CD8 α ⁺DC producing IL-12 (1). Mouse treatment with SCFA results in the recruitment of bone marrow-derived DC with impaired capacity to induce Th2 responses in the lung (2). Probiotics administered by gavage reduces the severity of allergic airway inflammation and AD through the reduction of eosinophil infiltration into the lung and skin, respectively (3).

Figure 2. Microbiota engineering. The three main approaches to microbiota engineering are depicted: left, interventions to modify the microbiota species composition; right, engineering biosynthetic pathways for the production of compounds benefiting human health and homeostasis; bottom, engineering selected commensal strains for therapeutic protein live delivery. At the center of each circled approach a single bacterial cell is depicted as

commensal representative. These approaches have possible overlaps that allow better tailored design. All approaches can be envisaged to ameliorate or even prevent both allergy and cancer.

Table 1. Definitions

Term	Definition
Microbiota	Microorganisms (bacteria, viruses, fungi, protozoa and archaea) populating the inner and outer surfaces of the human body
Microbiome	Collective genome of all microorganisms
Mutualism	A relationship between two organisms of different species resulting in benefits for both organisms from the interaction
Symbionts	Two different organisms of the same or of different species with a close and persistent biological interaction




Table 2. Prevalent bacteria at different sites under normal homeostasis







	Phylum	Class	Order	Family	Species
Skin	Actinobacteria	Actinobacteria	Actinomycetales	Propionibacteriaceae	Propionibacterium acnes
	Firmicutes	Bacilli	Bacillales	Staphylococcaceae	Staphylococcus epidermis
Gut					
upper part	Firmicutes	Bacilli	Lactobacillales	Streptococcaceae	Streptococcus spp.
				Lactobacillaceae	Lactobacilli spp.
distal part	Firmicutes	Clostridia	Clostridiales	Clostridaceae	Clostridium spp
	Actinobacteria	Actinobacteria	Bifidobacteriales	Bifidobacteriaceae	Bifidobacterium spp.
	Bacteroidetes	Bacteroidia	Bacteriales	Bacteroidaceae	Bacteroides spp.
Lung	Bacteroidetes	Bacteroidia	Bacteriales	Prevotellaceae	Prevotella spp.
	Firmicutes	Bacilli	Lactobacillales	Streptococcaceae	Streptococcus spp.
	Proteobacteria	Gammaproteobacteria	Pasteurellales	Pasteurellaceae	Actinobacillus spp., etc.
Nose	Actinobacteria	Actinobacteria	Actinomycetales	Corynebacteriaceae	Corynebacterium spp.
	Firmicutes	Bacilli	Bacillales	Staphylococcaceae	Staphylococcus spp.

Table 3. Known modulators of the bacterial composition

	Phylum	Class	Order	Family	Species
Allergy					
<i>rather increased</i>	Firmicutes	Clostridia	Clostridiales	Clostridaceae	Clostridium spp.
				Ruminococcaceae	Ruminococcus spp.
<i>rather decreased</i>	Bacteroidetes	Bacteroidia	Bacteriodales	Bacteroidaceae	Bacteroides spp.
Diet					
<i>omnivore</i>	Firmicutes	Clostridia	Clostridiales	Lachnospiraceae	
<i>oligosacharrides</i>	Actinobacteria	Actinobacteria	Bifidobacteriales	Bifidobacteriaceae	Bifidobacterium spp.
<i>Western diet/fat</i>	Firmicutes	Erysipelotrichia	Erysipelotrichales	Erysiopelotrichaceae	
<i>Vitamin D</i>					
<i>rather decrease</i>	Firmicutes	Clostridia	Clostridiales	Lachnospiraceae	Coprococcus
	Actinobacteria	Actinobacteria	Bifidobacteriales	Bifidobacteriaceae	Bifidobacterium spp.

Table 4. Diet and microbiota metabolites in allergy and cancer

Compound	Structure	Effects in allergy and cancer
Miscellaneous		
Iron	Fe	Poor iron associated with allergy and high serum iron increases risk of several cancers
Folate		Serum folate is not correlated with allergic reactions, but with serum IgE. High plasma folate decreases risk of wheeze in children
Vitamins		
Vitamin A		Retinol supplementation has no effect on allergy. Vit A increases Proteobacteria. Increased serum retinol is associated with prostate cancer.
Vitamin D		Vitamin D supplementation reduces respiratory infections (not associated with allergy) and possibly reduces asthma incidence in offspring. Vitamin D reduces abundance of Bifidobacterium and Coprococcus and increases Proteobacteria. Vitamin D may prevent hematologic malignancies and advanced colorectal adenomas.

Non-digestible oligosaccharides		
Galacto-oligosaccharides		Bacterial fermentation of NDO induces a shift from <i>Bacteriodes</i> and <i>Prevotella</i> species to beneficial <i>Bifidobacterium</i> and <i>Lactobacillus</i> species
Fructo-oligosaccharides		Specific NDO feeding reduced the development of AD in infants at risk of allergy, which was associated with increased <i>Bifidobacterium</i> and <i>Lactobacilli</i> .
Agaro-oligosaccharides		Specific NDO reduce allergic manifestations in experimental models for allergy Specific NDO influence microbiota diversity and prevent the progression of colorectal cancer.
Short chain fatty acids		
Butyrate		Butyrate and propionate inhibit HDAC activity and promote Treg cells.
Propionate		
Acetate		Acetate does not inhibit HDAC.

