



King's Research Portal

DOI:
[10.1002/mdc3.12728](https://doi.org/10.1002/mdc3.12728)

Document Version
Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Martinez-Martin, P., Schrag, A-E., Weintraub, D., Rizos, A., Rodriguez-Blazquez, C., Ray Chaudhuri, K., & on behalf of EUROPAR and the IPMDS Non Motor PD Study Group (2019). Pilot study of the International Parkinson and Movement Disorder Society-sponsored Non-Motor Rating Scale (MDS-NMS). *Movement Disorders Clinical Practice*, 6(3), 227-234. <https://doi.org/10.1002/mdc3.12728>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Pilot study of the International Parkinson and Movement Disorder Society-sponsored Non-Motor Rating Scale (MDS-NMS)

Pablo Martinez-Martin, MD, PhD;¹ Anette Schrag, MD, PhD;² Daniel Weintraub, MD;^{3,4} Alexandra Rizos, MSc;^{5,6} Carmen Rodriguez-Blazquez, PhD;¹ Kallol Ray Chaudhuri MD, PhD;^{5,6} on behalf of the IPMDS Non Motor PD Study Group

1. National Center of Epidemiology and CIBERNED, Carlos III Institute of Health, Madrid, Spain
2. Department of Clinical Neurosciences, Royal Free Campus Institute of Neurology, University College London, London, United Kingdom
3. Departments of Psychiatry and Neurology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania, USA
4. Parkinson's Disease Research, Education and Clinical Center (PADRECC), Philadelphia Veterans Affairs Medical Center, Philadelphia, Pennsylvania, USA
5. Parkinson's Foundation International Centre of Excellence, King's College Hospital, London, United Kingdom
6. Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

Correspondence to: Dr. P. Martinez-Martin

National Center of Epidemiology
Carlos III Institute of Health
Avenida Monforte de Lemos, 5
28029 – Madrid
Spain

E-mail: pmartinez@isciii.es

Word counts: Abstract – 248; Text – 3000

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/mdc3.12728

Running title: Pilot study of the MDS-NMS

Key words: Parkinson's disease; Non-motor symptoms; New Non-Motor Symptoms Scale; MDS-NMS; Pilot study

Abstract

Background: Non-motor symptoms (NMS) are integral to Parkinson's disease (PD), and have a detrimental effect on patients and their caregivers. Clinical quantification has been aided by the development of comprehensive assessments such as the Non-Motor Symptoms Questionnaire (NMSQuest) and Scale (NMSS). The NMSS has been widely used in clinical studies and trials; however, since its validation in 2007 our understanding of NMS has changed substantially. With the support of the International Parkinson and Movement Disorder Society (IPMDS), after a detailed peer review an initiative to develop an updated version of NMSS, the MDS-NMS, was launched in 2015.

Objective: This paper encapsulates the data from the pre-validation phases, carried out under the auspices of the IPMDS Non-Motor PD Study Group.

Methods: Item selection and wording (formatted as a rater-based tool), were based on the NMSS, literature review, and expert consensus. Neurologists, PD patients, and healthy controls were included in the cognitive pretesting and administration of the preliminary version of the MDS-NMS. Primary data on acceptability and reliability were obtained.

Results: The pilot study, carried out in English in the United Kingdom and United States, demonstrated that the preliminary version of the MDS-NMS was comprehensive, understandable, and appropriate. Data quality was excellent; moderate floor effect was present in patients for most MDS-MNS domains, with some components showing weak internal consistency. The results led to additional instrument modifications.

Conclusion: Qualitative and quantitative research results have led to an updated NMSS, the definitive version of the MDS-NMS, which is currently being validated.

Introduction

Although James Parkinson described a range of non-motor symptoms (NMS) in his description of the “Shaking Palsy” 200 years ago, the impact of NMS on patients with Parkinson’s disease (PD) was largely overlooked for many years.^{1,2} Since the development and validation of assessment instruments such as the NMS Questionnaire (NMSQ)³ and the PD Non-Motor Symptoms Scale (NMSS),^{4,5} assessment of NMS has become more acceptable in clinical practice and in some countries (e.g., the UK) it is considered a quality standard as a Patient Reported Experience Measure (UK Parkinson’s Audit, 2017).⁶

The impact of the burden of NMS on quality of life has been articulated in a large number of publications,⁷⁻¹¹ and spans prodromal to the late stages of PD.¹²⁻¹⁵ The impact of the NMS has also underpinned the concept of the syndromic nature of PD,¹⁶ and non-motor endophenotypes have been postulated.¹⁷

The NMSS has been used as an outcome measure in several international clinical trials and has demonstrated sensitivity to change,¹⁸⁻²⁵ with published cut-off scores describing the different levels of NMS burden.^{26,27} However, the NMSS is now dated, as it was developed in 2005 and published in 2007. Key limitations in the current version of NMSS include, for example, deficiencies in the structure of this scale (e.g., items for depression, anxiety and apathy are in the same domain, as are sleep disorders and fatigue), lack of questions related to drug-induced issues such as non-motor fluctuations and impulse control disorders, and limited coverage of cognitive deficits.

After a detailed consultation, the International Parkinson and Movement Disorder Society (IPMDS) proposed the development of an updated scale to assess NMS, the IPMDS-Non-Motor Rating Scale (MDS-NMS), to take into account the following issues: (1) a critical appraisal of the experience of using NMSS in clinical trials and studies in over 1000 patients since 2007, including a diversity of epidemiological studies and a first approach to data mining;²⁸ (2) feedback from clinical raters who have used the

NMSS regarding individual questions in the NMSS; and (3) feedback from patient groups in relation to the use of NMSS in “real life”.

This multicentre effort showed that the current NMSS requires both consolidation and expansion, as well as refinement and improvement in relation to scoring and ease of application. Also, there is a growing demand for NMSS to be used in the context of new clinical trials, both those with and without a primary focus on NMS. Examples of clinical trials using the NMSS are RECOVER,¹⁹ PANDA,²² and TOLEDO,²⁹ and worldwide registries such as GLORIA, related to intrajejunal levodopa infusion therapy.³⁰

The development and cognitive pretesting of MDS-NMS has been completed³¹ and an international (UK and US) validation of the original, English version through IPMDS sponsorship is underway. The overall goal is to develop a standard, global PD NMS rating scale to be used in all clinical trials and epidemiological studies assessing clinical features of PD, regardless if the primary focus is on motor or non-motor symptoms.

The aims of the pilot study reported here were to explore the feasibility/acceptability, appropriateness, rank of the response options, identification of flaws (e.g., ambiguity, irrelevance and redundancy), and review of the format of the new MDS-NMS.

Methods

Development of the preliminary version of the MDS-NMS

The preliminary version of the MDS-NMS was drafted and refined by PD clinical researchers with expertise in non-motor symptoms and scale development in a series of person-meetings, video-conferences and by e-mail (from October 2013 to January 2016). It was composed of 63 items in 15 domains: A. Depression (5 items), B. Anxiety (4 items), C. Apathy (4 items), D. Psychosis (5 items), E. Impulse control & related disorders (4 items), F. Other neuropsychiatric symptoms (NPS) (4 items), G. Cognition (6 items), H. Orthostasis (2 items), I. Urinary (5 items), J. Sexual (2 items), K. Gastrointestinal (4 items), L. Thermoregulatory (2 items), M. Sleep & wakefulness (7 items), N. Pain (4 items), and O. Others (5 items). Items were selected from the NMSS

and other scales relating to NMS, review of the literature, and expert consensus. Items were phrased as questions on the presence of the symptoms, designed for administration by healthcare professionals (i.e., rater-administered).

Each item was scored twice based on five options, for frequency (0= Never, to 4= Majority of time) and severity (0= Not present to 4= Severe). Specific explanations for each of these anchors were provided. Item score was calculated by multiplying frequency x severity, total domain scores were calculated by the sum of their respective item scores, and a total instrument score generated by summing the domain scores represented total NMS burden. The MDS-NMS total score ranges in theory from 0 to 1008, although reaching the maximum score is very unlikely, given the low possibility of simultaneously reaching the maximum frequency and severity scores on all instrument items. This is similar to findings using the NMSS.²⁶

The MDS-NMS also includes an optional section to rate non-motor fluctuations (NMFs), including depression, anxiety, thinking or cognitive abilities, bladder symptoms, restlessness, pain, and fatigue. Each item was scored on a 5-point scale for “Typical degree of change” from “on” to “off” periods (0= No change; 4= Large) and for “Time spent in “off” state with non-motor symptoms” (0= No “off” time; 4= Majority of time, ≥51% of waking day), and the item score was calculated by multiplying the two components. The sum of individual item scores generated the total NMFs score.

The time frame evaluated is the “past two weeks”, as a longer time frame may be difficult to remember accurately for PD patients.

Design of the Pilot study

Open, cross-sectional international study, on a sample of users (neurologists), target population (PD patients), and healthy controls.

Participants

Three categories of subjects participated in the study: (1) Neurologists with expertise in PD, operationalized as neurologists attending ≥ 300 PD patients per year; (2) Patients with a diagnosis of idiopathic PD given by a neurologist, based on international criteria (Movement Disorder Society criteria),³² without significant cognitive impairment according to the judgment of the evaluating neurologist; (3) healthy community-dwelling controls from 50 to 80 years of age, and without PD, dementia, neurological or psychiatric disorders. Control participants were included as the scale may be used in patients in a diagnostic evaluation.

Exclusion criteria

Neurologists without experience in PD; patients with parkinsonism other than idiopathic PD, presence of significant cognitive impairment, psychiatric or medical conditions that would interfere with ability to complete study assessments, and current medication with known effect on mental state (cognitive or emotional); and controls with a comorbid disease or disorder at a moderate or severe level, as well as institutionalized persons. Also excluded were patients and controls unable to consent, with inability to read and write English or to complete written questionnaires correctly.

Sample size

Usually sample sizes from 20 to 40 representative participants are proposed for pilot studies on development of a scale or questionnaire.^{33,34} Due to the complexity of the preliminary version of the MDS-NMS a sample >50 neurologists and patients was proposed.

Other assessments

In addition to socio-demographic data, feedback questionnaires about the MDS-NMS were prepared for neurologists ($n=50$), patients ($n=15$), and controls ($n=15$). These questionnaires include questions about the word choice, understanding, relevance, appropriateness, comprehensiveness, length, comfortableness, issues with response options, and a section for additional comments or suggestions.

Procedures

Neurologists and patients participating in the study were informed of the objective of the study. Each neurologist completed the MDS-NMS for one patient and the “Questionnaire for the Neurologist”. In addition to being administered the MDS-NMS, each patient or control also completed the “Questionnaire for the Patient” or the “Questionnaire for Controls” about the instrument.

Data from the scale application and questionnaires were entered into a database for analyses. After analysis (see below), the developers made decisions for changes based on the results of the pilot study to create the final MDS-NMS for use in the formal validation study.

Ethical aspects

Each site contributing to the study received approval of their respective Ethics Committee/IRB for participation. In the UK, the study was approved for adoption to the UK clinical research network (UKCRN No 18003). Patients and controls provided written informed consent.

Data analysis

For depicting the characteristics of the sample, descriptive statistics (mean, standard deviation, percentage) were used. Total daily levodopa equivalent dose (LEDD) was calculated according to Tomlinson et al., 2010.³⁵ Data did not follow a normal distribution (Shapiro-Francia test).

For the MDS-NMS: Data quality (standard values: missing data <10%; full computable scores, >90%) and floor and ceiling effects (standard, $\leq 15\%$) and skewness (standard, from -1 to +1)^{36,37} were determined in PD patients and healthy controls.

Preliminary results of reliability (Cronbach’s alpha, standard value >0.70; inter-item correlation, 0.20-0.75; item homogeneity coefficient, >0.15; and corrected item-total correlation, ≥ 0.30) were explored only in patients.³⁷⁻⁴¹

The questionnaires for neurologists, patients, and controls were analyzed in a descriptive manner to assess their opinions and criticisms of the scale.

Results

Participant characteristics

Fifty-two neurologists and 69 PD patients (69.6% male) were included in the study. Patients' mean (\pm SD) age was 67.2 ± 9.5 (range: 37-90) years and the average years of education was 14.9 ± 4.5 (range: 8-30). Most of patients were married (72.5%) or single (15.9%), and retired (72.5%) or employed (18.8%). Mean age at PD onset was 59.9 ± 10.6 years (range: 32-84) and disease duration was 7.2 ± 5.2 years. LEDD was 665.0 ± 434.7 mg (range: 100-2080). Six patients (8.7%) had undergone deep brain stimulation surgery (4.7 years previously on average; range: 1-20).

Nineteen healthy controls (89.5% female) participated in the study. Their age was 53.4 ± 13.9 years (range: 24-68) and years of education were 17.0 ± 4.7 (range: 10-24) years. Most were married (68.4%) and employed (77.8%).

MDS-NMS scores

MDS-NMS scores for PD patients and healthy controls are shown in Table 1. Scores were higher in all domains for PD patients. Among patients, the possible maximum total score for a domain was reached only for the Sexual and Thermoregulatory domains, and among controls only for the Thermoregulatory domain

There were four missing data points (out of 4347; 0.1%) in the patient group: 1 in Item D3. Visual hallucinations; 2 in Item J2. Difficulty with sex; 1 in Item O2. Impaired olfaction. Fully computable data were available for 97.1%–100% of cases for the domain scores and 95.6% for the total score. The part P. Non-Motor Fluctuations was completed for 49 patients (71.0%). There were no missing data in the control group.

Acceptability

In PD patients, the MDS-NMS showed moderate floor effects, but no ceiling effect for the domains as the maximum total score of the domain was reached by only three patients (4.5%) in Sexual and by one patient (1.5%) in Thermoregulatory. The MDS-NMS total score showed no floor or ceiling effects (Table 2). In controls, floor effects ranged from 31.6% (M. Sleep & wakefulness) to 89.5% (domains C. Apathy, D. Psychosis, and E. Impulse control and related disorders), and ceiling effect was absent for all domains (Table 2). Differences in floor effects between patients and controls were statistically significant for domains A. Depression, B. Anxiety, C. Apathy, H. Orthostasis, I. Urinary, K. Gastrointestinal, N. Pain, and O. Others (p from 0.04 to <0.001) (Table 2). A moderate skewness was present in all scores of both groups (Table 2).

Internal consistency

Cronbach's alpha index was ≥ 0.70 for 10/15 domains and for the Non-motor fluctuations section (Table 3). The domains Impulse control and related disorders (0.45); Other NPS (0.58); Gastrointestinal (0.66); Thermoregulatory (0.57); and Others (0.63) obtained alpha values <0.70 . The item homogeneity coefficient for domains ranged from 0.17 (ICD & related disorders) to 0.73 (Apathy) (Table 3).

The individual items with the lowest intra-domain inter-item correlation were the Dopamine dysregulation syndrome, Pseudobulbar affect, Urinary incontinence, Drooling of saliva, Constipation, Restlessness of limbs, Snoring or apnea, and Double vision items (data not shown). Item-total correlation was lower than the standard 0.30 for seven items: Other hallucinations (auditory, etc.), Hobbyism, Dopamine dysregulation syndrome, Pseudobulbar affect, Hypomania or mania, Drooling of saliva, and Decreased sense of smell.

Qualitative responses regarding the instrument

Table 4 shows the distribution of the responses to the questions regarding the scale provided by the neurologists and 15 patients. Responses were "positive" regarding the characteristics of the scale in $>80\%$ of survey respondents, with the highest proportion

of “negative” opinions for the neurologists (23.1%) related to the long length of the scale. Particular comments of participants are summarized in the Supplementary material.

After careful consideration of the comments on the scale and the results of acceptability and internal consistency, the scale was modified as follows: (1) instructions for navigating across the scale scoring options were added; (2) item scoring assignment was more clearly specified; (3) 14 items were reworded; (4) several items were deleted: one from the domain Apathy, two from the domain Urinary, and one from the domain Sleep and wakefulness; (5) two domains (Other NPS and Thermoregulatory) were deleted; and (6) one item was moved from Thermoregulatory to Others domain, and one item was added to the subscale of Non-motor fluctuations. In summary, the revised, final version of the MDS-NMS was consolidated to a total 52 items across 13 domains, plus the subscale for Non-motor fluctuations with 8 items.

Discussion

NMS in PD can have a major and detrimental effect on quality of life of PD patients as well as their caregivers. NMS occur from the prodromal stage of PD right through to the palliative phase and appear to have a distinctive trajectory of progression which may not parallel motor progression.⁴² Much of this is related to a multi–neurotransmitter and varied brain involvement in the pathophysiological processes that may be operative in PD, and therefore clearer understanding of the range, nature and impact of NMS is crucial for improving management of PD. As an example, contrary to previous perception cognition is now thought to be affected even in the prodromal stage of PD, and dementia at diagnosis is thus no longer an exclusion factor for diagnosis of PD.^{13,14,43}

Value-based healthcare is now a vital aspect of healthcare delivery worldwide, and measurement of symptoms and their burden, are likely to become of day-to-day relevance in the management of PD. For example, the International Consortium for

Health Outcomes Measurement (ICHOM) has now published recommendations of quality standards for PD, and NMS form an integral part of the outcome process.⁴⁴

The NMSS has been used widely since its creation in 2005, highlighting the value of a comprehensive PD NMS rating scale. However, in light of the evolution of the field over the last 10+ years, updates and revisions were needed. The MDS-NMS was designed to overcome this unmet need as well as address non-motor fluctuations in PD, which are a frequently overlooked, yet very important, clinical syndrome. Furthermore, the scale also incorporates recognition and identification of impulse control disorders and related behavioral problems, now recognized to be a major clinical challenge in the management of PD worldwide.⁴⁵

Data from the pilot study of MDS NMS are presented in this paper and data quality was deemed excellent, with less than 1/1000 missing data points in the patient group. Over 95% of the domains and scale scores were directly computable without any imputation. In the control group there were no missing data at all.

The relatively high floor effect and skewness values observed in both groups is explained by the proportion of subjects who did not experience the respective symptoms. As the preliminary MDS-NMS was a broad, comprehensive scale including 63 symptoms in 15 domains, it was expected that a considerable proportion of patients would be free of many of those symptoms. NMS, as a whole, can be present in otherwise healthy populations, but with lower prevalence of these manifestations than in PD.³ Consistent with this reasoning, the floor effect was higher in the control group (except for the domain Other NPS).

Importantly, the floor effect was negligible for the total score, as was the ceiling effect for domains and total score in both groups. These results suggest an appropriate acceptability of this preliminary MDS-NMS version.

Internal consistency of the scale was, as a whole, adequate. Most of the domains showed alpha coefficients higher than or close to the standard 0.70. Most of the items showed appropriate inter-relationship with the others in their domain and with the respective corrected total score, but some others showed deficient performance regarding these properties and were deleted (e.g., Pseudobulbar affect). Nonetheless, after review of their wording, if necessary, some of these items (e.g., Drooling of saliva) were kept given their importance from a clinical perspective. The ICD & related disorders had low internal consistency, which is not surprising given the various symptoms or disorders included in this domain are distinct and often not co-morbid.

Cognitive pretesting comments and criticisms of the scale were gathered from user stakeholders (Supplementary material). Some critical comments related to item content and wording, led to revisions. Comments conflicted with each other, predictable (e.g., embarrassment related to questions about sex), or expressing opinions solely based on personal experience (e.g., item considered not relevant because the subject has not that symptom) were not considered when modifying the MDS-NMS.

A limitation of this study was the predominance of women in the healthy controls, a fact that can be influencing the differences observed between groups. Nonetheless, the main objectives of this pilot study should be barely affected by such discrepancy.

In conclusion, this pilot study of the preliminary version of the MDS-NMS showed that the scale was comprehensive, understandable, and appropriate, according to the vast majority of participants. Complementary statistical analysis showed that feasibility and acceptability of the preliminary MDS-NMS scale were satisfactory, with floor effect in most of the domains, due to the proportion of patients not experiencing those symptoms, but not in the total score. Most of the internal consistency parameters were adequate, but some items showed clear flaws in this attribute, a fact that promoted their deletion. Both the information furnished by the participants and the primary analytic data led to modifications that resulted in the definitive version of the MDS-NMS to be validated in the next phase of the project.

Acknowledgements

Data capture was aided by involvement of the National Institute of Health Research (NIHR) central research network support and staff in the UK: in King's College Hospital: Anna Sauerbier, Lauren Perkins, Dhaval Trivedi, Miriam Parry, Anne Martin, Nikolay Dimitrov, and Mubasher Quamar; in Royal Free Hospital: Kareem Khan, Olaitan Okunuye, and Joy Read; in Luton and Dunstable Hospital: Yvynne Croucher and Susanne Tluk; in Lister Hospital: Norman Kock, Diana Cedron, and Tracey Smith. In the US data capture was performed by: Eugenia Mamikonyan, Jacqui Rick, Benjamin Deck and Sam Rudovsky.

For authors KRC and AR, this paper represents independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

The authors also acknowledge advisory inputs from several members of the MDS Non-Motor PD Study Group.

Author Roles

1. Research project: A. Conception: KRC, PMM, AS, DW; B. Organization: KRC, PMM, AS, DW, AR; C. Execution: KRC, AS, DW, AR.
2. Statistical Analysis: A. Design: PMM, CRB; B. Execution: PMM, CRB; C. Review and Critique: KRC, AS, DW, AR.
3. Manuscript Preparation: A. Writing of the first draft: PMM; B. Review and Critique: KRC, AS, DW, AR, CRB.

Funding Sources and Conflict of Interest

The authors declare that there are no conflicts of interest relevant to this work.

This study was funded by the International Parkinson and Movement Disorder Society (IPMSD).

Financial Disclosures for the previous 12 months

PMM – Honoraria: from Editorial Viguera for lecturing in courses; International Parkinson and Movement Disorder Society for management of the Program on Rating Scales; Air Liquide, Abbvie, Zambon, and HM Hospitales de Madrid for advice in clinic-epidemiological studies. License fee payments for the King's Parkinson's Disease Pain scale.

AS – Research funding from National Institute of Health (NIHR), UCLH Biomedical Research Centre, Parkinson's UK, International Parkinson and Movement Disorder Society, GE Healthcare, Economic and Social Research Council UK, European Commission; honoraria for consultancy from Medtronic and Roche Pharmaceuticals, travel funding from Bial Pharmaceuticals.

DW – Research funding or support from Michael J. Fox Foundation for Parkinson's Research, National Institutes of Health (NINDS), Department of Veterans Affairs, Alzheimer's Therapeutic Research Initiative, Alzheimer's Disease Cooperative Study, and the International Parkinson and Movement Disorder Society; honoraria for consultancy from Acadia, Alkahest, Anavex Life Sciences, BlackThorn Therapeutics, Bracket, Clintrex LLC, Sunovion, Theravance Biopharma, and the CHDI Foundation; license fee payments from the University of Pennsylvania for the QUIP and QUIP-RS; royalties from Wolters Kluweland; and fees for legal consultation for three lawsuits related to medication prescribing in patients with Parkinson's disease.

AR – Salary support from the NIHR (National Institute for Health Research) Clinical Research Network South London.

CRB – Nothing to disclose.

KRC – Advisory Boards or consultancy with AbbVie, UCB, Sunovion, Pfizer, Jazz Pharma, GKC, Bial, Cynapsus, Lobsor, Stada, Zambon, Profile. Honoraria for lectures in symposium by AbbVie, Britannia, Bial, Zambon. Grants: Britannia Pharmaceuticals, AbbVie, GKC, Bial. Academic grants: Parkinson's UK, NIHR, PDNMG, EU (Horizon 2020), IMI, Kirby Laing Foundation, NPF. Intellectual property, Royalties: KPP scale, PDSS-2 scale with Mapi Institute; books, with Elsevier and Fastfacts.

Compliance with Journal Ethical Publication Guidelines Statement

- Institutional review board or ethics committee that approved the study: (1) National Research Ethics Service (NRES) Committee East Midlands – Northampton, UK; (2) Institutional Review Board at the Perelman School of Medicine at the University of Pennsylvania, USA.
- Declaration of patient consent - All patients and controls participating in the study received information about the objective of the study and provided written informed consent.
- We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this work is consistent with those guidelines."

References:

1. Chaudhuri KR, Jenner P. Two hundred years since James Parkinson's essay on the shaking palsy-Have we made progress? Insights from the James Parkinson's 200 years course held in London, March 2017. *Mov Disord.* 2017; 32: 1311-1315.
2. Obeso JA, Stamelou M, Goetz CG, et al. Past, present, and future of Parkinson's disease: A special essay on the 200th Anniversary of the Shaking Palsy. *Mov Disord.* 2017; 32: 1264-1310.
3. Chaudhuri KR, Martinez Martin P, Schapira AHV et al. An international multicenter pilot study of the first comprehensive self-completed non-motor symptoms questionnaire for Parkinson's disease: The NMSQuest study. *Mov Disord* 2006; 21: 916-923.
4. Chaudhuri KR, P Martinez-Martin, RG Brown, et al. The metric properties of a novel non-motor symptoms scale for Parkinson's disease: results from an international pilot study. *Mov Disord.*2007; 22: 1901-1911.
5. Martinez-Martin P, Rodriguez-Blazquez C, Abe K, et al. International study on the psychometric attributes of the non-motor symptoms scale in Parkinson disease. *Neurology.* 2009; 73: 1584-1591.

6. UK Parkinson's Audit, 2017. Transforming care.
<https://www.parkinsons.org.uk/professionals/uk-parkinsons-audit-transforming-care>.
Accessed 27 Aug 2018.
7. Martinez-Martin P, Rodriguez-Blazquez C, Kurtis MM, Ray Chaudhuri K, on Behalf of the NMSS Validation Group. The impact of non motor symptoms on health-related quality of life of patients with Parkinson's disease. *Mov Disord* 2011; 26: 399-406.
8. Santos-García D, de la Fuente-Fernández R. Impact of non-motor symptoms on health-related and perceived quality of life in Parkinson's disease. *J Neurol Sci.* 2013; 332: 136-140.
9. Kim SR, So HY, Choi E, Kang JH, Kim HY, Chung SJ. Influencing effect of non-motor symptom clusters on quality of life in Parkinson's disease. *J Neurol Sci.* 2014; 347: 310-315.
10. Erro R, Picillo M, Vitale C, et al. The non-motor side of the honeymoon period of Parkinson's disease and its relationship with quality of life: a 4-year longitudinal study. *Eur J Neurol.* 2016; 23: 1673-1679.
11. Prakash KM, Nadkarni NV, Lye WK, Yong MH, Tan EK. The impact of non-motor symptoms on the quality of life of Parkinson's disease patients: a longitudinal study. *Eur J Neurol.* 2016; 23: 854-860.
12. Iranzo A, Fernández-Arcos A, Tolosa E, et al. Neurodegenerative disorder risk in idiopathic REM sleep behavior disorder: study in 174 patients. *PLoS One.* 2014; 9: e89741.
13. Schrag A, Horsfall L, Walters K, Noyce A, Petersen I. Prediagnostic presentations of Parkinson's disease in primary care: a case-control study. *Lancet Neurol.* 2015; 14: 57-64.
14. Chahine LM, Weintraub D, Hawkins KA, et al. Cognition in individuals at risk for Parkinson's: Parkinson associated risk syndrome (PARS) study findings. *Mov Disord.* 2016; 31: 86-94.
15. Schrag A, Siddiqui UF, Anastasiou Z, Weintraub D, Schott JM. Clinical variables and biomarkers in prediction of cognitive impairment in patients with newly diagnosed Parkinson's disease: a cohort study. *Lancet Neurol.* 2017; 16: 66-75.
16. Titova N, Martinez-Martin P, Katunina E, Chaudhuri KR. Advanced Parkinson's or "complex phase" Parkinson's disease? Re-evaluation is needed. *J Neural Transm (Vienna).* 2017; 124: 1529-1537.

17. Titova N, Chaudhuri KR. Non-motor Parkinson disease: new concepts and personalised management. *Med J Aust.* 2018; 208: 404-409.
18. Martinez-Martin P, Reddy P, Antonini A, et al. Chronic subcutaneous infusion therapy with apomorphine in advanced Parkinson's disease compared to conventional therapy: a real life study of non motor effect. *J Parkinsons Dis.* 2011;1: 197-203.
19. Trenkwalder C, Kies B, Rudzinska M, et al. Rotigotine effects on early morning motor function and sleep in Parkinson's disease: a double-blind, randomized, placebo-controlled study (RECOVER). *Mov Disord.* 2011; 26: 90-99.
20. Fasano A, Ricciardi L, Lena F, Bentivoglio AR, Modugno N. Intrajejunal levodopa infusion in advanced Parkinson's disease: long-term effects on motor and non-motor symptoms and impact on patient's and caregiver's quality of life. *Eur Rev Med Pharmacol Sci.* 2012; 16: 79-89.
21. Martinez-Martin P, Reddy P, Katzenschlager R, et al. EuroInf: a multicenter comparative observational study of apomorphine and levodopa infusion in Parkinson's disease. *Mov Disord.* 2015; 30: 510-516.
22. Trenkwalder C, Chaudhuri KR, Martinez-Martin P, et al. Prolonged-release oxycodone-naloxone for treatment of severe pain in patients with Parkinson's disease (PANDA): a double-blind, randomised, placebo-controlled trial. *Lancet Neurol.* 2015; 14: 1161-1170.
23. Dafsari HS, Reddy P, Herchenbach C, et al. Beneficial Effects of Bilateral Subthalamic Stimulation on Non-Motor Symptoms in Parkinson's Disease. *Brain Stimul.* 2016; 9: 78-85.
24. Kurcova S, Bardon J, Vastik M, et al. Bilateral subthalamic deep brain stimulation initial impact on nonmotor and motor symptoms in Parkinson's disease: An open prospective single institution study. *Medicine (Baltimore).* 2018; 97: e9750.
25. Dafsari HS, Silverdale M, Strack M, et al. Nonmotor symptoms evolution during 24 months of bilateral subthalamic stimulation in Parkinson's disease. *Mov Disord.* 2018; 33: 421-430.
26. Ray Chaudhuri K, Rojo JM, Schapira AHV, et al. A proposal for a comprehensive grading of Parkinson's disease severity combining motor and non-motor assessments: meeting an unmet need. *PLoS ONE* 2013; 8: e57221.
27. Martinez-Martin P, Ray Chaudhuri K. Comprehensive grading of Parkinson's disease using motor and non-motor assessments: addressing a key unmet need. *Expert Rev Neurother.* 2018; 18: 41-50.

28. Mu J, Chaudhuri KR, Bielza C, de Pedro-Cuesta J, Larrañaga P, Martinez-Martin P. Parkinson's Disease Subtypes Identified from Cluster Analysis of Motor and Non-motor Symptoms. *Front Aging Neurosci.* 2017; 9: 301.
29. Katzenschlager R, Poewe W, Rascol O, et al. Apomorphine subcutaneous infusion in patients with Parkinson's disease with persistent motor fluctuations (TOLEDO): a multicentre, double-blind, randomised, placebo-controlled trial. *Lancet Neurol.* 2018; 17: 749-759.
30. Antonini A, Poewe W, Chaudhuri KR, et al. Levodopa-carbidopa intestinal gel in advanced Parkinson's: Final results of the GLORIA registry. *Parkinsonism Relat Disord.* 2017; 45: 13-20.
31. Ray Chaudhuri K, Weintraub D, Schrag A, Martinez-Martin P, On behalf of EUROPAR, The MDS Non-Motor PD Study Group. The International Parkinson and Movement Disorder Society–Non-Motor rating Scale (MDS-NMS): Results from the cognitive pre-testing and phases 1 and 2 of an international validation. *Mov Disord* 2016; 31(Suppl 2): S516.
32. Postuma RB, Berg D, Stern M, et al. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord.* 2015; 30: 1591-1601.
33. Johanson GA, Brooks GP. Initial scale development: Sample size for pilot studies. *Educ Psychol Measurement* 2010; 70: 394-400.
34. Tilley BC, LaPelle NR, Goetz CG, Stebbins GT; MDS-UPDRS Task Force. Using cognitive pretesting in scale development for Parkinson's disease: the Movement Disorder Society Unified Parkinson's Disease Rating Scale (MDS-UPDRS) example. *J Parkinsons Dis.* 2014; 4: 395-404.
35. Tomlinson CL, Stowe R, Patel S, Rick C, Gray R, Clarke CE. Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Mov Disord.* 2010; 25: 2649-2653.
36. McHorney CA, Tarlov AR. Individual-patient monitoring in clinical practice: are available health status surveys adequate? *Qual Life Res* 1995; 4: 293-307.
37. Hobart JC, Riazi A, Lamping DL, Fitzpatrick R, Thompson AJ. Improving the evaluation of therapeutic interventions in multiple sclerosis: development of a patient-based measure of outcome. *Health Technol Assess* 2004; 8: 9.
38. Scientific Advisory Committee of the Medical Outcomes Trust. Assessing health status and quality-of-life instruments: Attributes and review criteria. *Qual Life Res* 2002; 11: 193–205.

39. Piedmont RL. Inter-item correlations. In: Michalos AC. Encyclopedia of Quality of Life and Well-Being Research. Dordrecht: Springer Netherlands, 2014; pp. 3303-3304.
40. Smith SC, Lamping DL, Banarjee S, et al. Measurement of health-related quality of life for people with dementia: development of a new instrument (DEMQOL) and an evaluation of current methodology. *Health Technol Assess* 2005; 9: 10.
41. Clark LA, Watson D. Constructing validity: Basic issues in objective scale development. *Psychol Assessm* 1995; 7: 309-319.
42. Schapira AHV, Chaudhuri KR, Jenner P. Non-motor features of Parkinson disease. *Nature Rev Neurosci* 2017; 18: 435-450.
43. Postuma RB, Berg D, Stern M, et al. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord.* 2015 ;30: 1591-601.
44. de Roos P, Bloem BR, Kelley TA, et al. A consensus set of outcomes for Parkinson's disease from the International Consortium for Health Outcomes Measurement. *J Parkinsons Dis.* 2017; 7: 533-543.
45. Weintraub D, Koester J, Potenza MN, et al. Impulse control disorders in Parkinson disease: a cross-sectional study of 3090 patients. *Arch Neurol.* 2010; 67: 589-595.

Table 1 – Movement Disorder Society-Non-Motor Symptoms scale scores

PD patients	% F.C.	Mean	SD	Median	IQR	Observed		Theoret.
						Min	Max	Max
A. Depression	100	8.17		3.00	0-8	0	72	80
B. Anxiety	100	6.94	13.98	3.00	0-8	0	45	64
C. Apathy	100	7.32	11.01	3.00	0-8	0	60	64
D. Psychosis	98.5	1.85	13.02	0.00	0-0	0	25	80
E. ICD & related disorders	100	1.19	5.05	0.00	0-0	0	18	64
F. Other NPS	100	1.81	3.26	0.00	0-2	0	21	64
G. Cognition	100	10.80	3.67	4.00	1-14.5	0	64	96
H. Orthostasis	100	2.80	15.04	0.00	0-3.5	0	21	32
I. Urinary	100	10.28	5.01	4.00	0-13	0	72	80
J. Sexual	97.1	4.06	15.56	0.00	0-4	0	32	32
K. Gastrointestinal	100	4.51	7.97	2.00	0-6	0	50	64
L. Thermoregulatory	100	3.68	7.08	0.00	0-5.5	0	32	32
M. Sleep & wakefulness	100	13.33	6.19	9.00	3.5-17.5	0	88	112
N. Pain	100	8.91	15.66	4.00	0-15	0	56	64
O. Others	98.5	9.40	11.90	7.00	1-14.8	0	67	80
MDS-NMS Total score	95.64	91.55	10.78	54.00	28.5-125.5	10	491	1008
P. NM fluctuations	71.01	7.71	100.50	0.00	0-9	0	112	112
Controls								
A. Depression	100	2.37	6.00	0.00	0-1	0	22	80
B. Anxiety	100	1.95	4.12	0.00	0-2	0	18	64
C. Apathy	100	1.11	3.35	0.00	0-0	0	12	64
D. Psychosis	100	0.21	0.63	0.00	0-0	0	2	80
E. ICD & related disorders	100	0.68	2.21	0.00	0-0	0	9	64
F. Other	100	2.11	3.49	0.00	0-2	0	12	64
G. Cognition	100	4.89	9.69	2.00	0-4	0	35	96
H. Orthostasis	100	0.26	0.73	0.00	0-0	0	3	32
I. Urinary	100	2.74	9.12	0.00	0-1	0	40	80
J. Sexual	100	0.63	1.86	0.00	0-0	0	8	32
K. Gastrointestinal	100	1.11	1.94	0.00	0-2	0	7	64
L. Thermoregulatory	100	2.89	7.74	0.00	0-1	0	32	32
M. Sleep & wakefulness	100	3.26	5.61	2.00	0-4	0	24	112

N. Pain	100	2.63	5.77	0.00	0-4	0	24	64
O. Others	100	1.63	5.74	0.00	0-0	0	25	80
MDS-NMS Total score	100	28.47	39.44	12.00	6-28	0	151	1008

PD: Parkinson's disease. SD: Standard deviation. MDS-NMS: Movement Disorder Society-Non-Motor Symptoms scale. NM fluctuations: Non-motor fluctuations.

% F.C.: Percentage of fully computable scores. IQR: Interquartile range. Theoret.: Theoretical.

Table 2 – Acceptability parameters of the preliminary Movement Disorder Society-Non-Motor Symptoms scale

	Parkinson's disease patients			Healthy controls		
	Floor effect (%)	Ceiling effect (%)	Skewness	Floor effect (%)	Ceiling effect (%)	Skewness
A. Depression	29.0**	0	2.67	68.4	0	2.85
B. Anxiety	27.5*	0	2.36	52.6	0	3.63
C. Apathy	39.1***	0	2.64	89.5	0	2.92
D. Psychosis	76.5	0	3.21	89.5	0	2.80
E. ICD & related disorders	79.7	0	3.43	89.5	0	3.48
F. Other NPS	53.6	0	3.49	52.6	0	1.93
G. Cognition	18.8	0	1.89	36.8	0	2.63
H. Orthostasis	55.1*	0	2.16	84.2	0	3.31
I. Urinary	33.3**	0	1.95	68.4	0	4.22
J. Sexual	61.2*	4.5	2.45	63.2	0	3.84
K. Gastrointestinal	31.9*	0	4.21	63.2	0	2.03
L. Thermoregulatory	55.1	1.5	2.24	73.7	0	3.38
M. Sleep & wakefulness	7.3**	0	2.57	31.6	0	3.14
N. Pain	34.8*	0	1.82	63.2	0	3.19
O. Others	22.1***	0	2.53	78.9	0	4.18
MDS-NMS total	0**	0	2.43	10.5	0	2.17
P. Non-motor fluctuations	51.0***	2.0	4.64	--	--	--

MDS-NMS: Movement Disorder Society-Non-Motor Symptoms scale.

Statistically significant difference between patients and controls for floor effect:

*, p<0.05; **, p<0.01; ***, p<0.001. The rest of differences was not significant.

Table 3 – Data of internal consistency of the preliminary MDS-NMS in Parkinson’s disease patients

	Cronbach’s alpha	Inter-item correlation	Item homogeneity Index	Item-total correlation
A. Depression	0.91	0.41 – 0.57	0.67	0.73 – 0.84
B. Anxiety	0.85	0.37 – 0.68	0.59	0.53 – 0.82
C. Apathy	0.91	0.41 – 0.70	0.73	0.76 – 0.83
D. Psychosis	0.59	0.19 – 0.62	0.38	0.07 – 0.69
E. ICD & related disorders	0.42	-0.04 – 0.56	0.17	0.04 – 0.48
F. Other NPS	0.61	-0.06 – 0.59	0.25	0.19 – 0.76
G. Cognition	0.85	0.19 – 0.62	0.48	0.51 – 0.87
H. Orthostasis*	0.79	0.70	0.65	-
I. Urinary	0.84	0.05 – 0.64	0.51	0.42 – 0.83
J. Sexual*	0.72	0.43	0.56	-
K. Gastrointestinal	0.63	0.03 – 0.22	0.33	0.23 – 0.60
L. Thermoregulatory*	0.54	0.32	0.40	-
M. Sleep & wakefulness	0.75	-0.03 – 0.41	0.31	0.33 – 0.55
N. Pain	0.77	0.13 – 0.56	0.45	0.40 – 0.66
O. Others	0.49	-0.01 – 0.41	0.26	0.04 – 0.43

P. Non-motor fluctuations	0.93	0.10 – 0.80	0.64	0.68 – 0.85
---------------------------	------	-------------	------	-------------

MDS-NMS: Movement Disorder Society–Non-Motor Symptoms scale

* Each of these domains has only two items

Table 4 – Synthesis of the responses of neurologists and patients about the MDS-NMS

	Neurologists (n=52)			Patients (n=15)		
		N	%		N	%
1. Do you find the scale relevant?	No	1	1.9	No	1	6.7
	Yes	51	98.1	Yes	13	86.6
	NR	0	0.0	NR	1	6.7
2. Does this scale help you to better understand your patient's / your current health state?	No	6	11.5	No	2	13.3
	Yes	46	88.5	Yes	13	86.7
	NR	0	0.0	NR	0	0.0
3. Do you think this scale is comprehensive?	No	5	9.6	No	2	13.3
	Yes	46	88.5	Yes	13	86.7
	NR	1	1.9	NR	0	0.0
4. Do you think this scale is too long?	No	40	76.9	No	13	86.7
	Yes	12	23.1	Yes	2	13.3
	NR	0	0.0	NR	0	0.0
5. Do you find the questions easy to understand?	No	3	5.8	No	1	6.7
	Yes	48	92.3	Yes	14	93.3
	NR	1	1.9	NR	0	0.0
6. Did you find any question(s) embarrassing?	No	49	94.2	No	14	93.3
	Yes	3	5.8	Yes	1	6.7
	NR	0	0.0	NR	0	0.0
7. Did you find any particular question(s) difficult to answer?	No	42	80.8	No	14	93.3
	Yes	10	19.2	Yes	1	6.7
	NR	0	0.0	NR	0	0.0

MDS-NMS: Movement Disorder Society Non-Motor Symptoms scale.

NR: No response.