



King's Research Portal

DOI:

[10.1136/annrheumdis-2018-214676](https://doi.org/10.1136/annrheumdis-2018-214676)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Dregan, A., Matcham, F., Harber-Aschan, L., Rayner, L., Brailean, A., Davis, K., Hatch, S. L., Pariente, C., Armstrong, D., Stewart, R., & Hotopf, M. (2019). Common mental disorders within chronic inflammatory disorders. A primary care database prospective investigation. *Annals of the Rheumatic Diseases*, 78(5), 688-695. <https://doi.org/10.1136/annrheumdis-2018-214676>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**Common mental disorders within chronic inflammatory disorders. A primary care
database prospective investigation**

**Dregan Alex PhD,¹ Matcham Faith PhD,¹ Harber-Aschan Lisa PhD,² Rayner Lauren
PhD,¹ Brailean Annamaria PhD,¹ Davis Katrina MSc,¹ Hatch Stephani L. PhD,¹
Pariante Carmine, PhD,¹ Armstrong David MD,³ Stewart Rob, MD^{1,4} Hotopf Matthew,
PhD^{1,4}**

¹ Department of Psychological Medicine, Institute of Psychiatry Psychology and
Neuroscience, King's College London.

² Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden.

³ School of Population Health and Environmental Sciences, King's College London.

⁴ South London and Maudsley NHS Foundation Trust, London, UK.

Correspondence to: Dr Alex Dregan, Department of Psychological Medicine, Institute of
Psychiatry Psychology and Neuroscience, De Crespigny Park, King's College London,
London, SE5 8AF, UK. Email: alexandru.dregan@kcl.ac.uk.

Word count

Abstract: 250

Text: 3000

Objectives: There is inconsistent evidence about the association between inflammatory disorders with depression and anxiety onset in a primary care context. The study aimed evaluate the risk of depression and anxiety within multisystem and organ-specific inflammatory disorders.

Methods: Prospective cohort study with primary care patients from the UK Clinical Practice Research Datalink diagnosed with an inflammatory disorder between 1st of January 2001 and 31st of December 2016. These patients were matched on age, gender, practice, and index date with patients without an inflammatory disorder. The study exposures were seven chronic inflammatory disorders. Clinical diagnosis of depression and anxiety represented the outcome measures of interest.

Results: Among 538,707 participants, the incidence of depression ranged from 14 per 1000 person-years (severe psoriasis) to 9 per 1000 person-years (systemic vasculitis), substantively higher compared to their comparison group (5 to 7 per 1000 person-years). Hazard ratios (HR) of multiple depression and anxiety events were 16% higher within inflammatory disorders (HR, 1.16, 95%CI 1.12-1.21, $p < 0.001$) compared to the matched comparison group. The incidence of depression and anxiety was strongly associated with the age at inflammatory disorder onset. The overall HR estimate for depression was 1.90 (95%CI, 1.66-2.17, $p < 0.001$) within early onset disorder (<40 years of age) and 0.93 (95%CI, 0.90-1.09, $p = 0.80$) within late-onset of disorder (≥ 60 years of age).

Conclusions: Primary care patients with inflammatory disorders have elevated rates of depression and anxiety incidence, particularly those patients with early onset inflammatory disorders. This finding may reflect the impact of the underlying disease on patients' quality of life, although the precise mechanisms requires further investigation.

Keywords: Autoimmune diseases, Epidemiology, Inflammation, Mental Health, Primary care

Key messages

What is already known on this subject?

- Several cross-sectional studies suggested but did not establish a contributory role of inflammation in the initiation of depression and anxiety within patients diagnosed with chronic inflammatory disorders.

What does this study add?

- In a prospective cohort study with 538,707 patients from primary care, a significant increment in the onset of new of depression and anxiety events was documented within organ-specific and multisystemic inflammatory disorders.
- The incidence of depression or anxiety varied with the age at inflammatory disorder onset.

How might this impact on clinical practice or future developments?

- The elevated risk of depression and anxiety means clinicians should be vigilant for early symptoms of depressive or anxiety in this highly at-risk group of patients.
- The risk was greater among patients with younger age at inflammatory disorder onset, supporting tailored preventative approaches early in the course of a chronic disorder.
- The study, however, does not demonstrate a causal relationship between inflammation with depression and anxiety.

Introduction

A growing body of evidence indicated that low-grade inflammation may play an influential role in the onset of depression and anxiety.¹ Past research has linked upregulated pro-inflammatory cytokines and increased levels of acute-phase reactants with changes in neurotransmitter and neuroendocrine functioning related to psychiatric disorders.^{2,3} This evidence supports a link between depression and anxiety with inflammatory disorders (e.g. rheumatoid arthritis (RA), psoriasis, ankylosing spondylitis (AS)), and cross-sectional studies are in line with this suggestion.⁴⁻⁷ Evidence from prospective studies exploring the role of inflammatory disorders in depression and anxiety onset were, however, inconsistent.^{8,9} Little is known about the incidence of depression or anxiety across clinically diverse inflammatory disorders. Differences in genetic influences and treatment choices across inflammatory disorders may lead to variation in depression or anxiety onset.^{10,11} The genetic association with human leukocyte antigen (HLA) alleles, for instance, was stronger within AS compared to RA.¹⁰ Quantifying the extent to which the link between inflammatory disorders with depression vary by individual disorders may suggest mechanisms underlying specific relationships and ultimately facilitate targeted preventative approaches. There is substantive variation in the age of onset across individual inflammatory disorders that may also lead to differential association with depression or anxiety,^{12,13} in turn more prevalent in early adult years. The incidence of depression or anxiety, thus, may be lower across disorders with late age at onset (e.g. RA) than those with early age at onset (e.g. Crohn's disease). The detection of disparities in mental health burden could guide treatment choice and effective tailoring of healthcare resources. The aim of the present study was to implement a prospective cohort study within a large primary care database to test the hypothesis that the incidences of depression or anxiety varied across specific inflammatory disorders. It was also hypothesised that depression or anxiety risk was greatest within people with an early age at disorder onset.

Methods

Data

A prospective matched cohort study design was implemented in the Clinical Practice Research Datalink (CPRD), among the world largest electronic medical records database. CPRD collects routine primary care data on over 14 million patients (≈ 6.7 million active) from around 675 practices throughout the UK National Health Service (NHS). All patients in the NHS are registered with a general practice that provides all their primary care and coordinates secondary and community care. Important diagnostic and therapy information from referrals to secondary or community care services are captured by primary care records. Patients included in the CPRD are broadly representative of the UK's wider population in terms of age, gender, and ethnicity.¹⁴ The validity and accuracy of CPRD diagnostic and prescription data have been demonstrated across a wide range of disorders including cancer,¹⁵ stroke,¹⁶ COPD,¹⁷ depression and anxiety,¹⁸ rheumatoid arthritis,¹⁹ inflammatory bowel disorders,²⁰ and autoimmune disorders.^{21 22}

Study population

A cohort of primary care patients aged >18 years with a first-ever diagnosis of a chronic inflammatory disorder (psoriasis, Crohn's disease (CD) and ulcerative colitis (UC), RA, SLE, ankylosing spondylitis (AS), and systemic vasculitis (SV)) recorded between 1st of January 2001 and 30th of September 2016, who were depression or anxiety disorders free at the time of inflammatory disorder diagnosis were sampled from the CPRD. The date of diagnosis was defined as the index date. The index date for patients transferring into the practice was their practice registration date and the practice up-to-standard (UTS) date was used if a practice joined the data base during the recruitment period. The end of recruitment was the earliest of 30th of September 2016 or the death date or transferred out of the practice date. Patients

below the age of 18 at the time of diagnosis were excluded from the study sample because the presentation and course of inflammatory disorders might be different in younger people.²³ All diagnoses were derived from the medical codes recorded by family physicians in patients' electronic health records. These patients were matched (a 1:2 ratio of inflammatory exposed to 2 matched non-exposed) on age (year of birth), gender, practice, and index date with a group of patients without a chronic inflammatory disorder selected for this study during the recruitment period. Matched controls were assigned the index date of the inflammatory disorder diagnosis of the matched case. Similar to the inflammatory patients, matched controls with a diagnosis of depression or anxiety before the assigned index date were excluded from the analyses. Psoriasis patients are commonly classified into severe if they were prescribed a systemic therapy (i.e. methotrexate, azathioprine, cyclosporine, hydroxyurea) or phototherapy (psoralen and ultraviolet A) during the study period, or into mild psoriasis if no such treatment was recorded.^{24 25} This classification has been validated with similar databases²⁶⁻²⁸ and has also been used in this study. Data were extracted from the CPRD in September 2017.

Outcome

The study primary outcome measures were a new Read medical code for a diagnosis of depression or anxiety used as binary variables (yes/no).²⁹ The date of the first outcome code following an inflammatory disorder diagnosis was referred to as the outcome index date. Depression was broadly defined to include single episode of depression, recurrent depression events, and bipolar depressive events to allow for the possibility that chronic inflammation is implicated across the wider spectrum of the depressive disorder. In keeping with other studies,³⁰ anxiety was broadly defined to include generalised anxiety disorders, phobias, panic attacks, and panic disorders.

Covariates

Factors known to be associated with chronic inflammation and depression or anxiety were adjusted for in the analyses. These covariates included age (continuous), gender (male vs female), body mass index (<18.5, 18.5 to 25, >25 to <30, 30 to <35, and ≥ 35 kg/m²), index of multiple deprivation (quintiles), blood pressure (BP) (<120 mmHG, normal; 120-139 mmHg, borderline; ≥ 140 mmHg, hypertension), smoking (ex- or current vs never), drinking (ex- or current vs never), physical comorbidities (yes/no) (i.e. cancer, diabetes, stroke, coronary heart disease (CHD), dementia, epilepsy, chronic obstructive pulmonary disease (COPD), liver disorders, kidney disorders, insomnia), stressful life events (e.g. stress at home or at work), together with prescription of statins, antihypertensive, anti-diabetic, and hypnotics. Previous studies^{31 32} linked corticosteroids and non-steroidal anti-inflammatory drugs (NSAIDs) with increased risk of depression and were therefore also included as covariates. For each potential confounder the value closer to the index date and within the five years period prior to a chronic inflammation diagnosis was included. For instance, if a patient had two BP measures within 5 years prior to the baseline (e.g. at 4 years and at 2 years prior to baseline), the value closer to the study baseline (e.g. at 2 years) was included in the analyses. Expanding the baseline period to available data was found to enhance covariate sensitivity by capturing data that would otherwise be missed.³³

Statistical analysis

The analyses were conducted in a time-to-event framework. Failure was classed as a new diagnosis of depression or anxiety. Participants contributed person-time to the analysis from the study start date (the later of the start of the participant's record in CPRD or the diagnosis date for a chronic inflammatory condition). Follow-up ended at the earliest of the study outcome date, the end of a participant's registration, the last date of CPRD data collection, or

date of death. All participants had at least 12 months of follow-up recorded had at least one medical event recorded from the study start date to the study end date.

A Cox proportional-hazards model for clustered data based on the matched pairs was implemented with the use of a multiple-failure events to allow for the possibility that each patient may experience more than one outcome event.³⁴ This approach permits analysis of data for each of multiple outcomes in a single model, allowing the most efficient use of each patient's data and reducing problems of multiple testing.²⁴ The multiple-failure model avoids the need to censor records at earlier outcome events or to test hypotheses separately for each outcome. Robust variance estimator was used to adjust for the dependency introduced by the matching variables. This approach is considered³⁵ superior to matched stratification as it allows for unbiased estimation of hazard ratios. Because confounding by matching variables cannot be excluded,³⁶ the estimation models adjusted for: matching variables (age, gender, practice, index year) and all study covariates listed above. A similar estimation was performed to estimate whether depression or anxiety onset varied with the age (<40, 40-49, 50-59, and 60 or over) at inflammatory disorder diagnosis. Additional analyses estimated the specific associations between each inflammatory condition with depression and anxiety in separate Cox regression models with robust estimate variance. The analyses used the Efron method to handle tied events. Forest plots were used to present measures of association for age subgroups and individual inflammatory disorder. A random-effects meta-analysis was implemented to evaluate heterogeneity by chronic inflammatory disorder and overall.³⁷ The proportionality assumption was tested and confirmed using Schoenfeld residuals. As this was an exploratory study no adjustment for multiple comparisons was made, and therefore marginally significant results may be type I errors. Several sensitivity analyses were conducted. Firstly, alternative follow-up times were used by starting the follow-up immediately after the inflammatory disorder diagnosis. Secondly, depression and anxiety

were redefined to include both a clinical diagnosis code plus a relevant prescription (i.e. antidepressant or anxiolytic drugs, respectively). Thirdly, stratification by matched pairs was implemented to account for the matching. Fourthly, to test the robustness of psoriasis findings, data on systemic therapy were used to classify RA and systemic vasculitis patients (the only sufficiently powered disorders) into mild (no systemic therapy) and severe (systemic therapy). The effect of competing risk on mortality was also assessed. Multiple imputation by chained equation was used to handle missing data. The analyses were implemented using Stata version 15.

Results

The analyses included 180163 patients with chronic inflammatory disorders (see Table 1) that were individually matched for age, gender, practice, and index date with 358544 control patients without a diagnosis of chronic inflammation. The median duration of follow-up was around 4 years for patients and controls. While clinical diagnosis and therapy data were comprehensive, among lifestyle factors missing information ranged from around 6% for smoking to 22% for alcohol status. Selective baseline characteristics of study participants are described in Table 1 (See Table S1 in Supplementary material for full data description).

Insert Table 1 about here

Figure 1 shows the across all inflammatory disorders, the incidence of both depression or anxiety was greater within cases compared to the matched controls. The highest incidence rate was observed within severe psoriasis (14 per 1000 person-years), followed by those diagnosed with CD and AS (12 per 1000 person-years). Similar trends emerged with regards to the incidence of anxiety (Figure S1 in the Supplement).

Insert Figure 1 about here

Table 2 presents the results of the analyses by study outcomes indicating increased hazard rates of depression and anxiety across all chronic inflammatory disorders. The strongest association was observed for severe psoriasis, being associated with a 71% increased rate of new depression onset (hazard ratio (HR) 1.71; 95% confidence interval (CI), 1.52-1.93, p -value <0.0001) compared to the matched comparisons. Regarding new anxiety onset, patients diagnosed with AS presented with the largest hazard rates (1.36, 1.23-1.51, p <0.0001) compared to their matched comparison group. Age-related analyses revealed higher depression and anxiety incidence among persons with early inflammatory disorder onset. Kaplan-Meier survival curves are presented in Figure S2, Supplementary material.

Insert Table 2 about here

Figure 2 displays the results from the multiple outcome models, with patients being allowed to experience either depression or anxiety in a random order. Compared to the matched group, patients with an inflammatory disorder experienced a 16% (1.16, 1.12-1.21, p <0.001) increased risk of depression or anxiety events. Patients diagnosed with CD presented with the highest hazard ratio (1.23, 1.13-1.33, p <0.001), while those with mild psoriasis with the lowest hazard ratio (1.08, 1.03-1.13, p <0.001). Age-based analyses (Figure 3) indicated that the pooled hazard rate for multiple depression or anxiety incidence was 1.71(1.59-1.84, p <0.001) among patients with early inflammatory disorder onset (<40 years of age), which declined to 0.93 (0.85-1.01, $p=0.080$) among those with late disorder onset (≥ 60 years) (See Figure S3 in the Supplement for findings among 40 to 49 and 50 to 59 years of age groups).

Insert Figure 2 about here

Insert Figure 3 about here

Sensitivity analyses

Sensitivity analyses using a more stringent criteria for depression and anxiety definition (e.g. clinical diagnosis plus corresponding drug prescriptions) resulted in modestly higher estimates, validating the robustness of the main findings (Figure S4 and Table S1). Analyses stratified by matched pairs endorsed the estimates and associations of the study findings. Systemic therapy-based sensitivity analyses indicated that both severe RA (1.43, 1.28-1.59, $p < 0.001$) and systemic vasculitis (1.65, 1.20-2.25, $p < 0.001$) presented greater risk of depression incidence relative to mild RA (1.36, 1.25-1.49, $p < 0.01$) or mild systemic vasculitis (1.42, 1.27-1.60, $p < 0.001$).

Discussion

The main aim of the present study was to provide a comprehensive understanding about the burden of common mental disorders across specific inflammatory disorders within a primary care context. Within a large prospective design, several clinically diverse inflammatory disorders presented with a consistently elevated risk of depression and anxiety incidence. In particular, a 16% overall increased risk of multiple depression and anxiety events emerged across seven specific chronic inflammatory disorders (RA, psoriasis, CD, UC, SLE, SV, and AS). Associations were observed between incident depression with each specific inflammatory disorder, although the effect size was of lower magnitude than suggested by findings based on secondary care-based populations.⁴ The reason for this discrepancy may be that a smaller proportion of patients with inflammatory disorders, those with most severe or active disease, are seen in secondary care.³⁸ In our study, the incidence of depression and

anxiety was higher for patients with severe psoriasis relative to those with mild psoriasis. This suggestion was substantiated in sensitivity analyses among RA and systemic vasculitis disorders.

The pooled incidence of depression and anxiety was considerably increased (71% increment) among primary care patients with early onset inflammatory disorder (<40 years of age) and less so (-7%) among those with late disorder onset (≥ 60 years of age). Early onset inflammatory disorders are associated with more widespread inflammation, increased frequency of active disease, and more aggressive disease manifestation and treatment compared to late-onset disorder.³⁹ Whether the increased incidence of depression or anxiety within early disorder onset was caused by increased disease activity or delay in disorder diagnosis and treatment (or their combined effect), needs further exploration.

All seven chronic disorders analysed in this study are connected by common underlying inflammatory mechanism and the consistently elevated rates of depression and anxiety incidence across them might support a potential role of inflammation in the pathogenesis of these disorders, though this suggestion was not directly tested in this study. The main alternative hypothesis that cannot be excluded from this study design is that depression and anxiety may represent emotional responses to the experience of living with a distressing and often debilitating inflammatory disorder. The psychosocial and physical effects of the inflammatory disorder might therefore contribute to the onset of depressive and anxiety symptoms. For example, increased depression and anxiety incidence among primary care patients with early disorder onset, as found in this study, may reflect these patients presenting with more extensive and severe manifestations of the inflammatory disorder.⁴⁰ The elevated rates of depression events among patients with severe psoriasis relative to those with mild

psoriasis seem to be in line with a disease response hypothesis. Pain, disfigurement, loneliness, and stigma associated with severe inflammatory disease indicators (e.g. eruptive psoriasis, multiple nail lesions), for example, could worsen patients' sleep quality and prevent them from full social participation, leading to the onset of depressive symptoms.⁴¹

The results of the present study raise important questions about the assessment and management of common mental health disorders among younger patients diagnosed with specific inflammatory disorders. Irrespective of whether psychological problems are the consequence of the emotional reaction to disease and disability or of a common inflammatory etiology, there seems a clear association between inflammatory disorders and depression or anxiety, especially for younger early-onset patients. Routine assessments of patients' mental health could lead to improved prevention and treatment outcomes. If further research supports the common inflammatory etiology hypothesis, then clinical intervention might target the inflammatory response itself. Renewed interest in the potential effectiveness of immunomodulatory therapies (e.g. new biologics, methotrexate) for the prevention of treatment-resistant depression may indicate one way forward.

Previous prospective studies explored the association between depression and anxiety with specific inflammatory disorders.⁹⁴² Marrie et al.,⁹⁴³ for example, documented somewhat higher incidence rates of depression and anxiety among patients with RA and IBD. Marie et al.' studies did not adjust for differences in chronic illnesses (e.g. CVD, diabetes, CKD) at baseline, did not account for matching in their analyses, used a different case definition (e.g. exclusion of cases within a 5-year period from index date), and relied on a more local population. These variations may account for the observed differences in effect size between our and Marrie et al. findings. Meesters et al.⁴⁴ also documented higher incidence rates of

depression events among AS patients from primary care compared to our findings, possibly due to previous study failure to adjust for other covariates apart from age and gender. An earlier study found no increased risk of depression among patients diagnosed with CD or ulcerative colitis⁴²: this may reflect previous study's lack of a comparison group or shorter follow-up (<5 years). Recent studies^{6,27} indicated greater incidence of depression among patients diagnosed with severe psoriasis relative to those with mild psoriasis, as observed in this study. The decline in depression incidence with age at disorder onset is in line with an earlier systematic review among RA patients,⁴ and extends previous findings to anxiety.

Strengths and limitations

The present study has several strengths including nationally representative primary care population, prospective study design, and clinically valid diagnoses of inflammatory disorders, depression and anxiety. The inclusion of primary care patients with systemic and organ-specific inflammatory disorders, ensures the generalisability of the study findings to real-world clinical practice. While our data possibly contains all diagnoses issued within primary care, it may be less complete with regards to diagnoses made in secondary or community care.⁴⁵ Nine out of ten adults with mental health disorders are supported in primary care in the UK, implying that only a small number of cases are not captured by the CPRD. The use of antidepressant and anxiolytics therapies as sensitivity analyses may have also mitigated against diagnostic bias, given that drug prescribing is often considered a reliable marker for case identification.⁴⁶ Clinicians may be more alert (or ask different questions) to depressive or anxiety symptoms among patients with inflammatory disorders due to increased contact with the healthcare system, and thus more likely to identify relevant cases. The mean number of primary care consultations, however, was similar between inflammatory patients and matched controls (data not shown). The precise timings of the

onset of exposure or outcome measures cannot be determined precisely in observational data, precluding robust causal inferences. To mitigate against this concern, the analyses excluded outcome measures that occurred during the first 12 months following an inflammatory disorder diagnosis. Our large study sample comprised a heterogeneous group of patients with distinctive underlying disease severity and symptomatology, potentially masking subgroups of patients that could present with clinically significant mental disorders. This suggestion is supported by our finding with regards to severe psoriasis and age at inflammatory disorder diagnosis. A method of analysis that did not allow for matching might give slightly wider CIs and larger p values than a matched analysis.⁴⁷ Sensitivity analyses that adjusting for matching validated the study main findings. We cannot exclude the possibility that the comparison group included patients diagnosed with other less common inflammatory disorders (e.g. bullous skin diseases, Sjogren syndrome). This concern may have attenuated the true risk of depression or anxiety within chronic inflammatory disorders. The study primary aim was to model initial inflammatory disorder status (e.g. psoriasis, RA, SLE) and therapy (e.g. NSAIDs, corticosteroids), along with patients' socio-demographic and clinical data to patients' overall risk for future depression or anxiety onset. The analyses, however, did not model potential post-diagnosis mediators and moderators for depression or anxiety onset, including temporary changes in underlying disease severity, treatment choices, and inflammatory responses. These are clinically relevant questions that deserve detailed investigation with future prospective studies. The study only differentiated between mild and severe psoriasis. The smaller sample of patients within the rest of inflammatory disorders precluded a similar classification. This concern also applied to patients with psoriatic arthritis that were classified as psoriasis. Given that the definition of severe of psoriasis was based on DMARD exposure, however, it is possible that patients with psoriatic arthritis were included in the severe psoriasis subgroup. Sensitivity analyses within RA and systemic vasculitis

disorders endorsed psoriasis severity results increasing confidence in the robustness of the study findings. Future studies with larger IBD, SLE, and AS samples are also required to confirm the link between inflammatory disorder severity with study outcomes. Missing data on lifestyle covariates can compromise the results of statistical analysis but use of multiple imputation and appropriate sensitivity analyses should have mitigated some of this risk. A larger proportion of women were diagnosed with AS in this study, which is contrary to other studies showing higher AS rates among men.⁴⁸ The study findings about the incidence of depression or anxiety may, thus, not be generalisable to the wider AS population. This concern was likely caused by the matching of patients and controls on gender, leading to intentional non-representativeness. In analytical studies where the aim is to explore exposure-outcome association (as in this study), however, population representativeness is not considered necessary or desirable.⁴⁹ Richiardi et al.,⁵⁰ for instance, suggested that non-representativeness increases the power to assess main effects and effect modification, and that valid statistical inferences can be made when adjusting for confounders. Primary care patients diagnosed with an inflammatory disorder were at greater risk of new depression and anxiety onset compared to matched patients without an inflammatory disorder, a risk that was particularly elevated among patients with early onset of chronic inflammatory disorder. These findings may reflect either a response to the physical effects of living with a chronic inflammatory disorder, or a role for inflammation in the genesis of depression and anxiety. The latter hypothesis deserves further attention as it may offer the opportunity for new therapeutic approaches to anxiety and depression, but first the question of whether depression is a consequence of inflammation or is a reaction to experiencing a chronic illness deserves further exploration. Studies evaluating modifiable mediators for depression and anxiety incidence across specific inflammatory disorders are also warranted.

Acknowledgment

This study is based in part on data from the CPRD obtained under license from the UK Medicines and Healthcare Products Regulatory Agency. However, the interpretation and conclusions contained in this report are those of the authors alone.

The study was approved by the Independent Scientific Advisory Committee (reference No. 17_036RA).

Contributors AD, MF, AB, KD, CP, DA, RS, SH, and MH were responsible for study design. AD was responsible for data acquisition. AD analysed the data and wrote the manuscript. All authors critically revised the manuscript and approved the final version.

Competing interest: None declared

Funding/Support: No funding

References

1. Kiecolt-Glaser JK, Derry HM, Fagundes CP. Inflammation: depression fans the flames and feasts on the heat. *Am J Psychiatry* 2015;172(11):1075-91. doi: 10.1176/appi.ajp.2015.15020152.
2. Cepeda MS, Stang P, Makadia R. Depression Is Associated With High Levels of C-Reactive Protein and Low Levels of Fractional Exhaled Nitric Oxide: Results From the 2007-2012 National Health and Nutrition Examination Surveys. *J Clin Psychiatry* 2016;77(12):1666-71. doi: 10.4088/JCP.15m10267.
3. Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom Med* 2009;71(2):171-86. doi: 10.1097/PSY.0b013e3181907c1b.
4. Matcham F, Rayner L, Steer S, et al. The prevalence of depression in rheumatoid arthritis: a systematic review and meta-analysis. *Rheumatology (Oxford)* 2013;52(12):2136-48. doi: 10.1093/rheumatology/ket169.
5. Lamb RC, Matcham F, Turner MA, et al. Screening for anxiety and depression in people with psoriasis: a cross-sectional study in a tertiary referral setting. *Br J Dermatol* 2017;176(4):1028-34. doi: 10.1111/bjd.14833.
6. Jensen P, Ahlehoff O, Egeberg A, et al. Psoriasis and New-onset Depression: A Danish Nationwide Cohort Study. *Acta Derm Venereol* 2016;96(1):39-42. doi: 10.2340/00015555-2183.
7. Ng A, Tam WW, Zhang MW, et al. IL-1beta, IL-6, TNF- alpha and CRP in Elderly Patients with Depression or Alzheimer's disease: Systematic Review and Meta-Analysis. *Sci Rep* 2018;8(1):12050. doi: 10.1038/s41598-018-30487-6.

8. Watad A, Bragazzi NL, Adawi M, et al. Anxiety disorder among rheumatoid arthritis patients: Insights from real-life data. *J Affect Disord* 2017;213:30-34. doi: 10.1016/j.jad.2017.02.007
9. Marrie RA, Walld R, Bolton JM, et al. Increased incidence of psychiatric disorders in immune-mediated inflammatory disease. *J Psychosom Res* 2017;101:17-23. doi: 10.1016/j.jpsychores.2017.07.015
10. van der Horst-Bruinsma IE, Lems WF, Dijkmans BA. A systematic comparison of rheumatoid arthritis and ankylosing spondylitis. *Clin Exp Rheumatol* 2009;27(4 Suppl 55):S43-9.
11. Braun J, Sieper J. Treatment of rheumatoid arthritis and ankylosing spondylitis. *Clin Exp Rheumatol* 2009;27(4 Suppl 55):S146-7.
12. Cullen AE, Holmes S, Pollak TA, et al. Associations Between Non-neurological Autoimmune Disorders and Psychosis: A Meta-analysis. *Biol Psychiatry* 2019;85(1):35-48. doi: 10.1016/j.biopsych.2018.06.016.
13. Amador-Patarroyo MJ, Rodriguez-Rodriguez A, Montoya-Ortiz G. How does age at onset influence the outcome of autoimmune diseases? *Autoimmune Dis* 2012;2012:251730. doi: 10.1155/2012/251730.
14. Herrett E, Gallagher AM, Bhaskaran K, et al. Data Resource Profile: Clinical Practice Research Datalink (CPRD). *Int J Epidemiol* 2015;44(3):827-36. doi: 10.1093/ije/dyv098.
15. Dregan A, Moller H, Murray-Thomas T, et al. Validity of cancer diagnosis in a primary care database compared with linked cancer registrations in England. Population-based cohort study. *Cancer Epidemiol* 2012;36(5):425-9. doi: 10.1016/j.canep.2012.05.013

16. Dregan A, Toschke MA, Wolfe CD, et al. Utility of electronic patient records in primary care for stroke secondary prevention trials. *BMC Public Health* 2011;11:86. doi: 10.1186/1471-2458-11-86
17. Quint JK, Mullerova H, DiSantostefano RL, et al. Validation of chronic obstructive pulmonary disease recording in the Clinical Practice Research Datalink (CPRD-GOLD). *BMJ Open* 2014;4(7):e005540. doi: 10.1136/bmjopen-2014-005540
18. John A, McGregor J, Fone D, et al. Case-finding for common mental disorders of anxiety and depression in primary care: an external validation of routinely collected data. *BMC Med Inform Decis Mak* 2016;16:35. doi: 10.1186/s12911-016-0274-7
19. Thomas SL, Edwards CJ, Smeeth L, et al. How accurate are diagnoses for rheumatoid arthritis and juvenile idiopathic arthritis in the general practice research database? *Arthritis Rheum* 2008;59(9):1314-21. doi: 10.1002/art.24015
20. Lewis JD, Brensinger C, Bilker WB, et al. Validity and completeness of the General Practice Research Database for studies of inflammatory bowel disease. *Pharmacoepidemiol Drug Saf* 2002;11(3):211-8. doi: 10.1002/pds.698
21. Watts RA, Al-Taiar A, Scott DG, et al. Prevalence and incidence of Wegener's granulomatosis in the UK general practice research database. *Arthritis Rheum* 2009;61(10):1412-6. doi: 10.1002/art.24544
22. Schoonen WM, Kucera G, Coalson J, et al. Epidemiology of immune thrombocytopenic purpura in the General Practice Research Database. *Br J Haematol* 2009;145(2):235-44. doi: 10.1111/j.1365-2141.2009.07615.x
23. Charpentier C, Salleron J, Savoye G, et al. Natural history of elderly-onset inflammatory bowel disease: a population-based cohort study. *Gut* 2014;63(3):423-32. doi: 10.1136/gutjnl-2012-303864

24. Dregan A, Charlton J, Chowienczyk P, et al. Chronic inflammatory disorders and risk of type 2 diabetes mellitus, coronary heart disease, and stroke: a population-based cohort study. *Circulation* 2014;130(10):837-44. doi: 10.1161/CIRCULATIONAHA.114.009990
25. Yeung H, Takeshita J, Mehta NN, et al. Psoriasis severity and the prevalence of major medical comorbidity: a population-based study. *JAMA Dermatol* 2013;149(10):1173-9. doi: 10.1001/jamadermatol.2013.5015
26. Egeberg A, Hansen PR, Gislason GH, et al. Risk of self-harm and nonfatal suicide attempts, and completed suicide in patients with psoriasis: a population-based cohort study. *Br J Dermatol* 2016;175(3):493-500. doi: 10.1111/bjd.14633
27. Wu JJ, Penfold RB, Primatesta P, et al. The risk of depression, suicidal ideation and suicide attempt in patients with psoriasis, psoriatic arthritis or ankylosing spondylitis. *J Eur Acad Dermatol Venereol* 2017;31(7):1168-75. doi: 10.1111/jdv.14175
28. Kurd SK, Troxel AB, Crits-Christoph P, et al. The risk of depression, anxiety, and suicidality in patients with psoriasis: a population-based cohort study. *Arch Dermatol* 2010;146(8):891-5. doi: 10.1001/archdermatol.2010.186
29. Bhattarai N, Charlton J, Rudisill C, et al. Prevalence of depression and utilization of health care in single and multiple morbidity: a population-based cohort study. *Psychol Med* 2013;43(7):1423-31. doi: 10.1017/S0033291712002498
30. Ruscio AM, Chiu WT, Roy-Byrne P, et al. Broadening the definition of generalized anxiety disorder: effects on prevalence and associations with other disorders in the National Comorbidity Survey Replication. *J Anxiety Disord* 2007;21(5):662-76. doi: 10.1016/j.janxdis.2006.10.004
31. Warrington TP, Bostwick JM. Psychiatric adverse effects of corticosteroids. *Mayo Clin Proc* 2006;81(10):1361-7. doi: 10.4065/81.10.1361

32. Gallagher PJ, Castro V, Fava M, et al. Antidepressant response in patients with major depression exposed to NSAIDs: a pharmacovigilance study. *Am J Psychiatry* 2012;169(10):1065-72. doi: 10.1176/appi.ajp.2012.11091325
33. Nakasian SS, Rassen JA, Franklin JM. Effects of expanding the look-back period to all available data in the assessment of covariates. *Pharmacoepidemiol Drug Saf* 2017;26(8):890-99. doi: 10.1002/pds.4210
34. Lindor KD, Dickson ER, Baldus WP, et al. Ursodeoxycholic acid in the treatment of primary biliary cirrhosis. *Gastroenterology* 1994;106(5):1284-90.
35. Austin PC. The performance of different propensity score methods for estimating marginal hazard ratios. *Stat Med* 2013;32(16):2837-49. doi: 10.1002/sim.5705
36. Cummings P, McKnight B, Greenland S. Matched cohort methods for injury research. *Epidemiol Rev* 2003;25:43-50.
37. Schmidt FL, Oh IS, Hayes TL. Fixed- versus random-effects models in meta-analysis: model properties and an empirical comparison of differences in results. *Br J Math Stat Psychol* 2009;62(Pt 1):97-128. doi: 10.1348/000711007X255327
38. Dean LE, Macfarlane GJ, Jones GT. Differences in the prevalence of ankylosing spondylitis in primary and secondary care: only one-third of patients are managed in rheumatology. *Rheumatology (Oxford)* 2016;55(10):1820-5. doi: 10.1093/rheumatology/kew228
39. Verstappen SM, Lunt M, Bunn DK, et al. In patients with early inflammatory polyarthritis, ACPA positivity, younger age and inefficacy of the first non-biological DMARD are predictors for receiving biological therapy: results from the Norfolk Arthritis Register. *Ann Rheum Dis* 2011;70(8):1428-32. doi: 10.1136/ard.2010.148106

40. Gudjonsson JE, Karason A, Runarsdottir EH, et al. Distinct clinical differences between HLA-Cw*0602 positive and negative psoriasis patients--an analysis of 1019 HLA-C- and HLA-B-typed patients. *J Invest Dermatol* 2006;126(4):740-5. doi: 10.1038/sj.jid.5700118
41. Dickens C, Creed F. The burden of depression in patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2001;40(12):1327-30. [published Online First: 2001/12/26]
42. Ananthakrishnan AN, Gainer VS, Cai T, et al. Similar risk of depression and anxiety following surgery or hospitalization for Crohn's disease and ulcerative colitis. *Am J Gastroenterol* 2013;108(4):594-601. doi: 10.1038/ajg.2012.471
43. Marrie RA, Hitchon CA, Walld R, et al. Increased Burden of Psychiatric Disorders in Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)* 2018 doi: 10.1002/acr.23539
44. Meesters JJ, Bremander A, Bergman S, et al. The risk for depression in patients with ankylosing spondylitis: a population-based cohort study. *Arthritis Res Ther* 2014;16(5):418. doi: 10.1186/s13075-014-0418-z]
45. Ettner SL, Azocar F, Branstrom RB, et al. Association of general medical and psychiatric comorbidities with receipt of guideline-concordant care for depression. *Psychiatr Serv* 2010;61(12):1255-9. doi: 10.1176/ps.2010.61.12.1255
46. Curtis HJ, Dennis JM, Shields BM, et al. Time trends and geographical variation in prescribing of drugs for diabetes in England from 1998 to 2017. *Diabetes Obes Metab* 2018;20(9):2159-68. doi: 10.1111/dom.13346
47. Austin PC. Comparing paired vs non-paired statistical methods of analyses when making inferences about absolute risk reductions in propensity-score matched samples. *Stat Med* 2011;30(11):1292-301. doi: 10.1002/sim.4200
48. Hill HF, Hill AG, Bodmer JG. Clinical diagnosis of ankylosing spondylitis in women and relation to presence of HLA-B27. *Ann Rheum Dis* 1976;35(3):267-70.

49. Rothman KJ, Gallacher JE, Hatch EE. Why representativeness should be avoided. *Int J Epidemiol* 2013;42(4):1012-4. doi: 10.1093/ije/dys223
50. Richiardi L, Pizzi C, Pearce N. Commentary: Representativeness is usually not necessary and often should be avoided. *Int J Epidemiol* 2013;42(4):1018-22. doi: 10.1093/ije/dyt103

Table 1*. Participants' characteristics at baseline assessment. Figures are numbers and percentages unless otherwise specified.

	RA	Psoriasis		CD	UC	SLE	Vasculitis	AS	Total	
		Mild	Severe						Exposed	Unexposed
N	37399	84184	6528	10453	23291	3604	14177	10363	180163	358544
Follow-up±	4(2,8)	5(2,8)	6(3,9)	4(2,8)	4(2,8)	4(2,8)	4(2,7)	5(2,9)	4(2,8)	4(2,7)
Age – M(sd)	60(16)	49(18)	49(16)	46(18)	54(19)	51(16)	65(17)	51(17)	53(18)	53(18)
Female	24929(67)	43757(52)	3474(53)	5678(54)	12664(54)	2936(81)	8974(63)	6301(61)	108713(57)	216287(57)
Cancer	3164(8)	4319(5)	301(5)	525(5)	1969(8)	261(7)	1673(12)	609(6)	12821(7)	24050(6)
CKD	4875(13)	5936(7)	529(8)	895(9)	2836(12)	436(12)	2574(18)	927(9)	19008(10)	28645(8)
Diabetes	4224(11)	6349(8)	564(9)	694(7)	2297(10)	259(7)	1934(12)	751(7)	17072(9)	26822(7)
CHD	3985(11)	5415(6)	403(6)	688(7)	2345(10)	242(7)	2082(15)	753(7)	15913(8)	25606(7)
COPD	2131(6)	2532(3)	200(3)	348(3)	1110(5)	105(3)	935(7)	269(3)	7630(4)	10428(3)
Stress	8006(21)	14598(17)	1239(19)	1880(18)	4505(19)	832(23)	2917(21)	2244(22)	36221(19)	57083(15)
Hypertension	14177(39)	25138(33)	1993(33)	2402(25)	6945(31)	1063(30)	6472(47)	3071(31)	61261(34)	112226(34)
Obesity	9710(29)	18926(27)	1837(34)	1792(20)	4339(21)	714(23)	3284(26)	2122(23)	42724(26)	69536(23)
Smoking	27738(76)	55278(69)	4364(70)	7236(82)	17927(80)	2419(69)	11124(81)	7365(74)	133451(73)	262379(76)
Alcohol	8060(24)	12930(19)	1103(20)	1826(21)	4195(22)	827(26)	3254(26)	1783(20)	33924(21)	62263(21)
AHT	19205(51)	28043(33)	2422(37)	3337(32)	10013(43)	1553(43)	8520(60)	4004(39)	77097(41)	128197(34)
Statins	9953(27)	13820(16)	1161(18)	1493(14)	5220(22)	621(17)	4585(32)	1749(17)	38602(20)	64919(17)
Hypnotics	5985(16)	9492(11)	934(14)	1366(13)	3361(14)	559(16)	2407(17)	1592(15)	25696(14)	37263(10)
NSAIDs	31894(85)	46984(56)	4477(69)	6089(58)	14115(61)	2513(70)	10080(71)	8571(83)	142723(66)	181991(48)
Steroids	16467(44)	14390(17)	1893(29)	3499(33)	7022(30)	1287(36)	8444(60)	2490(24)	55492(29)	54104(14)

Note: * - For ease of presentation some of the covariates data are not presented here. ± figures represent median and interquartile range. M=mean; sd=standard deviation; CD= Crohns disease, UC= ulcerative colitis, AS=ankylosing spondylitis, RA= rheumatoid arthritis, SLE=systemic lupus erythematosus. AHT=antihypertensive therapy; DMARDs= disease modifying anti-rheumatic drugs; NSADs= non-steroidal anti-inflammatory drugs; CKD= chronic kidney disease; COPD= chronic obstructive pulmonary disease; CHD= coronary heart disease.

Table 2. Adjusted hazard ratios (95% confidence interval) for depression and anxiety incidence among persons inflammatory disorders diagnosis compared to the matched comparison group.

	Overall sample HR(95% CI)	Age at diagnosis			
		<40 HR(95% CI)	40-49 HR(95% CI)	50-59 HR(95% CI)	≥60 HR(95% CI)
<i>Depression incidence</i>					
Psoriasis mild	1.30(1.26-1.35)	1.59(1.48-1.71)	1.32(1.20-1.45)	1.01(0.92-1.12)	0.88(0.81-0.95)
Psoriasis severe	1.71(1.52-1.93)	2.00(1.61-2.48)	1.77(1.39-2.24)	1.21(0.94-1.57)	0.87(0.66-1.13)
Rheumatoid arthritis	1.38(1.29-1.47)	2.40(2.07-2.79)	1.93(1.68-2.22)	1.40(1.23-1.59)	1.06(0.96-1.17)
Systemic lupus erythematosus	1.28(1.06-1.56)	1.27(0.91-1.78)	1.53(1.09-2.14)	1.02(0.68-1.54)	0.91(0.65-1.28)
Ankylosing spondylitis	1.44(1.30-1.60)	1.93(1.59-2.33)	1.62(1.30-2.01)	1.30(1.02-1.65)	1.07(0.87-1.30)
Systemic vasculitis	1.46(1.31-1.62)	2.52(1.98-3.20)	2.37(1.83-3.09)	1.73(1.37-2.20)	1.23(1.07-1.42)
Ulcerative colitis	1.39(1.29-1.49)	1.81(1.60-2.05)	1.31(1.09-1.56)	1.44(1.22-1.70)	0.96(0.84-1.09)
Crohn's disease	1.47(1.32-1.63)	1.84(1.55-2.19)	1.59(1.26-2.00)	1.28(0.99-1.65)	0.92(0.73-1.14)
<i>Anxiety incidence</i>					
Psoriasis mild	1.28(1.24-1.33)	1.51(1.40-1.63)	1.08(0.97-1.21)	1.03(0.93-1.15)	0.85(0.78-0.93)
Psoriasis severe	1.33(1.17-1.50)	1.40(1.10-1.80)	1.31(0.98-1.75)	1.04(0.77-1.41)	0.84(0.63-1.14)
Rheumatoid arthritis	1.10(1.03-1.18)	1.51(1.26-1.81)	1.20(1.01-1.43)	0.93(0.79-1.09)	0.80(0.71-0.90)
Systemic lupus erythematosus	1.28(1.06-1.55)	1.25(0.78-1.83)	1.61(1.07-2.42)	1.03(0.63-1.67)	0.78(0.49-1.22)
Ankylosing spondylitis	1.36(1.23-1.51)	1.54(1.25-1.90)	1.33(1.05-1.70)	1.32(1.03-1.70)	1.08(0.88-1.32)
Systemic vasculitis	1.19(1.07-1.32)	1.52(1.15-2.02)	1.45(1.04-2.02)	1.24(0.94-1.64)	1.01(0.86-1.18)
Ulcerative colitis	1.34(1.24-1.44)	1.57(1.35-1.83)	1.21(0.99-1.48)	0.94(0.76-1.17)	0.86(0.74-1.00)
Crohn's disease	1.35(1.21-1.50)	1.24(1.02-1.49)	0.97(0.73-1.28)	1.03(0.76-1.39)	0.78(0.61-1.00)

HR- hazard rate; CI- confidence interval. Adjusted for age, gender, deprivation, BP, BMI, smoking, alcohol, CHD, stroke, cancer, diabetes, cancer, dementia, epilepsy, CKD, liver disease, COPD, sleep disorders, AHT, statins, hypnotics, corticosteroids, NSAIDs, anti-diabetics.




Figure 1 Incidence of depression by condition for participants with chronic inflammatory disorders and matched controls.




Figure 2. Forest plot displaying random-effects meta-analysis of the influence of specific inflammatory disorders on the incidence of multiple depression and anxiety outcomes. CI indicates confidence interval; HR, hazard rate. Basic - adjusted for age and gender. Adjusted – adjusted for age, gender, deprivation, BP, BMI, smoking, alcohol, CHD, stroke, cancer, diabetes, cancer, dementia, epilepsy, CKD, liver disease, COPD, sleep disorders, AHT, statins, hypnotics, corticosteroids, NSAIDs, anti-diabetics.




Figure 3 Forest plot displaying random-effects meta-analysis of the influence of age at inflammatory disorders onset on the incidence of multiple depression and anxiety. CI indicates confidence interval; HR, hazard rate. Adjusted for age, gender, deprivation, BP, BMI, smoking, alcohol, CHD, stroke, cancer, diabetes, cancer, dementia, epilepsy, CKD, liver disease, COPD, sleep disorders, AHT, statins, hypnotics, corticosteroids, NSAIDs, anti-diabetics.