



King's Research Portal

Document Version
Other version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Grojnowski, I., & Shepherd-Barron, N. I. (2019). *Del Pezzo surfaces as Springer fibres for exceptional groups*. arXiv. <https://arxiv.org/abs/1507.01872>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

DEL PEZZO SURFACES AS SPRINGER FIBRES FOR EXCEPTIONAL GROUPS

I. GROJNOWSKI AND N. I. SHEPHERD-BARRON

1 Introduction

Since the discovery by Cayley and Salmon of the 27 lines on a cubic surface and their configuration and that by Killing and Cartan of the exceptional simple Lie algebra of type E_6 it has been clear that del Pezzo surfaces and exceptional simple algebraic groups have attached to them the same combinatorial objects. That is, there are well known constructions

$$(\text{del Pezzo surfaces}) \rightarrow (\text{exceptional root data}) \leftrightarrow (\text{exceptional simple groups}).$$

One of the results of this paper is a direct geometrical construction, for $5 \leq l \leq 8$, of the del Pezzo surfaces S of degree $d = 9 - l$ from the split simply connected exceptional simple¹ group G of type E_l , in the presence of the universal elliptic curve \mathcal{E} . Direct, that is, in the sense that it does not involve root data.

The del Pezzo surfaces arise out of the group as follows. Let P denote the weight lattice of G and B a Borel subgroup. Suppose that $\tilde{\mathcal{G}}_{\mathcal{E}}^{ss}$ is the stack that classifies semi-stable G -bundles over elliptic curves together with a reduction of degree 0 of the structure group to B . (The existence of such a reduction is a characterization of semi-stability.) Then there is a partial relative compactification of $\tilde{\mathcal{G}}_{\mathcal{E}}^{ss}$ in which the total space \mathcal{D}_1^- of a certain family $\mathcal{D}_1^- \rightarrow \mathcal{Y} = \text{Hom}(P, \mathcal{E})$ of these del Pezzo surfaces appears as a boundary divisor. We shall describe this in more detail later in this introduction. (The del Pezzo surfaces in the family $\mathcal{D}_1^- \rightarrow \mathcal{Y}$ are in fact *marked*; that is, a copy of \mathcal{E} is embedded in them as an anticanonical curve and their Néron–Severi groups are rigidified. The family is nearly, but not quite, universal; the fine moduli space is a non-separated union of copies of \mathcal{Y} , and \mathcal{Y} is then its maximal separated quotient.)

However, we have no suggestion for a similarly direct construction of the groups from the del Pezzo surfaces that would bypass consideration of the root data.

The title of this paper refers to our presentation of these results as an extension of the construction and theorems due to Brieskorn, Grothendieck, Slodowy and Springer (BGSS) and described in detail in [Sl] that, in good characteristics, reveal the deformations and simultaneous resolution of du Val singularities (= rational

¹We write $E_5 = D_5$ and regard D_5 as exceptional. We also refer to groups as “simple” rather than “almost simple”.

double points = simple singularities = Kleinian singularities) of type A , D or E inside the corresponding split simply connected simple algebraic group. In turn, this provides a characterization of du Val singularities. Of course, there are many others. One, that we shall use, is that a surface singularity is du Val if and only if its monodromy group, acting on the vanishing cohomology, is the relevant Weyl group.

Our results lead, as a by-product, to an identification of the unipotent singularity of E_8 in bad characteristics (that is, 2, 3 and 5). This is done in section 8. The proof depends upon specializing \mathcal{E} to a uniformizable elliptic curve E over a complete algebraically closed field of characteristic p . The picture is that the curve E acts as a vector to carry some of the geometry of the group into a del Pezzo surface S with a singularity isomorphic to the unipotent singularity, and we can then exploit the fact that the defining equations of such a surface are so simple that classification of the unipotent singularities becomes very easy and quick. Moreover, this leads to an embedding of the group-theoretical description of the deformations and resolutions of du Val singularities into their description in terms of del Pezzo surfaces, as expounded in [SSS].

This is, perhaps, some slight evidence supporting a view that our results tell us something about the group, and that further information about the group is to be found in higher codimension in suitable relative compactifications of $\mathcal{G}_{\mathcal{E}}^{ss}$.

In the rest of this introduction we give a more detailed survey of our results.

From the viewpoint of the geometry of algebraic surfaces and their degenerations the extension referred to above consists of showing that *simultaneous log resolutions*, which are defined in the next paragraph, of simply elliptic singularities of degrees $d = 1, 2, 3$ or 4 , which were first observed in the context of type II degenerations of K3 surfaces, although inspired by work of Brieskorn [B] and Looijenga [L], also have realizations in terms of simple algebraic groups. Here, only exceptional groups of type E_{9-d} can, and do, occur.

We define a *del Pezzo surface* to be a surface S whose anti-canonical divisor class is ample and that has at worst du Val singularities; a *weak del Pezzo surface* is smooth and its anti-canonical class is ample modulo finitely many (-2) -curves. That is, a weak del Pezzo surface is the minimal resolution of a del Pezzo surface and a del Pezzo surface is the anticanonical model of a weak del Pezzo surface.

Definition 1.1 *Given a flat family $X \rightarrow S$ of normal Gorenstein surfaces, a simultaneous log resolution of $X \rightarrow S$ is a commutative diagram*

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & X \\ \downarrow & & \downarrow \\ \tilde{S} & \longrightarrow & S \end{array}$$

where $\tilde{S} \rightarrow S$ is proper, dominant and generically finite, $\tilde{X} \rightarrow X \times_S \tilde{S}$ is proper and birational, \tilde{X} is smooth, $\tilde{X} \rightarrow \tilde{S}$ is flat and semi-stable, in the sense that

each geometric fibre is a reduced union of smooth surfaces with normal crossings, and the relative canonical class $K_{\tilde{X}/\tilde{S}}$ is equivalent to the pullback of $K_{X/S}$.

It is well known in the context of the birational geometry of degenerating families of K3 surfaces that a versal deformation $X \rightarrow S$ of a simply elliptic singularity of degree at most 4 possesses a simultaneous log resolution $\tilde{X} \rightarrow \tilde{S}$ with the following properties.

- (1) $\tilde{X} \rightarrow \tilde{S}$ is of type II, in the sense that, for each fibre X_s with an elliptic singularity, its inverse image \tilde{X}_s in \tilde{X} has two components, say $\tilde{X}_s = D_1 + D_2$. Moreover D_1 is the minimal resolution of X_s , D_2 is a weak del Pezzo surface, D_2 is contracted to a point under $\tilde{X} \rightarrow X$ and $D_1 \cap D_2$ is the elliptic curve that is the exceptional locus of $D_1 \rightarrow X_s$.
- (2) For each fibre X_s that has at most du Val singularities, its inverse image \tilde{X}_s in \tilde{X} is the minimal resolution of X_s .
- (3) the base change $\tilde{S} \rightarrow S$ is the composite of a ramified Galois covering of the base S whose Galois group is the corresponding finite Weyl group and a blow-up along the simply elliptic locus in S .

On the other hand, Helmke and Slodowy [HS1] and [HS3] have given a brief indication of a proof that miniversal deformations of simply elliptic singularities of degree d can be realized inside \mathcal{G} when $G = E_{9-d}$.

Recall from [HS2], Theorems 5.6 and 5.12, that a *regular* bundle on an elliptic curve E over a field is one whose automorphism group has minimal dimension, namely l , while for a *subregular* bundle the automorphism group has the next smallest dimension, $l + 2$. Up to translation by points of E there is a unique unstable regular bundle η and a unique unstable subregular bundle ξ ; in a miniversal deformation of ξ the unstable locus is, up to this translation, a surface S and all points of $S - \{\xi\}$ correspond to bundles isomorphic to η . Note that in order to construct these bundles it is necessary (and sufficient) for the group to be split and the curve E to have a point; it cannot be replaced by an arbitrary curve of genus 1. Then both ξ and η are constructed as bundles on \mathcal{E} .

Let's list the objects that appear in our main theorem. These are:

- (1) the stack $\mathcal{G} = \mathcal{G}_{\mathcal{E}}$ of principal G -bundles over the universal elliptic curve \mathcal{E} (defined over the moduli stack \mathcal{M}_{Ell} of elliptic curves) and the open substack \mathcal{G}^{ss} of semi-stable bundles;
- (2) the stack $\tilde{\mathcal{G}}^{ss}$ that classifies semi-stable G -bundles together with a reduction of the structure group to a Borel subgroup such that the associated torus bundle is of degree 0 (in the course of reaching our main result we shall show that $\tilde{\mathcal{G}}^{ss}$ is proper, representable and generically finite, of degree equal to the order of the Weyl group W of G , over \mathcal{G}^{ss} , and that, when we restrict attention to a uniformizable elliptic curve, this structure reproduces the BGSS picture);
- (3) stacks $\tilde{\mathcal{G}}$ and $\tilde{\mathcal{G}}^+$ (to be described in Section 5), a projective birational morphism $\tilde{\mathcal{G}} \rightarrow \tilde{\mathcal{G}}^+$ and a morphism which is proper but not representable (although its fibres have finite automorphism groups) $\tilde{\mathcal{G}}^+ \rightarrow \mathcal{G}$ such that $\tilde{\mathcal{G}} \rightarrow \mathcal{G}$ and $\tilde{\mathcal{G}}^+ \rightarrow \mathcal{G}$

are relative compactifications of $\tilde{\mathcal{G}}^{ss} \rightarrow \mathcal{G}^{ss}$;

(4) the abelian variety $\mathcal{Y} = \text{Hom}(P, \mathcal{E})$ over \mathcal{M}_{EU} and a certain ample W -linearized line bundle $\mathcal{L} \rightarrow \mathcal{Y}$, which gives a cone $\hat{\mathcal{Y}}$ over \mathcal{M}_{EU} by contracting the zero section $0_{\mathcal{L}}$ of \mathcal{L} such that the geometric quotient $[\hat{\mathcal{Y}}/W]$ of $\hat{\mathcal{Y}}$ by W , relative to \mathcal{M}_{EU} , is, by [L], an affine space bundle of rank $l + 1$ over \mathcal{M}_{EU} ;

(5) the unstable subregular bundle ξ mentioned above, that is defined over \mathcal{M}_{EU} and is unique modulo translation by \mathcal{E} ;

(6) the base \mathcal{Z} of a deformation that is minimally versal, modulo the translation by \mathcal{E} mentioned above, of ξ ;

(7) the products $\tilde{\mathcal{Z}} = \mathcal{Z} \times_{\mathcal{G}} \tilde{\mathcal{G}}$, and $\tilde{\mathcal{Z}}^+ = \mathcal{Z} \times_{\mathcal{G}} \tilde{\mathcal{G}}^+$ and a birational map (not a morphism) $\tilde{\mathcal{Z}}^{+-} \rightarrow \tilde{\mathcal{Z}}^-$ which is constructed as a succession of flops relative to \mathcal{Z} such that the morphisms $\tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$ and $\tilde{\mathcal{Z}}^{\pm} \rightarrow \mathcal{Z}$ are representable.

Moreover, let ${}^0\mathcal{G}$ be the open substack of \mathcal{G} that is the complement of the locus of those unstable bundles Ξ such that $\dim \text{Aut}(\Xi) > l + 4$. That is, the locus of unstable bundles that are “worse than subregular” has been deleted. In particular, \mathcal{G}^{ss} is contained in ${}^0\mathcal{G}$. Set ${}^0\tilde{\mathcal{G}} = \tilde{\mathcal{G}} \times_{\mathcal{G}} {}^0\mathcal{G}$ and ${}^0\tilde{\mathcal{G}}^+ = \tilde{\mathcal{G}}^+ \times_{\mathcal{G}} {}^0\mathcal{G}$, so that \mathcal{Z} is, up to turning off $\text{Pic}^1(\mathcal{E})$, a chart in ${}^0\mathcal{G}$ and $\tilde{\mathcal{Z}}$ is a chart in ${}^0\tilde{\mathcal{G}}$. Then the birational morphism $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{Z}}^+$ and the birational maps $\tilde{\mathcal{Z}}^{+-} \rightarrow \tilde{\mathcal{Z}}^-$ and $\tilde{\mathcal{Z}}^- \rightarrow \tilde{\mathcal{Z}}^-$ are all isomorphisms over \mathcal{G}^{ss} ; that is, all birational modifications are made inside the boundary of ${}^0\tilde{\mathcal{G}}$ and the commutative square

$$\begin{array}{ccc} \tilde{\mathcal{Z}} & \longrightarrow & \tilde{\mathcal{Z}}^+ \\ \downarrow & \swarrow & \downarrow \\ \tilde{\mathcal{Z}}^- & \longrightarrow & \mathcal{Z} \end{array}$$

of rational maps and morphisms is a local description of a commutative square

$$\begin{array}{ccc} {}^0\tilde{\mathcal{G}} & \longrightarrow & {}^0\tilde{\mathcal{G}}^+ \\ \downarrow & \swarrow & \downarrow \\ {}^0\tilde{\mathcal{G}}^- & \longrightarrow & {}^0\mathcal{G} \end{array}$$

of rational maps and morphisms between stacks.

What is missing from this picture is a suitable stack $\tilde{\mathcal{G}}^-$ of which ${}^0\tilde{\mathcal{G}}^-$ is a natural open substack.

The result of Helmke and Slodowy concerning elliptic singularities that we mentioned above is that there is a flat morphism $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$ that is a minimally versal deformation of a simply elliptic singularity of degree d . We give a detailed proof of this as part of our main result, which is that there is also an analogue of the BGSS construction for the situation involving \mathcal{E} , as in the following theorem, which summarizes the results of the paper. The main result is given more concisely in Theorem 7.12.

Theorem 1.2 *Suppose that $G = E_{9-d}$.*

(1) *Let $\tilde{\mathcal{Z}}^\sharp$ denote any one of $\tilde{\mathcal{Z}}, \tilde{\mathcal{Z}}^\pm$. Then there is a commutative diagram*

$$\begin{array}{ccc} \tilde{\mathcal{Z}}^\sharp & \longrightarrow & \mathcal{Z} \\ \downarrow \pi^\sharp & & \downarrow \\ \mathcal{L} & \longrightarrow & [\hat{\mathcal{Y}}/W] \end{array}$$

where $\pi^\sharp : \tilde{\mathcal{Z}}^\sharp \rightarrow \mathcal{L}$ is semi-stable and smooth over $\mathcal{L} - 0_{\mathcal{L}}$, the complement of the zero section $0_{\mathcal{L}}$.

(2) *Suppose that 0 is a point of $0_{\mathcal{L}}$ lying over the origin $0_{\mathcal{Y}}$ of \mathcal{Y} . Then $\tilde{\mathcal{Z}}_0 = D_0 + D_1 + Q$, $\tilde{\mathcal{Z}}_0^\pm = D_0^\pm + D_1^\pm$, each $D_0^\sharp \cap D_1^\sharp$ is a copy of \mathcal{E} , \mathcal{Z}_0 is a cone over \mathcal{E} of degree d and $\tilde{\mathcal{Z}}^-$ is its minimal resolution. Q is a copy of $\mathbb{P}^1 \times \mathbb{P}^1$ and the morphism $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{Z}}^+$ contracts Q to a curve via a projection $Q \rightarrow \mathbb{P}^1$.*

(3) *When $\tilde{\mathcal{Z}}^\sharp = \tilde{\mathcal{Z}}^-$ then the diagram is a type II simultaneous log resolution of $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$.*

(4) *The rational map $\tilde{\mathcal{Z}}^+ \dashrightarrow \tilde{\mathcal{Z}}^-$ is a sequence of flops and the centre of each flop is the total space of a smooth family of rational curves over $0_{\mathcal{L}}$.*

(5) *$\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$ is a miniversal deformation of a simply elliptic singularity of degree d over $\mathbf{Spec} \mathbb{Q}$, and over $\mathbf{Spec} \mathbb{Z}[1/d]$ if $d \leq 3$.*

(6) *The morphisms $\tilde{\mathcal{Z}}^\sharp \rightarrow \mathcal{Z} \times_{[\hat{\mathcal{Y}}/W]} \mathcal{L}$ are birational.*

(7) *The exceptional locus \mathcal{D}_1^- in \mathcal{Z}^- is the total space of a family $\mathcal{D}_1^- \rightarrow \mathcal{Y} = 0_{\mathcal{L}}$ of weak del Pezzo surfaces of degree d . In particular, the fibre over a point (y, ξ) of the morphism $\mathcal{Z}^- \rightarrow \mathcal{Z} \times_{[\hat{\mathcal{Y}}/W]} \mathcal{L}$, where y is a point in the zero section $0_{\mathcal{L}}$ of \mathcal{L} , is a weak del Pezzo surface of degree d .*

(8) *Locally on \mathcal{M}_{Ell} there is an isomorphism $\tilde{\mathcal{Z}}^- \rightarrow \omega_{\mathcal{D}_1^-/\mathcal{Y}}$.*

The fibres of 1.2 7 are the Springer fibres of the title.


Figure 1 below shows the structure of the fibres $\tilde{\mathcal{Z}}_0, \tilde{\mathcal{Z}}_0^\pm$ and \mathcal{Z}_0 .

The crucial part of the proof is Theorem 6.7, where we exhibit a surface D_1 in $\tilde{\mathcal{G}}_{\mathcal{E}}$ whose transform under a natural flop is a weak del Pezzo surface. It follows (this is Corollary 6.18) that that the fibre of $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$ over the origin of the affine space $[\hat{\mathcal{Y}}/W]$ has a simply elliptic singularity. We then recover the result of Helmke and Slodowy, that $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$ is a miniversal deformation of this singularity. In some sense, this is a ‘‘top down’’ approach which depends upon being able to recognize a singularity from its resolution.

Remark: In particular, we have made a non-trivial birational modification of $\tilde{\mathcal{G}}_{\mathcal{E}}$, centred on a locus in the boundary $\tilde{\mathcal{G}}_{\mathcal{E}} - \tilde{\mathcal{G}}_{\mathcal{E}}^{ss}$, which is of a kind that is dictated by the needs of birational geometry but which has no obvious modular interpretation, in order to arrive at $\tilde{\mathcal{Z}}^-$.

However, this is a common phenomenon.

For example, consider the coarse moduli space A_g of principally polarized abelian varieties over \mathbb{C} and its perfect compactification A_g^P . Then [SB16] the

Figure 1: The fibres $\tilde{\mathcal{Z}}^\#$ and \mathcal{Z}_0

exceptional locus of the \mathbb{Q} -divisor $12M - D$, where M is the bundle of modular forms of weight 1 and D is the boundary, is the image of $A_1^P \times A_{g-1}^P$ in A_g^P , the locus of ppav's with an elliptic factor. So, if $7 \leq g \leq 10$, then A_g^P has terminal singularities [AS16] and is of general type, while $K_{A_g^P} \sim (g+1)M - D$ is not nef. Therefore the first step in running the Minimal Model Program on A_g^P is to contract and flip the extremal ray generated by the curve $A_1^P \times \{x\}$, where x is a point A_{g-1}^P . This is a non-trivial birational modification which is dictated by the MMP and which is centred in a locus that penetrates the interior A_g of A_g^P that has no obvious modular interpretation, unless the notion of ppav can be modified.

It is important to point out that, beyond the work of Brieskorn and Looijenga mentioned above, and that of Pinkham [P], what we do builds very much on the results of Helmke and Slodowy, and, to an equal extent, on those of Friedman and Morgan [FM].

We will revisit the results of this paper in [GS], where amongst other things we will analyse the moduli stacks in a neighbourhood of infinity by using the Tate curve, give uniform descriptions of the combinatorics in terms of the affine Weyl groups and extend the main theorem of this paper to the cases where $5 \leq d \leq 9$.

Finally, if H is a group then $\mathbb{B}H$ denotes its classifying stack and $[\mathcal{X}]$ denotes the geometric quotient of an algebraic stack \mathcal{X} when that quotient exists.

Acknowledgements

We are very grateful to Dougal Davis for many conversations on these matters and particularly for pointing out an error in our proof of the previous version of Proposition 6.4. We are also grateful to Igor Dolgachev for his question about unipotent singularities and to Simon Salamon for many kindnesses, which include drawing the picture.

2 The BGSS construction

To set the notation we begin by recalling a little of the machinery of reductive groups in a way that emphasizes the flag variety, and its intersection theory, of an affine group rather than the group itself. This approach is very well known, and details can be found in, for example, [MOSW].

As the starting point of the discussion, fix a smooth connected reductive² algebraic group G over a field k . Then [SGA3 XXII, Th. 5.8.1 and Cor. 5.8.3, pp. 228-230] there is an associated *flag variety* F over k that parametrizes the maximal soluble subalgebras (the Borel subalgebras) of $\mathfrak{g} = \text{Lie } G$; it is constructed as a subscheme of the Grassmannian $Gr(\mathfrak{g})$. It is a smooth projective k -variety and is homogeneous under the obvious G -action; it is maximal among the set of homogeneous projective k -varieties the stabilizers of whose points are smooth, and the stabilizer of each point of F is a Borel subgroup (that is, a maximal connected reduced soluble subgroup of G). If G has a Borel subgroup B defined over K then $F = G/B$, but we do not yet assume that such a subgroup exists; that is, we do not assume that F has a k -point. By definition, this last condition is equivalent to G being *quasi-split*.

Locally in the étale topology on $\mathbf{Spec} k$ there is an array of projective homogeneous G -varieties, 2^l in number where l is the semi-simple rank of G , and G -morphisms between them; in particular, there are G -maps $F \rightarrow F_1, \dots, F \rightarrow F_l$ that are étale \mathbb{P}^1 -bundles and there are G -maps $F \rightarrow X_i$ where X_i is *minimal*, that is, of Picard number 1. We shall assume that all these varieties, and the morphisms in the lattice, are defined over k ; this assumption is fulfilled automatically if the Dynkin diagram attached to G has no symmetries. Indeed, the Galois group that acts on the array of varieties above is a subgroup of the symmetry group of the Dynkin diagram. If G is quasi-split, this is, by definition, the same as G being *split*.

For each minimal variety X_i there is a unique line bundle ϖ_i on $X_i \otimes \bar{k}$ that is a positive generator of $\text{Pic}_{X_i \otimes \bar{k}}$; we assume that ϖ_i is defined over K . This condition holds automatically if G is split. We also let ϖ_i denote the pullback of ϖ_i to F ; then $\varpi_1, \dots, \varpi_l$ are the *fundamental dominant weights* of G and form

²This assumption was omitted in a previous version of this paper. Otherwise, as was pointed out by a referee, the field k must be assumed to be perfect.

a \mathbb{Z} -basis of Pic_F . (So, for groups of type E_8 the bundles ϖ_i are always defined over k , because here the root lattice equals the weight lattice.) They also span the nef cone of F , which is also the effective cone of F . The semi-simple rank of G is the Picard number of F .

Set $\alpha_j = T_{F/F_j}$, the relative tangent bundle. This is a G -linearized line bundle, so defines a class in Pic_F^G , the group of classes of G -linearized line bundles on F . (The rank of this group is the rank of G .) These classes are the *simple roots* of G . The *simple coroots* are $\alpha_1^\vee, \dots, \alpha_l^\vee$ where α_i^\vee is the class of a fibre of $F \rightarrow F_i$; such a curve is a conic (a smooth curve of genus zero).

So some of the various intersection numbers between these curves and divisor classes are given by $(\alpha_i^\vee, \varpi_j) = \delta_{ij}$ and $(\alpha_i^\vee, \alpha_i) = 2$.

The numbers $1, \dots, l$ are the nodes of the associated Coxeter–Dynkin diagram D ; the edges of D are labelled in a way that is determined by the numbers $(\alpha_i^\vee, \alpha_j)$. This diagram leads to a Coxeter system $(W, S = \{s_1, \dots, s_l\})$ acting on Pic_F^G in the usual way: $s_i(x) = x + (x, \alpha_i^\vee)\alpha_i$. There is also a root datum $(M, M^\vee, \Delta, \Delta^\vee)$ derived from this picture: $M = \text{Pic}_F^G$, Δ is the W -orbit of the set of simple roots and Δ^\vee is the W -orbit of the set of simple coroots. More intrinsically, the roots arise as the G -linearized line bundles in a natural filtration of the tangent bundle T_F and the coroots appear as the classes of curves in the Bruhat decomposition of F . Conversely, each root datum leads to a unique split reductive group defined over \mathbb{Z} . (A smooth affine group G over a base S is reductive if all its geometric fibres are reductive.)

Now assume that G is quasi-split, so split. Fix a k -point of F and rigidify all line bundles and G -linearized line bundles on F at this point. Define the *weight lattice* P of G by $P = \text{Pic}_F^G$. Then there are universal torsors $\mathcal{T} \rightarrow F$ and $\mathcal{T}^G \rightarrow F$ under tori T and T^G , respectively, whose character groups are $\mathbb{X}^*(T) = \text{Pic}_F$ and $\mathbb{X}^*(T^G) = P$. Since G is reductive it acts effectively on \mathcal{T}^G .

If the forgetful homomorphism $\text{Pic}_F^G \rightarrow \text{Pic}_F$ is injective then G is *semi-simple* and if it is an isomorphism then G is *simply connected*. In this case the maps $F \rightarrow F_i$ are Zariski \mathbb{P}^1 -bundles. A semi-simple group is *simple* if its Dynkin diagram is connected.

Remark: It is clear from this that in fact a reductive group does not give rise only to a root datum, but rather to a *pinned root datum* [SGA3 XXIII, 1.5]. On the other hand, a del Pezzo surface S gives rise to a root datum (M, M^*, R, R^*) in three ways: M is one of K_S^\perp , $\text{NS}(S)/\mathbb{Z} \cdot [K_S]$ and $\text{NS}(S)$ and $R = R^*$ is the set of (-2) -vectors in $\text{NS}(S)$ or the image of that set modulo $\mathbb{Z} \cdot [K_S]$, but this datum is not naturally pinned, because of the existence of monodromy on $\text{NS}(S)$. This explains why the construction to be given in this paper leads to del Pezzo surfaces that are *split*, in the sense that this monodromy is trivial. To put it another way, when the Dynkin diagram has no symmetry, as in the case of E_8 , there is no monodromy on $\text{NS}(F) = \text{Pic}_F$ and whether F has a k -point is a non-trivial issue. However, del Pezzo surfaces of degree 1 always have a naturally defined k -point

(the base point of the anticanonical system) while the monodromy is significant.

For the rest of this paper G will be a split simply connected simple reductive group.

Define the torus T by the formula $\mathbb{X}^*(T) = \text{Pic}_F^G$, or $T = \text{Hom}(\text{Pic}_F^G, \mathbb{G}_m)$. By assumption, T is split but is not necessarily isomorphic to a subgroup of G .

Next, we recall the BGSS construction.

Define the incidence subvariety \tilde{G} of $G \times F$ by

$$\tilde{G} = \{(g, x) | g(x) = x\}.$$

Then $pr_2 : \tilde{G} \rightarrow F$ is smooth, since $pr_2^{-1}(x)$ is just the stabilizer $\text{Stab}_G(x)$, and $pr_1 : \tilde{G} \rightarrow G$ is projective. Moreover, there is a smooth morphism $\pi : \tilde{G} \rightarrow T$ given by

$$\pi(g, x)(L) = (\phi : g^*L \rightarrow L)(x),$$

where the isomorphism $\phi : g^*L \rightarrow L$ is part of the data of the G -linearization of the line bundle L .

Remark: Fix a k -point x on F . Take $B = pr_2^{-1}(x) = \text{Stab}_G(x)$. Then π restricts to a surjective homomorphism $B \rightarrow T$. This homomorphism is split, and choosing a splitting of it gives the standard set-up $T \hookrightarrow B \hookrightarrow G$ of the theory of reductive groups. The torus T that we have taken is an *abstract Cartan subgroup*.

In fact, $(pr_2, \pi) : \tilde{G} \rightarrow F \times T$ is smooth; each fibre is a translate of the unipotent radical of a Borel subgroup of G .

Let $\tilde{G} \rightarrow X \rightarrow G$ be the Stein factorization of pr_1 . Then $X = \mathbf{Spec} \Gamma(\tilde{G}, \mathcal{O}_{\tilde{G}})$ is the affine hull of \tilde{G} , so that $\pi : \tilde{G} \rightarrow T$ factors through X .

From the definitions of W and T there is an action of W on T and there is a commutative square

$$\begin{array}{ccc} \tilde{G} & \xrightarrow{pr_1} & G \\ \pi \downarrow & & \downarrow \rho \\ T & \longrightarrow & [T/W] \end{array}$$

The key point to be proved here is that the composite morphism $\tilde{G} \rightarrow T \rightarrow [T/W]$ factors through $pr_1 : \tilde{G} \rightarrow G$; this can be verified over \bar{k} , where it follows from (1) the existence of a maximal torus T_1 in G that is isomorphic to T and (2) the fact that the pair $(T_1, N_G(T_1))$ is a slice to the adjoint action of G on itself.

Proposition 2.1 *The morphism $\pi : \tilde{G} \rightarrow T$ is relatively symplectic and the canonical class $K_{\tilde{G}}$ is trivial.*

PROOF: The smooth morphism $pr_2 : \tilde{G} \rightarrow F$ gives, at each point y of \tilde{G} , an exact sequence

$$0 \rightarrow \mathfrak{b} \rightarrow T_{\tilde{G}}(y) \rightarrow \mathfrak{g}/\mathfrak{b} \rightarrow 0.$$

So there is an exact sequence

$$0 \rightarrow \mathfrak{u} \rightarrow T_\pi(y) \rightarrow \mathfrak{g}/\mathfrak{b} \rightarrow 0,$$

where \mathfrak{u} is the Lie algebra of the unipotent radical U of B . The Killing form shows that this sequence is self-dual, which gives the relative symplectic structure in question. Since $K_T \sim 0$, Then taking the Pfaffian of the relative symplectic structure gives the triviality of the relative canonical class $K_{\tilde{G}/T}$; since K_T is trivial, the proposition is proved. \square

From the viewpoint of algebraic surfaces, their singularities and their moduli, the significance of this diagram is that, if the edges of the Coxeter–Dynkin diagram are unlabelled (that is, if the group is of type ADE), then, the localization of the unipotent variety $Uni(G)$ at the geometric generic point of the subregular unipotent orbit in G has a du Val singularity of the same type as G and the morphism $\rho : G \rightarrow [T/W]$ has normal fibres (this can be proved over \bar{k} as a corollary of Steinberg’s theorem that in that context ρ has a section). Moreover, except in bad characteristic (taken to include any prime divisor of $n + 1$ in the case of type A_n), ρ yields a miniversal deformation of this singularity and $\pi : \tilde{G} \rightarrow T$ gives a miniversal deformation of the minimal resolution of the singularity.

3 BGSS for semi-stable bundles over uniformizable elliptic curves

Fix an elliptic curve E over a field k and a simply connected reductive group G over k . The word “bundle” will imply “locally trivial in the étale topology”. A principal G -bundle $\Xi \rightarrow E$ gives rise to an F -bundle $\mathcal{F}_\Xi = \mathcal{F} = \Xi \times^G F$ and line bundles $L_\varpi = \Xi \times^G \varpi$ on \mathcal{F} for any weight ϖ . Conversely, suppose that $p : \mathcal{F} \rightarrow E$ is an F -bundle and that $L_{\varpi_1}, \dots, L_{\varpi_l}$ are line bundles on \mathcal{F} such that L_{ϖ_i} restricts to ϖ_i on each geometric fibre of p . Then there is a reductive and simply connected group scheme G_1 over E defined as the automorphism group scheme of \mathcal{F} and all the given line bundles. If the sheaves $p_* L_{\varpi_i}$ are free on E then G_1 is constant and pulls back from a k -form of G .

Fix $\Xi \rightarrow E$. Every section σ of $\mathcal{F} \rightarrow E$ defines a co-character $[\sigma]$ by $([\sigma], \varpi) = \sigma.c_1(L_\varpi)$. We say that $[\sigma] \leq 0$ if $[\sigma].\varpi_i \leq 0$ for every fundamental dominant weight ϖ_i and $[\sigma] < 0$ if $[\sigma] \leq 0$ and $\sigma \neq 0$.

Definition 3.1 *A G -bundle $\Xi \rightarrow E$ is semi-stable if $\text{ad } \Xi$ is semi-stable as a coherent sheaf on E and is unstable if it is not semi-stable.*

This is well known ([HS2] Theorem 1.1.2) to be equivalent to other definitions. In particular, Ξ is semi-stable if and only if, for every section σ of $\mathcal{F}_\Xi \rightarrow E$, the associated cocharacter $[\sigma]$ satisfies $[\sigma] \geq 0$.

Recall also Theorem 1.3.1 of [HS2]: for any unstable principal G -bundle $\Xi \rightarrow E$, there is a proper parabolic subgroup P of G , a Levi subgroup Λ of P and a

principal Λ -bundle $\lambda \rightarrow E$ such that λ is semi-stable, $\Xi = \lambda \times^\Lambda G \rightarrow E$ and Λ is maximal among Levi subgroups with this property. Moreover, Λ is unique up to G -conjugacy and λ is determined up to isomorphism of principal bundles.

This is proved as a consequence of a unique decomposition $\text{ad } \Xi = \underline{\mathfrak{l}} \oplus \underline{\mathfrak{u}} \oplus \underline{\mathfrak{u}}^*$ of vector bundles, where $\underline{\mathfrak{l}} = \text{ad } \lambda$ consists of summands of degree 0, $\underline{\mathfrak{u}}$ consists of summands of positive degree and $\underline{\mathfrak{l}} \oplus \underline{\mathfrak{u}} = \text{ad}(\lambda \times^\Lambda P)$. In turn, this follows from Atiyah's classification of vector bundles on an elliptic curve.

Let \mathcal{G}_E^{ss} denote the stack of semi-stable G -bundles on E and $\tilde{\mathcal{G}}_E^{ss}$ the stack whose objects are pairs (Ξ, σ) where $\Xi \in \text{Ob } \mathcal{G}_E^{ss}$ and σ is a section of the F -bundle $\mathcal{F}_\Xi = \Xi \times^G F \rightarrow E$ whose associated cocharacter $[\sigma]$ is zero.

Lemma 3.2 $\tilde{\mathcal{G}}_E^{ss}$ is naturally isomorphic to the stack \mathcal{B}_E^0 of B -bundles $\beta \rightarrow E$ whose associated T -bundle $\beta \times^B T \rightarrow E$ is of degree 0.

PROOF: This follows from the tautology that giving a section of \mathcal{F}_Ξ is the same as giving a reduction of Ξ to a B -bundle. \square

In the rest of this section we assume, until the end of the proof of Theorem 3.11, that the elliptic curve E is *uniformizable*; that is, that we can write $E = \mathbb{G}_m / \langle q^{\mathbb{Z}} \rangle$. For example, this holds if either $k = \mathbb{C}$ or k is an algebraically closed complete valued field, $\text{char } k \neq 2$ and $j(E)$ is not an integer of k [BGR, 9.7]. If $\text{char } k = 2$ then there do exist uniformizable curves, derived from the Tate curve. We will return to the Tate curve in the future.

Fix a co-ordinate z on \mathbb{G}_m ; then we claim that there is a morphism $\tilde{f} : \tilde{G} \rightarrow \tilde{\mathcal{G}}_E^{ss}$ defined by

$$\tilde{f}(g, x) = ((G \times \mathbb{G}_m) / \langle \gamma \rangle, \{x\} \times E) = (\Xi, \sigma),$$

where $\gamma(h, z) = (ghg^{-1}, qz)$. The F -bundle \mathcal{F}_Ξ associated to the G -bundle $\Xi = \tilde{f}(g, x)$ is the quotient $(F \times \mathbb{G}_m) / \langle \gamma \rangle$ where $\gamma(y, z) = (g(y), qz)$ for any point $y \in F$, so that indeed the image of $\{x\} \times E$ is a section of \mathcal{F}_Ξ .

To establish the claim that $\tilde{f}(g, x)$ is a point of $\tilde{\mathcal{G}}_E^{ss}$, note that the cocharacter $[\sigma]$ of the section $\sigma = \{x\} \times E$ of \mathcal{F}_Ξ vanishes by continuity, because after specializing to the identity element $g = 1$ of G it is certainly zero.

Lemma 3.3 *There is a commutative square*

$$\begin{array}{ccc} \tilde{G} & \xrightarrow{\tilde{f}} & \tilde{\mathcal{G}}_E^{ss} \\ \pi \downarrow & & \downarrow \tilde{\pi} \\ T & \xrightarrow{\phi} & Y \end{array}$$

where ϕ is the unramified covering induced from the uniformization $\mathbb{G}_m \rightarrow E \cong \text{Pic}^0(E)$.

PROOF: This follows immediately from the definitions of the morphisms involved. \square

Lemma 3.4 $\tilde{\pi}$ is smooth.

PROOF: By Lemma 3.2 we can identify $\tilde{\mathcal{G}}_E^{ss}$ with the stack \mathcal{B}_E^0 , and then $\tilde{\pi}$ is the morphism that maps a B -bundle $\beta \rightarrow E$ to its associated T -bundle $\tau = \beta \times^B T \rightarrow E$. The smoothness of $\tilde{\pi}$ follows from the surjectivity of the map $H^1(E, \text{ad } \beta) \rightarrow H^1(E, \text{ad } \tau)$ of tangent spaces, which follows in turn from the surjectivity of the morphism $\text{ad } \beta \rightarrow \text{ad } \tau$ of coherent sheaves on E and the fact that E is 1-dimensional. \square

Lemma 3.5 $[T/W] \rightarrow [Y/W]$ is étale in an analytic neighbourhood of 0_Y and $[Y/W]$ is smooth in a Zariski neighbourhood of 0_Y .

PROOF: The morphism $\phi : T \rightarrow Y$ is the quotient by the group P , which acts freely on T , so ϕ is étale. Moreover, the group $\widetilde{W} = P \rtimes W$ acts on T and the stabilizer $\text{Stab}_{\widetilde{W}}(1_T)$, where 1_T is the identity point of T , is W , which equals $\text{Stab}_W(0_Y)$. The first part of the lemma is now established, and the rest follows from the smoothness of $[T/W]$. \square

Now consider the commutative diagram (“the basic comparison”)

$$\begin{array}{ccc} \tilde{G} & \xrightarrow{\tilde{f}} & \tilde{\mathcal{G}}_E^{ss} \\ \text{pr}_1 \downarrow & & \downarrow s \\ G & \xrightarrow{f} & \mathcal{G}_E^{ss} \end{array}$$

where $f(g) = (G \times \mathbb{G}_m)/\langle \gamma \rangle$ with $\gamma(h, z) = (ghg^{-1}, qz)$. We shall show, in Theorem 3.8 below, that this is Cartesian when restricted to the analytic neighbourhood \mathcal{U} of the identity element e of G that is described in the next Lemma.

Lemma 3.6 There is an analytic (classical or rigid) open neighbourhood \mathcal{U} of e in G on which f is étale, and so surjective.

PROOF: The codifferential f^* is, via Serre duality, a map $H^0(E, (\text{ad } \Xi)^*) \rightarrow \mathfrak{g}^*$. For the trivial G -bundle $\Xi = E \times G$ this is, by inspection, an isomorphism. So f is smooth at e , and the result follows. \square

Note that \mathcal{U} maps to an analytic (classical or rigid) neighbourhood of the trivial bundle $E \times G$ in \mathcal{G}_E^{ss} .

Recall that pr_1 and s are projective, generically finite and dominant. In fact, s is finite over the locus $\tilde{\mathcal{G}}_{E, \text{reg}}^{ss}$ of regular semi-stable bundles (those whose automorphism group is of minimal dimension, namely the rank of G).

Lemma 3.7 $\deg s = \#W = \deg \text{pr}_1$.

PROOF: We can assume that there are subgroups $T \hookrightarrow B \hookrightarrow G$ as in the usual set-up for split reductive groups, with a surjection $B \twoheadrightarrow T$. Then there is a

commutative diagram

$$\begin{array}{ccccc}
 Y & \longrightarrow & \tilde{\mathcal{G}}_{E,reg}^{ss} & \longrightarrow & \mathcal{G}_{E,reg}^{ss} \\
 & \searrow & \downarrow & & \\
 & = & Y & &
 \end{array}$$

arising from the identification made above of $\tilde{\mathcal{G}}_E^{ss}$ with the stack \mathcal{B}_E^0 . Now $\mathcal{G}_{E,reg}^{ss}$ has a geometric quotient that can be identified with $[Y/W]$ and the fibres of $\tilde{\mathcal{G}}_{E,reg}^{ss} \rightarrow Y$ are point over the regular locus in Y . That is, over the maximal open subvariety Y^0 of Y on which Y acts freely.

In other words, the square

$$\begin{array}{ccc}
 \tilde{\mathcal{G}}_{E,reg}^{ss} & \longrightarrow & \mathcal{G}_{E,reg}^{ss} \\
 \downarrow & & \downarrow \\
 Y & \longrightarrow & [Y/W]
 \end{array}$$

is Cartesian when restricted to Y^0 and this is enough. \square

Theorem 3.8 *The square*

$$\begin{array}{ccc}
 \tilde{G} & \xrightarrow{\tilde{f}} & \tilde{\mathcal{G}}_E^{ss} \\
 pr_1 \downarrow & & \downarrow s \\
 G & \xrightarrow{f} & \mathcal{G}_E^{ss}
 \end{array}$$

in the “basic comparison” is Cartesian when restricted to the neighbourhood \mathcal{U} of e in G .

PROOF: Consider the diagram

$$\begin{array}{ccccc}
 \tilde{G} & & & & \\
 \swarrow r & \searrow \tilde{f} & & & \\
 & Z & \xrightarrow{h'} & \tilde{\mathcal{G}}_E^{ss} & \\
 pr_1 \downarrow & \downarrow q & & \downarrow s & \\
 & G & \xrightarrow{f} & \mathcal{G}_E^{ss} &
 \end{array}$$

where Z is the fibre product. Since $\deg pr_1 = \deg s$, it follows that r is birational; clearly, r is proper. Also, h' is smooth on $\mathcal{U}_1 = q^{-1}(\mathcal{U})$, since f is smooth on \mathcal{U} , so that \mathcal{U}_1 is smooth. Set $\mathcal{U}_2 = r^{-1}(\mathcal{U}_1)$, an open subvariety of \tilde{G} . So $r : \mathcal{U}_2 \rightarrow \mathcal{U}_1$ is a proper birational morphism of smooth analytic k -varieties.

By Proposition 2.1 the canonical divisor class $K_{\tilde{G}}$ is trivial, and then $K_{\mathcal{U}_2} \sim 0$.

Lemma 3.9 *If $g : X \rightarrow Y$ is a proper birational morphism of smooth (rigid analytic or classical) k -varieties and $K_X \sim 0$, then g is an isomorphism.*

PROOF: This is very well known in characteristic zero but possibly less so in positive characteristic. So we give a proof.

There is an open subvariety V of Y whose complement is of codimension at least 2 over which g is an isomorphism. Therefore the trace map $tr : g_*\omega_X \rightarrow \omega_Y$ is an isomorphism on V ; since $g_*\omega_X \cong \mathcal{O}_Y$ it follows that tr is an isomorphism and that $\omega_Y \cong \mathcal{O}_Y$. Then the natural homomorphism $g^*\omega_Y \rightarrow \omega_X$ is also an isomorphism. Therefore the determinant of the derivative $dg : T_X \rightarrow g^*T_Y$ is an isomorphism, and so dg is an isomorphism. Therefore g is smooth. Since it is also proper and birational it is an isomorphism. \square

In particular, the restriction of r to \mathcal{U}_2 is an isomorphism and we are done. \square

Corollary 3.10 *$s : \tilde{\mathcal{G}}_E^{ss} \rightarrow \mathcal{G}_E^{ss}$ is proper and surjective.*

PROOF: By Theorem 3.8 the morphism s is surjective in a neighbourhood of the trivial bundle in \mathcal{G}^{ss} . The properness follows from the valuative criterion and the fact that, if $\{\Xi_t\}$ is a family of semi-stable G -bundles over E , then a section σ_t of \mathcal{F}_{Ξ_t} whose cocharacter $[\sigma_t]$ vanishes can only specialize to a section $[\sigma_0]$ of \mathcal{F}_{Ξ_0} such that $[\sigma_0] \leq 0$. Since Ξ_0 is semi-stable, $[\sigma_0] = 0$. \square

Theorem 3.11 *In a neighbourhood of the origin in $[Y/W]$ the squares*

$$\begin{array}{ccc} \tilde{\mathcal{G}}_E^{ss} & \xrightarrow{s} & \mathcal{G}_E^{ss} \\ \downarrow & & \downarrow \\ Y & \longrightarrow & [Y/W] \end{array} \quad \text{and} \quad \begin{array}{ccc} \tilde{G} & \longrightarrow & G \\ \downarrow & & \downarrow \\ T & \longrightarrow & [T/W] \end{array}$$

are smoothly equivalent.

PROOF: Define $\mathcal{H} = \mathcal{G}_E^{ss} \times_{[Y/W]} [T/W]$ and $\tilde{\mathcal{H}} = \tilde{\mathcal{G}}_E^{ss} \times_Y T$, so that the square

$$\begin{array}{ccc} \tilde{\mathcal{H}} & \longrightarrow & \mathcal{H} \\ \downarrow & & \downarrow \\ T & \longrightarrow & [T/W] \end{array}$$

is the pull-back under $[T/W] \rightarrow [Y/W]$ of the square

$$\begin{array}{ccc} \tilde{\mathcal{G}}_E^{ss} & \xrightarrow{s} & \mathcal{G}_E^{ss} \\ \downarrow & & \downarrow \\ Y & \longrightarrow & [Y/W]. \end{array}$$

We have shown that, near e , the square

$$\begin{array}{ccc} \tilde{G} & \longrightarrow & G \\ \downarrow & & \downarrow \\ \tilde{\mathcal{G}}_E^{ss} & \longrightarrow & \mathcal{G}_E^{ss} \end{array}$$

is Cartesian; the same argument shows that, near e , $G \rightarrow \mathcal{H}$ is smooth and that

$$\begin{array}{ccc} \tilde{G} & \longrightarrow & G \\ \downarrow & & \downarrow \\ \tilde{\mathcal{H}} & \longrightarrow & \mathcal{H} \end{array}$$

is Cartesian in a neighbourhood of e . So, in a neighbourhood of e ,

$$\begin{array}{ccc} \tilde{G} & \longrightarrow & G \\ \downarrow & & \downarrow \\ T & \longrightarrow & [T/W] \end{array}$$

is smoothly equivalent to

$$\begin{array}{ccc} \tilde{\mathcal{H}} & \longrightarrow & \mathcal{H} \\ \downarrow & & \downarrow \\ T & \longrightarrow & [T/W]. \end{array}$$

Comparing these two descriptions of the last square gives the result. \square

We shall see later that this result fails when E is supersingular in characteristics 2, 3 and 5 and $G = E_8$.

We can now tie up some loose ends concerning semi-stability. We no longer assume the curve E to be uniformizable.

We have recovered, as Corollary 3.10, the well known fact that the first projection $s : \tilde{\mathcal{G}}_E^{ss} \rightarrow \mathcal{G}_E^{ss}$ is surjective and proper when the base is $\mathbf{Spec} \mathbb{C}$. This is true over any base.

Lemma 3.12 *The projection $s : \tilde{\mathcal{G}}_E^{ss} \rightarrow \mathcal{G}_E^{ss}$ is surjective and projective.*

PROOF: It follows from Corollary 3.10 that $s : \tilde{\mathcal{G}}_E^{ss} \rightarrow \mathcal{G}_E^{ss}$ is surjective if E is defined over \mathbb{C} , and so over any field of characteristic zero. Suppose then that Ξ is a semi-stable G bundle over E in characteristic p ; lift E and Ξ to $\tilde{\Xi} \rightarrow \tilde{E}$ in characteristic zero. There is then a section $\tilde{\sigma}$ of $\mathcal{F}_{\tilde{\Xi}}$ with $[\tilde{\sigma}] = 0$. Specializing back to characteristic p gives a section σ of \mathcal{F}_{Ξ} in characteristic p with $[\sigma] \leq 0$; since Ξ is semi-stable, $[\sigma] = 0$.

So s is surjective. Properness follows from the valuative criterion in the same way and then projectivity is a consequence of the projectivity of the Hilbert scheme. \square

Proposition 3.13 *If the base is an algebraically closed field, then Ξ is semi-stable if and only if \mathcal{F}_Ξ has a section σ with $[\sigma] = 0$ and Ξ is unstable if and only if \mathcal{F}_Ξ has a section σ with $[\sigma] < 0$.*

PROOF: Suppose that Ξ is semi-stable. Then the existence of a suitable σ is the surjectivity to which we have just alluded. Conversely, the existence of a section σ such that $[\sigma] = 0$ leads to a description of $\text{ad } \Xi$ as an extension of line bundles of degree 0, which implies its semi-stability.

According to [HS3], 3.3, every unstable G -bundle Ξ_0 can be deformed to a semi-stable bundle Ξ_t . Then \mathcal{F}_{Ξ_t} has a section σ_t with $[\sigma_t] = 0$; specializing to $t = 0$ gives a section σ_0 of \mathcal{F}_{Ξ_0} with $[\sigma_0] \leq 0$. Since Ξ_0 is unstable, $[\sigma_0] \neq 0$. \square

Remark: There is a special case of Theorem 1.3.1 of [HS2] that we can now spell out.

Suppose that Ξ can be reduced to a semi-stable Λ -bundle where Λ is a Levi subgroup of a maximal parabolic subgroup P of G . Let $\Xi_P = \Xi/P = \Xi \times^G G/P \rightarrow E$ be the associated (G/P) -bundle. Then $\Xi_P \rightarrow E$ has a section σ whose associated cocharacter $[\sigma]$ is a negative integer multiple of α^\vee , where α^\vee is the simple coroot belonging to P .

4 Some relative compactifications of $\tilde{\mathcal{G}}_\mathcal{E}^{ss}$

We denote by \mathcal{T}^0 the stack of T -bundles over \mathcal{E} of multi-degree zero, so that $\mathcal{T}^0 \cong \mathcal{Y} \times \mathbb{B}T$, where $\mathcal{Y} = \text{Hom}(P, \mathcal{E})$ and, for a group H , $\mathbb{B}H = \{*\}/H$ is the classifying stack of H -bundles. There are morphisms $s : \tilde{\mathcal{G}}_\mathcal{E}^{ss} \rightarrow \mathcal{G}_\mathcal{E}^{ss}$ and $\tilde{\pi} : \tilde{\mathcal{G}}_\mathcal{E}^{ss} \rightarrow \mathcal{Y}$ given by $\tilde{\pi}(\Xi, \sigma)(\varpi) = L_\varpi|_\sigma$ and these stacks and morphisms fit into a commutative square, analogous to the square in the BGSS construction,

$$\begin{array}{ccc} \tilde{\mathcal{G}}_\mathcal{E}^{ss} & \xrightarrow{s} & \mathcal{G}_\mathcal{E}^{ss} \\ \tilde{\pi} \downarrow & & \downarrow \\ \mathcal{Y} & \longrightarrow & [\mathcal{Y}/W] \end{array}$$

where $s = pr_1$ is projective (in particular, representable), by Proposition 4.1 below, and $[\mathcal{Y}/W]$ is the geometric quotient, relative to \mathcal{M}_{Ell} , of the open substack $\mathcal{G}_{\mathcal{E}, reg}^{ss}$ of *regular* semi-stable bundles. Here “regular” means that the automorphism group has minimal dimension l , the rank of G .

It is well known that the classifying morphism $\mathcal{G}_{\mathcal{E}, reg}^{ss} \rightarrow [\mathcal{Y}/W]$ extends to a morphism $\gamma : \mathcal{G}_\mathcal{E}^{ss} \rightarrow [\mathcal{Y}/W]$ given by sending a bundle to its S -equivalence class.

We now recall various relative compactifications of $\tilde{\mathcal{G}}_\mathcal{E}^{ss}$. There are several that are relevant, but the most useful for us here, because of its smoothness properties, will be denoted by $\tilde{\mathcal{G}}_\mathcal{E}$ or $\tilde{\mathcal{G}}_\mathcal{E}^{KM}$; it is based on the stack of stable maps introduced by Kontsevich (although it is also appropriate to attach the name of Mori). That

is, it relies on enlarging the source of a map, rather than on Drinfel'd's idea, which we now recall, of enlarging the target.

Define $R = \bigoplus_{\varpi} H^0(F, \varpi)$, the Cox ring of F , the sum being taken over all dominant weights. Let T denote the torus introduced previously and consider the singular and non-separated stack $\overline{F} = (\mathbf{Spec} R)/T$. There is an open G -equivariant embedding $F \hookrightarrow \overline{F}$ and Drinfel'd's idea, which is described in Section 1 of [BG], and recalled in more detail below, is to embed \mathcal{F}_{Ξ} into the \overline{F} -bundle $\Xi \times^G \overline{F} \rightarrow E$ and consider sections of this. Of course, any projective homogeneous G -variety X has a similar enlargement $X \hookrightarrow \overline{X}$. This Kontsevich-Mori compactification turns out to be slightly wrong for our purposes (roughly speaking, it needs to be contracted and then flopped) but the extra information that it contains turns out to be crucial for the proof of our main result. The Drinfel'd compactification $\widetilde{\mathcal{G}}_{\mathcal{E}}^D$, on the other hand, is too small and too singular.

Let $\mathcal{C}_{pre} \rightarrow \mathbf{Spec} \mathbb{Z}$ denote the stack whose objects over a point S are 1-marked pre-stable curves $C \rightarrow S$ whose canonical model is an elliptic curve over S . So, if S is a geometric point, then C is reduced with normal crossings, its dual graph is a tree, every irreducible component except one, say C_1 , a copy of \mathbb{P}^1 and C_1 is an elliptic curve. We refer to these as ‘‘elliptic curves with rational tails’’ and to \mathcal{C}_{pre} as the stack of elliptic curves with tails. Note that \mathcal{C}_{pre} is a smooth stack over \mathbb{Z} and that its discriminant, the locus of singular curves, is a divisor $\Delta = \Delta(\mathcal{C}_{pre})$ in \mathcal{C}_{pre} with normal crossings. The complement $\mathcal{C}_{pre} - \Delta$ is naturally isomorphic to \mathcal{M}_{Ell} and there is a retraction $\mathcal{C}_{pre} \rightarrow \mathcal{M}_{Ell}$ given by sending a pre-stable curve to its canonical model. This retraction is smooth, since it is surjective on tangent spaces.

Let $\widetilde{\mathcal{G}}_{\mathcal{E}} = \widetilde{\mathcal{G}}_{\mathcal{E}}^{KM}$ be the stack whose objects over a scheme S are triples $(\Xi, C, \sigma : C \rightarrow \Xi \times^G F)$, where Ξ is a G -bundle over $E = \mathcal{E} \times_{\mathcal{M}_{Ell}} S$, C is an S -object of \mathcal{C}_{pre} and σ is a stable map (in the sense of Kontsevich) such that the composite $C \rightarrow \Xi \times^G F \rightarrow E$ is the contraction of C to its canonical model and $\deg \sigma^* L_{\varpi_i} = 0$ for each fundamental dominant weight ϖ_i . (Here, degree means total degree, the sum of the degrees on each component of C .) Let $s : \widetilde{\mathcal{G}} \rightarrow \mathcal{G}$ be the forgetful map $(\Xi, C, \sigma : C \rightarrow \Xi \times^G F) \mapsto \Xi$.

Proposition 4.1 (1) $s : \widetilde{\mathcal{G}}_{\mathcal{E}} \rightarrow \mathcal{G}_{\mathcal{E}}$ is proper and has finite relative automorphism group schemes.

(2) $pr_2 : \widetilde{\mathcal{G}}_{\mathcal{E}} \rightarrow \mathcal{C}_{pre}$ is smooth.

(3) There is a smooth morphism $\pi : \widetilde{\mathcal{G}}_{\mathcal{E}} \rightarrow \mathcal{Y}$ that extends the morphism $\tilde{\pi} : \widetilde{\mathcal{G}}_{\mathcal{E}}^{ss} \rightarrow \mathcal{Y}$ described previously.

(4) $(pr_2, \pi) : \widetilde{\mathcal{G}}_{\mathcal{E}} \rightarrow \mathcal{C}_{pre} \times_{\mathcal{M}_{Ell}} \mathcal{Y}$ is also smooth.

(5) s is representable when restricted to the locus in $\widetilde{\mathcal{G}}_{\mathcal{E}}$ where, on each geometric component of C , either σ is of degree 1 onto its image or σ is constant.

PROOF: The only things which are neither obvious nor well known are (3) and (4).

For (3), it is enough to consider the universal curve $\Gamma \rightarrow \mathcal{C}_{pre}$ and then notice that the contraction $\Gamma \rightarrow \mathcal{E} \times_{\mathcal{M}_{Ell}} \mathcal{C}_{pre}$ is a projective and birational morphism of smooth stacks. It is well known that for such a morphism there is a blowing-down morphism from \mathbb{G}_m -bundles on Γ to \mathbb{G}_m -bundles on $\mathcal{E} \times_{\mathcal{M}_{Ell}} \mathcal{C}_{pre}$, given by pushing forward divisor classes; this extends to a blowing-down morphism from T -bundles on Γ to T -bundles on $\mathcal{E} \times_{\mathcal{M}_{Ell}} \mathcal{C}_{pre}$ for any split torus T , by induction on the rank of T . So π exists; its smoothness is an immediate consequence of the fact that curves are 1-dimensional.

(4): Suppose that k is an algebraically closed field and that C is a k -point of \mathcal{C}_{pre} whose canonical model is the elliptic curve E . Since $H^1(E, \mathcal{O}_E) \rightarrow H^1(C, \mathcal{O}_C)$ is an isomorphism, it is enough to show that the morphism $\mathcal{B}_C \rightarrow \mathcal{T}_C$ from the stack of B -bundles on C to the stack of T -bundles on C is smooth. As usual, the obstruction to smoothness lies in a group of the form $H^2(C, \text{ad } \alpha)$, where $\alpha \rightarrow C$ is a principal bundle under a unipotent group, so vanishes. \square

We shall see that 4.1 5 is enough to show that s is representable over some neighbourhood of the locus of regular or subregular unstable G -bundles.

The objects of Drinfeld's compactification $\tilde{\mathcal{G}}_{\mathcal{E}}^D$ are described as follows; we refer to [BG, section 1] for the details of what is described in the next paragraph.

Fix an elliptic curve E and a G -bundle $\Xi \rightarrow E$, with $q : \mathcal{F} = \mathcal{F}_{\Xi} = \Xi \times^G F \rightarrow E$ the associated F -bundle. For every dominant weight ϖ there is a line bundle L_{ϖ} on \mathcal{F} and vector bundle $V_{\varpi} = q_* L_{\varpi}$ on E . A reduction of Ξ to a B -bundle is a line sub-bundle M_{ϖ} of V_{ϖ} (the line generated by a vector of highest weight) such that the set of all subsheaves $M_{\varpi} \hookrightarrow V_{\varpi}$, as ϖ ranges over all dominant weights, satisfies the Plücker relations. An object of $\tilde{\mathcal{G}}_{\mathcal{E}}^D$ consists of a G -bundle Ξ over an elliptic curve E and a collection of subsheaves $\{M_{\varpi} \hookrightarrow V_{\varpi}\}_{\varpi}$ where M_{ϖ} is invertible, but not necessarily a sub-bundle, that satisfies the Plücker relations; we also demand that the associated T -bundle should have degree 0. (This T -bundle is constructed as follows: objects of $\tilde{\mathcal{G}}_{\mathcal{E}}^D$ are identified with sections of the \bar{F} -bundle $\bar{\mathcal{F}} = \Xi \times^G \bar{F} \rightarrow E$ that meet the open subscheme \mathcal{F} of $\bar{\mathcal{F}}$. The open embedding $F \hookrightarrow \bar{F}$ induces an isomorphism $\text{Pic}_{\bar{F}}^G \rightarrow \text{Pic}_F^G$, so there is a natural T -bundle associated to $\bar{\mathcal{F}}$; this is to be of degree 0.)


The projection $\tilde{\mathcal{G}}_{\mathcal{E}}^D \rightarrow \mathcal{G}_{\mathcal{E}}$ is projective, while $\tilde{\mathcal{G}}^{KM} \rightarrow \mathcal{G}$ is also proper but can have non-trivial, but finite, relative automorphism groups.

Proposition 4.2 *There is a natural morphism $\tilde{\mathcal{G}}_{\mathcal{E}} \rightarrow \tilde{\mathcal{G}}_{\mathcal{E}}^D$ relative to $\mathcal{G}_{\mathcal{E}}$.*

PROOF: An object of $\tilde{\mathcal{G}}_{\mathcal{E}}$ gives sub-line bundles M'_{ϖ} of V_{ϖ} on a pre-stable curve C where each M'_{ϖ} has degree 0; pushing these sheaves M'_{ϖ} forward to E gives an object of $\tilde{\mathcal{G}}_{\mathcal{E}}^D$. \square

5 Some deformations

and the stack $\widetilde{\mathcal{G}}_{\mathcal{E}}^+$. Assume that G is a split and simply connected reductive group of type E_l , where $l = 5, 6, 7$ or 8 , and that $T \subset G$ is a copy of $\mathrm{Hom}(\mathrm{Pic}_F^G, \mathbb{G}_m)$, a maximal torus in G . Fix also a Borel subgroup B containing T . We shall assume, as we may, that these data are all given over $\mathbf{Spec} \mathbb{Z}$. For E_6, E_7 and E_8 we shall number the nodes of Dynkin diagrams as in Bourkaki's planches: the branch node is numbered 4 and the node adjacent to it on the long arm (or one of the two longer arms in the case of E_6) is numbered 5. For $E_5 = D_5$ we also number the branch node by 4, but 5 will refer to a node on one of the short arms. Here is the diagram for E_8 .



Recall, from [FM], the construction of miniversal deformation spaces for certain unstable G -bundles over an elliptic curve $E \rightarrow S$, where S is some scheme. Suppose that $P \supset B$ is a maximal parabolic subgroup, with a Levi factor $\Lambda = P \cap P^-$, where $P^- \supset B^-$ are opposite to the pair $P \supset B$. Fix a Λ -bundle $\lambda \rightarrow E$ such that

- (i) λ is semi-stable,
- (ii) $\Xi = \lambda \times^\Lambda G$ is an unstable G -bundle and
- (iii) λ is regular, in the sense that for every geometric point of S the automorphism group of λ is of minimal dimension amongst all Λ -bundles that satisfy (i) and (ii).

Then consider the stack \mathcal{H} whose objects are triples $(\lambda, \mathcal{P}^-, \phi)$, where \mathcal{P}^- is a P^- -bundle over E and ϕ is an isomorphism $\phi : \mathcal{P}^-/U^- \rightarrow \lambda$, where U^- is the unipotent radical of P^- . The forgetful map $\mathcal{H} \rightarrow \Lambda_{\mathcal{E}, reg}^{ss}$, where $\Lambda_{\mathcal{E}, reg}^{ss}$ is the stack of regular semi-stable Λ -bundles over E , is represented by an affine space bundle over $\Lambda_{\mathcal{E}, reg}^{ss}$; the fibre over λ is the non-abelian cohomology set $H^1(E, \underline{U}^-)$, where \underline{U}^- is the principal U^- -bundle $\lambda \times^\Lambda U^-$. However, it is shown in [FM], section 4, that this set is naturally an affine space isomorphic to the cohomology vector space $H^1(E, \underline{\mathfrak{u}}^-)$, where $\underline{\mathfrak{u}}^-$ is the corresponding bundle of Lie algebras. (In *loc. cit.* the base is $\mathbf{Spec} \mathbb{C}$ but their argument is valid over any base. So if E is defined over a base scheme S then $H^1(E, \underline{U}^-)$ is an affine space bundle over S .) In particular, \mathcal{H} is algebraic and smooth.

Consider the morphism $\rho : \mathcal{H} \rightarrow \mathcal{G}_{\mathcal{E}}$ given by $\rho(\mathcal{P}^-) = \mathcal{P}^- \times^{P^-} G$. Note that there is an action of the centre $\zeta(\Lambda)$ on \mathcal{H} that covers the trivial action on $\mathcal{G}_{\mathcal{E}}$, coming from the inclusion $\zeta(\Lambda) \hookrightarrow \mathcal{P}^-$. The fixed locus of this action is the stack of triples $(\lambda, \lambda \times^\Lambda P^-, \phi_{can})$, which is a copy of Λ_{reg}^{ss} . That is, the affine bundle $\mathcal{H} \rightarrow \mathcal{L}_{reg}^{ss}$ has a \mathbb{G}_m -action and a section consisting of fixed points for the \mathbb{G}_m -action.

Now suppose that we start with an unstable G -bundle ξ , defined over some algebraically closed field k , that is either regular or subregular. Then, according to [HS2], Theorem 5.12, ξ is isomorphic, up to translation by a k -point of E , to the bundle $\Xi_r = \lambda_r \times^{\Lambda_r} G$ where $r = 4$ if ξ is regular and $r = 5$ if ξ is subregular, λ_r is a semi-stable Λ_r -bundle and Λ_r is a Levi subgroup of a maximal parabolic subgroup P_r that is associated to the node numbered r . From its description, the group Λ_r is defined over \mathbb{Z} .

Lemma 5.1 *The bundle λ_r is defined over \mathcal{M}_{EU} .*

PROOF: When $r = 4$ this follows from the description of λ_r on pp. 375–376 of [HS2] and the fact that \mathcal{E} has a natural line bundle of degree 1, corresponding to the origin $0_{\mathcal{E}}$.

When $r = 5$ there is a similar description: λ_5 is determined by a $GL_q \times GL_s$ bundle (η_q, η_s) with $\deg \det \eta_s = 1$ and $\det \eta_q \cong \det \eta_s^{\otimes 2}$, where $(q, s) = (5, l - 4)$. \square

Therefore there is a versal deformation space \mathcal{Z}^\dagger for ξ that is an affine space bundle, on which \mathbb{G}_m acts with strictly positive weights, over $\text{Pic}^1(\mathcal{E})$. The fixed locus of the \mathbb{G}_m -action is a section and we can projectivize to form $\mathbb{P}(\mathcal{Z}^\dagger) \rightarrow \text{Pic}^1(\mathcal{E})$.

Let $\mathcal{Z}^{\dagger ss}$ be the semistable locus in \mathcal{Z}^\dagger ; this is open, and its complement has codimension ≥ 2 . There is a classifying morphism $\gamma : \mathcal{Z}^{\dagger ss} \rightarrow [\mathcal{Y}/W]$.

Consider the induced morphism $\phi : \mathcal{Z}^{\dagger ss} \rightarrow [\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E})$. This morphism is constant on \mathbb{G}_m -orbits and therefore factors through a morphism $\psi : [\mathcal{Z}^{\dagger ss}/\mathbb{G}_m] \rightarrow [\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E})$.

Note that $[\mathcal{Z}^{\dagger ss}/\mathbb{G}_m]$ is an open substack of the weighted projective space bundle $\mathbb{P}(\mathcal{Z}^\dagger)$ over $[\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E})$ and its complement is of codimension at least 2.

For any line bundle \mathcal{L} on $[\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E})$ that is ample relative to $\text{Pic}^1(\mathcal{E})$, let $\widehat{\mathcal{L}} \rightarrow \text{Pic}^1(\mathcal{E})$ be the family of corresponding cones over $[\mathcal{Y}/W]$ obtained by contracting the 0-section $0_{\mathcal{L}}$ of the total space of \mathcal{L} . So there is a commutative diagram

$$\begin{array}{ccc} (\mathcal{L} - 0_{\mathcal{L}}) & \hookrightarrow & \widehat{\mathcal{L}} \\ \downarrow & & \downarrow \\ [\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E}) & \xrightarrow{pr_2} & \text{Pic}^1(\mathcal{E}). \end{array}$$

Lemma 5.2 *Suppose that \mathcal{L}_1 is a line bundle on $[\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E})$ that is ample relative to $\text{Pic}^1(\mathcal{E})$. Then there is a line bundle \mathcal{A} on $\text{Pic}^1(\mathcal{E})$ such that, if $\mathcal{L} = \mathcal{L}_1 \otimes pr_2^* \mathcal{A}$, then ϕ lifts to $\tilde{\phi} : \mathcal{Z}^{\dagger ss} \rightarrow \mathcal{L} - 0_{\mathcal{L}}$.*

PROOF: Write $\mathcal{M}^0 = \psi^* \mathcal{L}_1$. Then, after replacing \mathcal{L}_1 by \mathcal{L} as in the statement of the lemma, we have $\mathcal{M}^0 = \mathcal{O}_{\mathbb{P}(\mathcal{Z}^\dagger)}(n)|_{[\mathcal{Z}^{\dagger ss}/\mathbb{G}_m]}$ for some $n \geq 1$.

Write $\mathcal{M} = \mathcal{O}_{\mathbb{P}(\mathcal{Z}^\dagger)}(n)$. Then $\mathcal{M}^0 - 0_{\mathcal{M}^0} = \mathcal{Z}^{\dagger ss}/\mu_n$ and there is a commutative diagram

$$\begin{array}{ccccccc}
 \mathcal{Z}^{\dagger ss} & \longrightarrow & (\mathcal{M}^0 - 0_{\mathcal{M}^0}) & \xrightarrow{\Psi} & (\mathcal{L} - 0_{\mathcal{L}})^c & \xrightarrow{o} & \widehat{\mathcal{L}} \\
 & \searrow & \downarrow & & \downarrow & & \downarrow \\
 & & [\mathcal{Z}^{\dagger ss}/\mathbb{G}_m] & \xrightarrow{\psi} & [\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E}) & \xrightarrow{pr_2} & \text{Pic}^1(\mathcal{E}).
 \end{array}$$

Composing the arrows in the top row gives the result. \square

Lemma 5.3 *Suppose that \mathcal{T} is a stack, that \mathcal{X} and \mathcal{V} are flat and affine over \mathcal{T} , that \mathcal{X} is normal and that \mathcal{U} is an open substack of \mathcal{X} whose complement has codimension at least 2 in every fibre of $\mathcal{X} \rightarrow \mathcal{T}$. Then every morphism $f : \mathcal{U} \rightarrow \mathcal{V}$ over \mathcal{T} extends uniquely to a morphism $\mathcal{X} \rightarrow \mathcal{V}$.*

PROOF: Suppose that $T \rightarrow \mathcal{T}$ is a smooth cover and that T is affine. Write $\mathcal{U}_T = \mathcal{U} \times_{\mathcal{T}} T$ etc. and we have a diagram

$$\begin{array}{ccc}
 \mathcal{U}_T & \xrightarrow{o} & \mathcal{X}_T \\
 & \searrow & \\
 & & \mathcal{V}_T
 \end{array}$$

of schemes, where \mathcal{X}_T and \mathcal{V}_T are affine and \mathcal{X}_T is normal. Now

$$\begin{aligned}
 \text{Mor}_{Sch/T}(\mathcal{U}_T, \mathcal{V}_T) &= \text{Mor}_{Rings/\Gamma(\mathcal{O}_T)}(\Gamma(\mathcal{O}_{\mathcal{V}_T}), \Gamma(\mathcal{O}_{\mathcal{U}_T})) \\
 &= \text{Mor}_{Rings/\Gamma(\mathcal{O}_T)}(\Gamma(\mathcal{O}_{\mathcal{V}_T}), \Gamma(\mathcal{O}_{\mathcal{X}_T})) = \text{Mor}_{Sch/T}(\mathcal{X}_T, \mathcal{V}_T)
 \end{aligned}$$

since $\Gamma(\mathcal{O}_{\mathcal{U}_T}) = \Gamma(\mathcal{O}_{\mathcal{X}_T})$. So the lemma is proved in the scheme-theoretical context and then follows in full via descent. \square

Corollary 5.4 *The morphism $\gamma : \mathcal{Z}^{\dagger ss} \rightarrow [\mathcal{Y}/W] \times_{\mathcal{M}_{EU}} \text{Pic}^1(\mathcal{E})$ lifts to a morphism $\mathcal{Z}^\dagger \rightarrow \widehat{\mathcal{L}}$.*

PROOF: This follows from the previous two lemmas. \square

Looijenga [L1] considered ample W -linearized line bundles \mathcal{L} on \mathcal{Y} such that the linearization is trivial on the line $\mathcal{L}(0_{\mathcal{Y}})$ and the polarization defined by \mathcal{L} is the standard W -invariant quadratic form on the weight lattice P .

He proved that the set of such line bundles is naturally a torsor under the finite group scheme $\text{Hom}(P/Q, \mathcal{E})$ and that over any complex point of \mathcal{M}_{EU} the ring of invariants $\bigoplus_{n \geq 0} H^0(\mathcal{Y}, \mathcal{L}^{\otimes n})^W$ is a polynomial ring and is generated by homogeneous elements whose degrees are the coefficients of the biggest (highest) root when expressed in terms of simple roots. (Recall that we are only considering simply laced groups.)

Proposition 5.5 *There exists a line bundle on \mathcal{Y} that satisfies Looijenga's hypotheses.*

PROOF: Assume that $r \leq 8$ and let Λ denote the hyperbolic lattice $\Lambda = \mathbb{Z}e_0 \oplus \bigoplus_1^r \mathbb{Z}e_i$ with $e_0^2 = -1$, $e_i^2 = 1$ for $i \geq 1$ and $e_i \cdot e_j = 0$ otherwise. Put $\kappa = 3e_0 - \sum_1^r e_i$. Then $P \cong \Lambda/\mathbb{Z}\kappa$ and the exact sequence

$$0 \rightarrow \mathbb{Z}\kappa \rightarrow \Lambda \rightarrow P \rightarrow 0$$

gives an exact sequence

$$0 \rightarrow \mathcal{Y} \rightarrow \text{Hom}(\Lambda, \mathcal{E}) \rightarrow \mathcal{E} \rightarrow 0.$$

For $j = 0, 1, \dots, r$ we have $\mathbb{Z}e_j \hookrightarrow \Lambda$, so projections $pr_j : \text{Hom}(\Lambda, \mathcal{E}) \rightarrow \mathcal{E}$. Define $\mathcal{A} = \mathcal{O}_{\mathcal{E}}(0_{\mathcal{E}})$ and $\mathcal{A}_j = pr_j^* \mathcal{A}$. Then $\mathcal{L} = \mathcal{A}_0^{-1} \otimes \mathcal{A}_1 \otimes \dots \otimes \mathcal{A}_r$ is the line bundle required. \square

That is, each regular or subregular unstable bundle Ξ_4 or Ξ_5 has a versal deformation space that is an affine bundle over $\text{Pic}^1(\mathcal{E})$, and after turning off $\text{Pic}^1(\mathcal{E})$ the affine space that is the fibre admits a classifying morphism to the cone $[\widehat{\mathcal{Y}}/W]$ corresponding to the line bundle \mathcal{L} .

Now we construct the stack $\widetilde{\mathcal{G}}_{\mathcal{E}}^+$. Fix the G -homogeneous spaces H_4, H_5 and $H_{4,5}$ associated to the sets of nodes $\{4\}$, $\{5\}$ and $\{4, 5\}$, so that $F \rightarrow H_4$ and $F \rightarrow H_5$ are \mathbb{P}^1 -bundles, $H_4 \rightarrow H_{4,5}$ and $H_5 \rightarrow H_{4,5}$ are \mathbb{P}^2 -bundles and $F \rightarrow H_{4,5}$ is a bundle whose fibre is the flag variety SL_3/B of type A_2 .

Given $S \rightarrow \mathcal{M}_{EII}$, the S -objects of $\widetilde{\mathcal{G}}_{\mathcal{E}}^+$ are triples (Ξ, σ, τ) where Ξ is a G -bundle $\Xi \rightarrow \mathcal{E} \times_{\mathcal{M}_{EII}} S$, σ a stable map $\sigma : C \rightarrow \mathcal{H}_5 = \Xi \times^G H_5$ and τ a section of $\overline{\mathcal{F}}$ such that the associated T -bundle is of degree 0. Moreover, we require that under the projection $\overline{\mathcal{F}} \rightarrow \overline{\mathcal{H}}_5 = \Xi \times^G \overline{H}_5$ the curves σ and τ should agree over a dense open subset U of $\mathcal{E} \times_{\mathcal{M}_{EII}} S$ that meets every geometric fibre of $\mathcal{E} \times_{\mathcal{M}_{EII}} S \rightarrow S$. It follows from the definition that $\widetilde{\mathcal{G}}_{\mathcal{E}}^+$ is a closed substack of $\widetilde{\mathcal{G}}_{\mathcal{E}} \times_{\mathcal{G}_{\mathcal{E}}} \widetilde{\mathcal{G}}_{\mathcal{E}}^D$.

Proposition 5.6 *The morphism $\widetilde{\mathcal{G}}_{\mathcal{E}} \rightarrow \widetilde{\mathcal{G}}_{\mathcal{E}}^D$ factors through $\widetilde{\mathcal{G}}_{\mathcal{E}}^+$.*

PROOF: This is an immediate consequence of Proposition 4.2 and the existence of a projection $\mathcal{F} \rightarrow \mathcal{H}_5$ of bundles over $\mathcal{E} \times_{\mathcal{M}_{EII}} S$. \square

6 Subregular unstable bundles and del Pezzo surfaces

We keep the notation of Section 5. In particular, $r = 4$ or 5 .

There is a decomposition

$$\mathfrak{g}_r := \text{ad } \Xi_r = \mathfrak{l}_r \oplus \mathfrak{u}_r \oplus \mathfrak{u}_r^*,$$

of vector bundles on \mathcal{E} , where $\mathbf{l}_r = \text{ad } \lambda_r$ consists of summands of degree 0, \mathbf{u}_r consists of summands of positive degree and $\mathbf{l}_r \oplus \mathbf{u}_r = \text{ad}(\lambda_r \times^{\Lambda_r} P_r)$.

We know that there is a miniversal deformation space \mathcal{Z}^\dagger for Ξ_5 with a morphism $\mathcal{Z}^\dagger \rightarrow [\widehat{\mathcal{Y}}/W]$ that preserves \mathbb{G}_m -orbits; note that this is weaker than the statement that the morphism is \mathbb{G}_m -equivariant.

Define $\widetilde{\mathcal{Z}}^\dagger = \mathcal{Z}^\dagger \times_{\mathcal{G}_\mathcal{E}} \widetilde{\mathcal{G}}_\mathcal{E}$, with projection $\nu : \widetilde{\mathcal{Z}}^\dagger \rightarrow \mathcal{Z}^\dagger$. By restricting to $\mathcal{G}_\mathcal{E}^{ss}$ we see that ν is generically finite and that $\deg \nu = \#W$. The existence of the morphisms $\pi : \widetilde{\mathcal{G}}_\mathcal{E} \rightarrow \mathcal{Y}$ and $\mathcal{Z}^\dagger \rightarrow [\widehat{\mathcal{Y}}/W]$ gives a square

$$\begin{array}{ccc} \widetilde{\mathcal{Z}}^\dagger & \longrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow \\ [\widehat{\mathcal{Y}}/W] & \dashrightarrow & [\mathcal{Y}/W] \end{array}$$

where the lower broken arrow is a rational map, the projection of a cone from its vertex to its base. On the open substack $\nu^{-1}(\mathcal{Z}^{\dagger ss})$ this square is commutative, and so there is a factorization through the line bundle $\mathcal{L} \rightarrow \mathcal{Y}$ corresponding to the cone $\widehat{\mathcal{Y}}$:

$$\begin{array}{ccccc} & & \widetilde{\mathcal{Z}}^\dagger & & \\ & \swarrow \nu^\dagger & \downarrow & \searrow & \\ \mathcal{Z}^\dagger & & \mathcal{L} & \longrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow & & \\ [\widehat{\mathcal{Y}}/W] & \longleftarrow & \widehat{\mathcal{Y}} & & \end{array}$$

where $\deg \nu^\dagger = \#W$.

The next lemma is a slight strengthening of 4.1 4.

Lemma 6.1 $\widetilde{\mathcal{Z}}^\dagger \rightarrow \text{Pic}^1(\mathcal{E}) \times_{\mathcal{M}_{EU}} \mathcal{C}_{pre} \times_{\mathcal{M}_{EU}} \mathcal{Y}$ is smooth.

PROOF: $\widetilde{\mathcal{Z}}^\dagger$ classifies B -bundles over pre-stable curves. The obstruction to smoothness lies in an extension of groups each of the form $H^2(C, \mathcal{U})$ where C is a pre-stable curve and \mathcal{U} is a commutative unipotent group scheme over C . Since C is 1-dimensional this obstruction vanishes. \square

Now restrict to the point $[0_\mathcal{E}]$ of $\text{Pic}^1(\mathcal{E})$; that is, turn off $\text{Pic}^1(\mathcal{E})$. Let \mathcal{Z} and $\widetilde{\mathcal{Z}}$ denote the resulting spaces; they fit into a commutative diagram

$$\begin{array}{ccccc} & & \widetilde{\mathcal{Z}} & & \\ & \swarrow \nu & \downarrow \mu & \searrow & \\ \mathcal{Z} & & \mathcal{L} & \longrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow & & \\ [\widehat{\mathcal{Y}}/W] & \longleftarrow & \widehat{\mathcal{Y}} & & \end{array}$$

where also $\deg \nu = \#W$. Note that, by Lemma 6.1, $\tilde{\mathcal{Z}} \rightarrow \mathcal{C}_{\mathcal{M}_{Eu}} \mathcal{Y}$ is smooth and the unstable locus $\tilde{\mathcal{Z}}^u$ in $\tilde{\mathcal{Z}}$ (defined as the inverse image $\nu^{-1}(\mathcal{Z}^u)$ of the unstable locus \mathcal{Z}^u in \mathcal{Z}) equals $\mu^{-1}(0_{\mathcal{L}})$, where $0_{\mathcal{L}}$ is the zero-section of \mathcal{L} . Moreover, $\tilde{\mathcal{Z}}$ is smooth over the stack \mathcal{C}_{pre} of elliptic curves with rational tails, and $\tilde{\mathcal{Z}}^u$ is the inverse image of the discriminant divisor $\Delta_{\mathcal{C}_{pre}}$ in \mathcal{C}_{pre} . So $\mu^{-1}(0_{\mathcal{L}})$ has normal crossings. However, we do not yet know that $\mu^{-1}(0_{\mathcal{L}})$ is reduced; this will be proved in Theorem 6.7 below.

By construction, \mathcal{Z}^u is the inverse image of the vertex of the cone $[\hat{\mathcal{Y}}/W]$, so that \mathcal{Z}^u is a surface, relative to \mathcal{M}_{Eu} , with two strata: $\mathcal{Z}^u = \{[\Xi_5]\} \cup (\mathcal{Z}^u - \{[\Xi_5]\})$; the points of $\mathcal{Z}^u - \{[\Xi_5]\}$ correspond to regular unstable bundles, all of which are isomorphic to Ξ_4 .

The next result, giving a version of Steinberg's cross-section theorem which underlies the classical BGSS construction, is due to Friedman and Morgan [FM], Theorem 5.1.1; see also [Brue]. From it, they get a new proof of Looijenga's result [L1], that $[\hat{Y}/W]$ is an affine space. They all assume that the ground field is \mathbb{C} but specialization extends the result to all characteristics, as we show.

Theorem 6.2 (1) [FM] *For any geometric point η in $\mathcal{Z}^u - \{[\Xi_5]\}$ there is a section of the morphism $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$ through η .*

(2) [L1] *$[\hat{\mathcal{Y}}/W]$ is isomorphic to an affine space bundle over \mathcal{M}_{Eu} whose fibre is \mathbb{A}^{l+1} .*

PROOF: Friedman and Morgan construct, for the regular unstable bundle Ξ_4 , a chart V that is isomorphic to $H^1(\mathcal{E}, \text{ad } \Xi_4)$, a vector bundle over \mathcal{M}_{Eu} . Their main result is that the classifying morphism $\hat{h} : V \rightarrow [\hat{\mathcal{Y}}/W]$, which preserves \mathbb{G}_m -orbits, induces a morphism $h : \mathbb{P}(V) \rightarrow [\mathcal{Y}/W]$ defined over \mathcal{M}_{Eu} which is an isomorphism over $\mathcal{M}_{Eu, \mathbb{Q}} = \mathcal{M}_{Eu} \otimes_{\mathbb{Z}} \mathbb{Q}$.

Now the formation of geometric quotients does not always commute with reduction modulo p , so we know only that $[\mathcal{Y}/W]$ is normal and the geometric fibres of $[\mathcal{Y}/W] \rightarrow \mathcal{M}_{Eu}$ are generically reduced (on the locus where W acts freely). However, $h : \mathbb{P}(V) \rightarrow [\mathcal{Y}/W]$ provides a simultaneous normalization of the fibres of the family $[\mathcal{Y}/W] \rightarrow \mathcal{M}_{Eu}$ (without the requirement for any base change) and it follows that h is an isomorphism.

Moreover, the weights of the \mathbb{G}_m -action on V and on $[\hat{Y}/W]$ are the same on both sides, so that $h^*L \cong \mathcal{O}_{\mathbb{P}(V)}(1)$ and \hat{h} is an isomorphism.

By specialization, it follows that h is an isomorphism in characteristic p , and then we deduce similarly that \hat{h} is an isomorphism in characteristic p .

By the openness of versality, \mathcal{Z} is also a chart for η ; that is, $\mathcal{Z} \rightarrow \mathcal{G}_{\mathcal{E}}$ is smooth at η , so an appropriate slice of \mathcal{Z} through η gives the section required. \square

Now fix a geometric point of \mathcal{M}_{Eu} , corresponding to an elliptic curve E over an algebraically closed field k . Fix the origin 0_Y of $Y = \text{Hom}(P, E)$ and the copy of \mathbb{A}_k^1 that is the line in $L = \mathcal{L}|_Y$ lying over 0_Y . Let $\tilde{X} \rightarrow \mathbb{A}_k^1$ denote the restriction of $\tilde{\mathcal{Z}}$ to this line. That is, \tilde{X} is the fibre over 0_Y of the smooth morphism $\tilde{\mathcal{Z}} \rightarrow Y$,

so that \tilde{X} is smooth, of dimension $\dim \tilde{X} = \dim \tilde{Z} - \dim Y = l + 3 - l = 3$, and the zero fibre \tilde{X}_0 of $\tilde{X} \rightarrow \mathbb{A}_k^1$ has normal crossings.

Now \mathbb{A}_k^1 maps isomorphically to its image Γ in $[\hat{Y}/W]$, since W acts trivially on this line \mathbb{A}_k^1 , and so there is a commutative diagram

$$\begin{array}{ccc} \tilde{X} & & \\ f \downarrow & \searrow & \\ X & \longrightarrow & \mathbb{A}_k^1 \end{array}$$

where X is the image of \tilde{X} in Z and $\tilde{X} \rightarrow X$ is proper and birational. Moreover, this diagram is \mathbb{G}_m -equivariant, since f takes \mathbb{G}_m -orbits to \mathbb{G}_m -orbits and is birational, $0 \in \mathbb{A}_k^1$ is a fixed point and $\mathbb{A}_k^1 - \{0\}$ is homogeneous (but not necessarily a torsor; this will be proved later, in Lemma 6.23) under the \mathbb{G}_m -action.

Lemma 6.3 *X has local complete intersection (LCI) singularities and X_0 is a normal surface with a unique singular point, namely the point $[\xi]$ corresponding to ξ .*

PROOF: X is the inverse image in Z of the line Γ in $[\hat{Y}/W]$. Since $[\hat{Y}/W] \cong \mathbb{A}_k^{l+1}$ and Z is smooth, the singularities of X are LCI. That $[\xi]$ is isolated follows from the fact that for the exceptional groups there is (up to choosing a point on $\text{Pic}^1(E)$, which amounts to translating ξ by a point on E) just one subregular unstable bundle and the subregular unstable locus has codimension 2 in the unstable locus. \square

Write $\mathcal{F}_r = \Xi_r \times^G F$, the associated F -bundle over E . We know, from the computation of Helmke and Slodowy, that $\dim H^0(E, \underline{\mathfrak{g}}_5) = l + 4$.

Proposition 6.4 *Suppose that σ is a section of \mathcal{F}_r that gives a negative cocharacter $[\sigma]$.*

- (1) *If $r = 4$, then $[\sigma] = -\alpha_4^\vee$.*
- (2) *If $r = 5$, then $[\sigma] = -\alpha_4^\vee$ or $-\alpha_5^\vee$ or $-\alpha_4^\vee - \alpha_5^\vee$.*

PROOF: A section σ of \mathcal{F}_r corresponds to a B -bundle $\mathcal{B}_\sigma \rightarrow E$ such that the induced bundle $\mathcal{B}_\sigma \times^B G$ equals Ξ_r .

Suppose $\underline{\mathfrak{u}}_\Sigma$ is an ideal in $\text{Lie } B$, Σ the corresponding subset of the set of positive roots, $\underline{\mathfrak{u}}_\Sigma = \underline{\mathfrak{u}}_\Sigma \times^B \mathcal{B}$ the bundle attached to this by \mathcal{B} and $2\rho_\Sigma = \sum_{\alpha \in \Sigma} \alpha$. Then, writing $[\sigma] = -\sum_{i \in I} s_i \alpha_i^\vee$, $s_i \geq 0$, we have

$$\deg \underline{\mathfrak{u}}_\Sigma = (-[\sigma], 2\rho_\Sigma) = \left(\sum s_i \alpha_i^\vee, 2\rho_\Sigma \right).$$

As $\underline{\mathfrak{u}}_\Sigma$ is a sub-bundle of $\underline{\mathfrak{g}}$, we have $\dim H^0(E, \underline{\mathfrak{g}}_r) \geq \dim H^0(E, \underline{\mathfrak{u}}_\Sigma) \geq \deg \underline{\mathfrak{u}}_\Sigma$.

Lemma 6.5 $\dim H^0(E, \underline{\mathfrak{g}}_r) > \deg \underline{\mathfrak{u}}_\Sigma$.

PROOF: Suppose otherwise. Then $\deg \underline{\mathfrak{u}}_\Sigma = \dim H^0(E, \underline{\mathfrak{u}}_\Sigma)$, and so the only possible indecomposable summands of the vector bundle $\underline{\mathfrak{u}}_\Sigma$ with slope zero must have non-trivial determinant; the remaining summands have strictly positive slope.

Next, consider the canonical reduction of $\underline{\mathfrak{g}}_r$. This is $\underline{\mathfrak{g}}_r = \underline{\mathfrak{l}}_r + \underline{\mathfrak{n}}^+ + \underline{\mathfrak{n}}^-$, where $\underline{\mathfrak{l}}_r = \zeta(\underline{\mathfrak{l}}_r) + [\underline{\mathfrak{l}}_r, \underline{\mathfrak{l}}_r]$ is the decomposition of $\underline{\mathfrak{l}}_r$ as the direct sum of its centre and the bundle of its derived subalgebra, $\zeta(\underline{\mathfrak{l}}_r)$ is a non-zero direct sum of trivial line bundles, $[\underline{\mathfrak{l}}_r, \underline{\mathfrak{l}}_r]$ is a direct sum of semi-stable bundles of degree 0, and $\underline{\mathfrak{n}}^+$ (resp. $\underline{\mathfrak{n}}^-$) is a direct sum of vector bundles of strictly positive (resp. negative) slopes. Then $\underline{\mathfrak{u}}_\Sigma$ embeds into $[\underline{\mathfrak{l}}_r, \underline{\mathfrak{l}}_r] + \underline{\mathfrak{n}}^+$, and hence $\dim H^0(E, \underline{\mathfrak{u}}_\Sigma) \leq \dim H^0(E, [\underline{\mathfrak{l}}_r, \underline{\mathfrak{l}}_r] + \underline{\mathfrak{n}}^+) < \dim H^0(E, \underline{\mathfrak{g}}_r)$. \square

Now let J be a subset of the simple roots, and take Σ_J to be the roots of the unipotent radical of the standard parabolic P_J generated by B and the negative simple roots $-\alpha$ for $\alpha \notin J$, and write ρ_J for ρ_{Σ_J} , $\underline{\mathfrak{u}}_J$ for $\underline{\mathfrak{u}}_{\Sigma_J}$.

Suppose that α is simple and $\alpha \notin J$. Let \mathfrak{s}_α denote the reflection in α . Then $\mathfrak{s}_\alpha(\Sigma_J) = \Sigma_J$ and $\mathfrak{s}_\alpha(\alpha^\vee) = -\alpha^\vee$, so $(\alpha^\vee, \rho_J) = 0$, and hence $2\rho_J = \sum_{j \in J} m_j \varpi_j$ for some integers m_j depending on J . In particular, $2\rho_{\{j\}} = m_j \varpi_j$ and $2\rho_{\{j,k\}} = a_j \varpi_j + a_k \varpi_k$ for some integers m_j, a_j, a_k . Evaluating on ϖ_j^\vee , we get $m_j(\varpi_j^\vee, \varpi_j) = (\varpi_j^\vee, 2\rho_{\{j\}}) = (\varpi_j^\vee, 2\rho)$, and so the numbers m_j can be read off from Bourbaki's planches; the vector $m = (m_1, \dots, m_l)$ is given as follows, and $\dim H^0(E, \underline{\mathfrak{g}}_r) > \deg \underline{\mathfrak{u}}_{\{j\}} = s_j m_j$ for all j .

$$\begin{aligned} E_8 : m &= (23, 17, 13, 9, 11, 14, 19, 29); \\ E_7 : m &= (17, 14, 11, 8, 10, 13, 18); \\ E_6 : m &= (12, 11, 9, 7, 9, 12); \\ D_5 : m &= (8, 7, 8, 6, 8). \end{aligned}$$

(1): $r = 4$. Then $l + 2 = \dim H^0(E, \underline{\mathfrak{g}}_4)$, $s_j = 0$ if $j \neq 4$, and $s_4 \leq 1$. But $[\sigma] < 0$, so $s_4 > 0$ and then $s_4 = 1$. This proves (1) of the Proposition.

(2): $r = 5$. Ξ_5 is induced from a semi-stable Λ_5 -bundle [HS2], and $l + 4 = \dim H^0(E, \underline{\mathfrak{g}}_5)$, so again $s_j \leq 1$.

To finish, we need to show that if $j \neq 4$ or 5 , then $s_j = 0$. If $G = E_7$ or E_8 , this follows from $l + 4 = \dim H^0(E, \underline{\mathfrak{g}}) > \dim H^0(E, \underline{\mathfrak{u}}_{\{j\}}) = s_j m_j$ and the values given above for the vector m . For $G = E_6$ or D_5 we can (eschewing science) calculate $\deg \underline{\mathfrak{u}}_{\{5,j\}}$ case by case, as necessary. If $G = E_6$ and $s_3 = 1$, we get $\deg \underline{\mathfrak{u}}_{\{5,j\}} = 10 = l + 4$, while for $G = D_5$ we get $\deg \underline{\mathfrak{u}}_{\{5,j\}} = 11, 9, 10$ when $j = 1, 2, 3$, which furnishes a contradiction and thereby concludes the proof of the Proposition. \square

We remark that a uniform treatment of all the combinatorics in this paper will be found in [GS].

Corollary 6.6 *The morphism $p : \tilde{\mathcal{G}}_\mathcal{E} \rightarrow \mathcal{G}_\mathcal{E}$ is representable over a neighbourhood of Ξ_5 that contains $\mathcal{G}_\mathcal{E}^{ss}$.*

PROOF: From Proposition 6.4, any rational component of a stable map $C \rightarrow \mathcal{F}_5$ maps birationally to its image if it is not contracted to a point. So the map $C \rightarrow \mathcal{F}_5$ has no automorphisms. \square

The next result is the crux of our paper. It is this that shows how exceptional groups lead naturally to weak del Pezzo surfaces. The strategy of the proof is to show first, by considering which components can meet, that the only triple points on the closed fibre \tilde{X}_0 lie on certain copies Q of $\mathbb{P}^1 \times \mathbb{P}^1$ and then to compute intersection numbers in as many ways as possible and use the adjunction formula as much as possible in order to derive the full result.

Note that, although the surface Q that appears might appear to be annoying, its presence enables us to recognize the other surface D_1 as a weak del Pezzo surface, because the intersection $D_1 \cap Q$ is a line on D_1 : if Q , and so that line, were missing then our argument would only show that D_1 is a surface whose first Chern class has a sign (modulo (-2) -curves). In the course of the proof we shall see that there is a smooth birational contraction where Q is contracted onto that line.

Theorem 6.7


(1) *The fibre \tilde{X}_0 is semi-stable. That is, it is a reduced union of smooth surfaces and it has normal crossings.*

(2) *\tilde{X}_0 consists of three components: $\tilde{X}_0 = D_0 + D_1 + Q$, where D_0 is the strict transform of the affine surface X_0 , Q is a copy of $\mathbb{P}^1 \times \mathbb{P}^1$, the normal bundle $\mathcal{N}_{Q/\tilde{X}}$ is isomorphic to $\mathcal{O}(-1, -1)$ and D_1 is a weak del Pezzo surface.*

(3) *Set $D_0 \cap D_1 = A$. Then A is isomorphic to E and A is an anticanonical divisor on both D_0 and on D_1 .*

(4) *$Q \cap D_0$ is a fibre β of one ruling on Q , $Q \cap D_1$ is a fibre γ of the other ruling, γ is the unique (-1) -curve on D_1 and β is a (-1) -curve on D_0 .*

(5) *There is a number t with $0 \leq t \leq l - 1$ that depends only upon G such that the dual graph of the exceptional locus of the contraction $D_0 \rightarrow X_0$ is a chain*



of transverse curves where each ϵ_i is a (-2) -curve.

(6) *There is a birational map $\tilde{X}^- \rightarrow X^-$, where X^- is smooth and $X^- \rightarrow \mathbb{A}^1$ is also semi-stable, which is constructed by first contracting Q onto γ and then successively flopping the strict transforms of $\epsilon_1, \dots, \epsilon_t$.*

(7) *$X_0^- = D_0^- + D_1^-$ where D_0^- is the minimal resolution of X_0 , D_1^- is a weak del Pezzo surface of degree $9-l$ and D_1^- contains a configuration Γ_0 of (-2) -curves of type G .*

Remark: We do not need to know the value of t , since the curves β, ϵ_i will be flopped to give a family of del Pezzo surfaces of the correct degree. This issue will be addressed in Davis' thesis [D].

PROOF: The points x of \tilde{X}_0 correspond to configurations $C = \sigma + \phi$, where σ is a section of either $\xi \times^G F$ or $\eta \times^G F$, where η is regular unstable. That is, $\eta \in X_0 - \{\xi\}$.

Since X_0 is reduced, by Theorem 6.2, it follows from Proposition 6.4 that

$$\tilde{X}_0 = D_0 + \sum_{i=1}^r m_i D_i + \sum_{j=1}^s n_j Q_j + \sum r_k F_k,$$

where D_0 is the strict transform of X_0 and the other components are projective. Moreover, points in the interior of the various components correspond to configurations $\sigma + \phi$ where σ is a copy of E and the class of ϕ is given as follows:

- (1) D_0 : $[\phi] = \alpha_4^\vee$ or α_5^\vee ;
- (2) D_i^o for $i \geq 1$: $[\phi] = \alpha_5^\vee$;
- (3) F_k^o : $[\phi] = \alpha_4^\vee$;
- (4) Q_j^o : $[\phi] = \alpha_4^\vee + \alpha_5^\vee$.

Lemma 6.8 *For $j \geq 1$ the D_j are smooth and are mutually disjoint, as are the Q_j and the F_k .*

PROOF: If two branches of the divisor $\sum_{j>0} D_j$ were to meet, then they would meet along a curve δ that parametrized sections τ of \mathcal{F}_ξ with $[\tau] = -\alpha_4^\vee - \alpha_5^\vee$; more precisely, δ would parametrize stable maps whose image is $\tau + \phi_4 + \phi_5$ with $[\phi_i] = \alpha_i^\vee$. Since $\tilde{\mathcal{G}} \rightarrow \mathcal{C}_{pre}$ is smooth, the locus of stable maps whose image contains a curve of class α_4 is of pure codimension; however, it contains δ and the surface D_0 . So $\sum_{j>0} D_j$ is smooth.

The argument for $\sum Q_j$ and $\sum F_k$ is similar. \square

Lemma 6.9 *Q_j is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$.*

PROOF: Every point of Q_j corresponds to a stable map $f : C = \sigma \cup \phi \rightarrow \mathcal{F}_\xi$ where $\sigma \cap \phi$ consists of a single point, say x , and $[\phi] = \alpha_4^\vee + \alpha_5^\vee$. Therefore Q_j is isomorphic to the variety V of $(1, 1)$ curves on the flag variety $H = SL_3/B$ of type A_2 that pass through a fixed point v_0 of H (v_0 is also here a fixed point on σ).

So it is enough to prove that $V \cong \mathbb{P}^1 \times \mathbb{P}^1$.

Recall that H is the incidence variety $\{(x, l) | x \in l\}$, where x is a point in \mathbb{P}^2 and l is a line. The projections $p : H \rightarrow \mathbb{P}^2$, $q : H \rightarrow (\mathbb{P}^2)^\vee$ map V to the set of points on a fixed line and to the set of lines through a fixed point, respectively. This gives a morphism $a : V \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$.

Suppose that γ, δ are $(1, 1)$ curves on H with $p(\gamma) = p(\delta) = m$ and $q(\gamma) = q(\delta) = n$. So m, n are lines. Then γ, δ are curves in $p^{-1}(m)$ with equal image under q . Now $p^{-1}(m)$ is a copy of the Hirzebruch surface \mathbb{F}_1 and $q : \mathbb{F}_1 \rightarrow (\mathbb{P}^2)^\vee$ exhibits \mathbb{F}_1 as the blow-up of $(\mathbb{P}^2)^\vee$ at a point. Then $\gamma = \delta$, so that a is an isomorphism, as required. \square

Lemma 6.10 $Q_j \cap (\tilde{X}_0 - Q_j) = \beta \cup \gamma$ where β, γ are opposite rulings on $\mathbb{P}^1 \times \mathbb{P}^1$.

PROOF: $Q_j \cap (\tilde{X}_0 - Q_j)$ parametrizes stable maps which are *either* of the form $f : C = \sigma \cup \phi_4 \cup \phi_5 \rightarrow \mathcal{F}_\xi$ or of the form $f : C = \sigma \cup \psi \cup \phi_4 \cup \phi_5 \rightarrow \mathcal{F}_\xi$ where in each case $[f(\phi_i)] = \alpha_i^\vee$ and $f(\psi)$ is a point. Also $\sigma \cap (\phi_4 \cup \phi_5)$ is a single point, resp., $\sigma \cap (\psi \cup \phi_4 \cup \phi_5)$ is a single point. There are three possibilities.

- (1) σ and ϕ_4 remain fixed and ϕ_5 moves along ϕ_4 . This gives β .
- (2) σ and ϕ_5 remain fixed while ϕ_4 moves along ϕ_5 . This gives γ .
- (3) All remain fixed. This gives $\beta \cap \gamma$.

□

Lemma 6.11 Each triple point of $(\tilde{X}_0)_{red}$ lies in some Q_j .

PROOF: A triple point is a stable map $f : C \rightarrow \mathcal{F}_x$ where C has at least three double points. The only possibility is $C = \sigma \cup \psi \cup \phi_4 \cup \phi_5$, which corresponds to a point $\beta \cap \gamma$ as in the proof of the previous lemma. □

We now quote what seems to be folklore.

Theorem 6.12 \mathcal{M}_{EU} is simply connected.

PROOF: This can be found at

<https://mathoverflow.net/questions/105047/fundamental-group-of-the-moduli-stack-of-elliptic-curves> □

Lemma 6.13 Every double curve of \tilde{X}_0 that does not lie in $\sum Q_j$ meets at least one Q_j and dominates E .

PROOF: We can work with the universal curve \mathcal{E} . Since \mathcal{M}_{EU} is simply connected, every geometric component of \tilde{X}_0 and every double curve is defined over \mathcal{M}_{EU} .

Suppose that A is a double curve meeting no Q_j . Then A parametrizes stable maps $f : \sigma \cup \phi_4 + \phi_5 \rightarrow \mathcal{F}_\xi$ where $\sigma \cap \phi_4$ is never equal to $\sigma \cap \phi_5$. This defines a subtraction morphism $t : A \rightarrow E$ by $t(f) = (\sigma \cap \phi_4) - (\sigma \cap \phi_5)$ which is never zero, and so is constant and defines a non-zero point on E . But on the universal curve the only rational point is zero. □

Lemma 6.14 \tilde{X}_0 is reduced and $K_{\tilde{X}} \sim \sum Q_j$.

PROOF: We argue by induction.

Start by considering a neighbourhood in \tilde{X} of some $Q_j = Q$. It meets two components L, M of \tilde{X}_0 . Assume, as our induction hypothesis, that L has multiplicity 1 in \tilde{X}_0 . Since $\tilde{X}_0 \sim 0$ as divisors on \tilde{X} , we can write

$$\tilde{X}_0 = L + mM + qQ, \quad K_{\tilde{X}} \sim pM + rQ,$$

where $m, q, p, r \in \mathbb{Z}$ and $m, q > 0$. Say $Q \cap M = \gamma$, $Q \cap L = \beta$. Note that $L.\beta = (\beta_Q)^2 = 0$ and $M.\gamma = (\gamma_Q)^2 = 0$.

So $\tilde{X}_0.\beta = 0$ gives $m = qa$ and $\tilde{X}_0.\gamma = 0$ gives $qb = 1$, so that $q = b = 1$ and $m = a$.

The adjunction formula for the curve γ in the 3-fold \tilde{X} is $2g(\gamma) - 2 = K_{\tilde{X}}.\gamma + \deg \mathcal{N}_{\gamma/\tilde{X}}$, which gives $-2 = r.(-1) - 1$, so $r = 1$ and we have

$$\tilde{X}_0 = L + mM + Q, \quad K_{\tilde{X}} \sim pM + Q.$$

The adjunction formula applied to β gives $p = 2(m - 1)$, so that $K_{\tilde{X}} \sim 2(m - 1)M + Q$.

Suppose next that A is a double curve that lies in L and meets Q . So A is a connected component of $L \cap M$, meeting Q_1, \dots, Q_s , say. Pick a sufficiently small neighbourhood U of A that contains Q_1, \dots, Q_s .

The adjunction formula for L gives $K_L \sim (m - 2)A$ (in the neighbourhood $U \cap L$ of A in L). There are (-1) -curves β_1, \dots, β_s in this neighbourhood; they satisfy $\beta_i.A = 1$ and $\beta_i.K_L = -1$, and so $m = 1$ and then $K_{\tilde{X}} \sim Q$. Since \tilde{X}_0 is connected in codimension one, this completes the proof of Lemma 6.14. \square

Now drop the distinction between the D_i and the F_k . The adjunction formula shows that $-K_{D_i} \sim \left(\sum_{j \neq i} D_j\right)|_{D_i}$.

Lemma 6.15 *If L is a smooth projective surface and $B \in |-K + L|$ is smooth, but maybe disconnected, then there are three possibilities.*

- (1) B is connected, $g(B) = 1$ and L is rational.
- (2) B is connected, $g(B) = 1$, L is elliptic ruled and B is a bisection of the ruling.
- (3) B has just two connected components B_1 and B_2 , $g(B_i) = 1$, L is elliptic ruled and each B_i is a section of the ruling.

PROOF: This is easy and well known. \square

Since X_0 has a \mathbb{G}_m -action, D_0 contains at most one curve of genus ≥ 1 . Since D_0 meets $\tilde{X}_0 - D_0$, the divisors D_0, \dots, D_r form a chain of surfaces, where, for $r \geq i \geq 1$, each $D_i \cap D_{i-1}$ is a curve A_i of genus 1. Moreover, A_1 is an anticanonical divisor on D_0 , A_r is an anticanonical divisor on D_r and, for $1 \leq i \leq r - 1$, $A_i + A_{i+1}$ is an anticanonical divisor on D_i . So D_1, \dots, D_{r-1} are elliptic ruled and D_r is rational or elliptic ruled.

Lemma 6.16 *Each A_i meets exactly one Q_j and is isomorphic to E .*

PROOF: If A_i meets s of the Q_j then the subtraction morphism $t : A_i \rightarrow E$ has a fibre $t^{-1}(0)$ consisting of s points. So t is étale of degree s in characteristic zero. Then t is étale of degree s universally. Then the kernel of the dual morphism

$t^\vee : E \rightarrow \text{Pic}^0(A_i)$ is a finite étale subgroup scheme of E of order s . But on the universal curve the only such subgroup scheme is trivial, so $s = 1$. \square

So $\tilde{X}_0 = \sum_0^r D_i + \sum_1^r Q_i$ where the D_i form a chain of surfaces that intersect in copies A_1, \dots, A_r of E and Q_j intersects D_{j-1} in the curve β_j and D_j in the curve γ_j . Both β_{j+1} and γ_j are (-1) -curves on D_j . The dual complex of \tilde{X}_0 is shown in Figure 2.




Figure 2: The dual complex of \tilde{X}_0

Lemma 6.17 D_r is geometrically rational.

PROOF: If not, then A_r is a bisection of the ruling on D_r , so that there is a universally defined morphism $E \rightarrow E$ of degree 2, which is impossible. \square

Corollary 6.18 X_0 has a simply elliptic singularity. Say its degree is d ; then $d \leq 4$. Its multiplicity is $\max\{2, d\}$ and its embedding dimension is $\max\{3, d\}$.

PROOF: We know that $K_{D_0} \sim -A_1$; it follows at once that X_0 is simply elliptic. That is, the exceptional locus in its minimal resolution is a single elliptic curve. Since X_0 has a \mathbb{G}_m -action it is isomorphic to a line bundle over A_1 , which is isomorphic to E .

There is a Cartesian diagram

$$\begin{array}{ccccc} X_0 & \hookrightarrow & X & \hookrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ \{0\} & \hookrightarrow & \mathbb{A}^1 & \hookrightarrow & \mathbb{A}^{l+1} \end{array}$$

whose horizontal arrows are closed embeddings; it exhibits X_0 as an LCI singularity. From the well known structure of these homogeneous co-ordinate rings it

follows that $d \leq 4$ and that the multiplicity and embedding dimension of X_0 are as described. \square

We now let d denote the degree of the simply elliptic singularity X_0 . Recall that l is the rank of the group G .

Proposition 6.19 $d = 9 - l$.

PROOF: The point is to show that X_0 has, first, the correct embedding dimension and, then, the correct multiplicity.

Recall that $Z \cong \mathbb{A}^{l+3}$ and that $X \rightarrow \mathbb{A}^1$ is obtained from a morphism $\pi : Z \rightarrow [\widehat{Y}/W]$ that preserves \mathbb{G}_m -orbits by restricting to the line in $[\widehat{Y}/W]$ that goes through the origin of Y . So X_0 is embedded in Z as the fibre over the vertex of $[\widehat{Y}/W]$. By Looijenga's result (Theorem 6.2 above), $[\widehat{Y}/W]$ is isomorphic to \mathbb{A}^{l+1} and the weights of the \mathbb{G}_m -action on $[\widehat{Y}/W]$ are the coefficients in the biggest root in the affine Dynkin diagram. These are given by Table 1, in which each exponent denotes the multiplicity of the relevant weight.

Table 1: Coefficients in the biggest root in the affine diagram

$$\begin{aligned} E_8 &: 1 \quad 2^2 \quad 3^2 \quad 4^2 \quad 5 \quad 6 \\ E_7 &: 1^2 \quad 2^3 \quad 3^2 \quad 4 \\ E_6 &: 1^3 \quad 2^3 \quad 3 \\ D_5 &: 1^4 \quad 2^2 \end{aligned}$$

We also need to know the weights of the \mathbb{G}_m -action coming from the action of the centre $\zeta(\Lambda_5)$ of Λ_5 . Note that $\zeta(\Lambda_5) \cong \mathbb{G}_m$.

Lemma 6.20 *The weights of the $\zeta(\Lambda_5)$ -action on Z are given in Table 2.*

Table 2: Weight multiplicities for the action of $\zeta(\Lambda_5)$ on Z

$$\begin{aligned} E_8 &: 1^2 \quad 2^3 \quad 3^3 \quad 4^2 \quad 5 \\ E_7 &: 1^4 \quad 2^4 \quad 3^2 \\ E_6 &: 1^6 \quad 2^3 \\ D_5 &: 1^8 \end{aligned}$$

PROOF: As before, we have $\mathfrak{g}_5 = \mathfrak{l}_5 \oplus \mathfrak{n}^+ \oplus \mathfrak{n}^-$. This represents the fact that the G/P_5 -bundle associated to Ξ_5 has a section σ with $[\sigma] = -\alpha_5^\vee$ (see Remark 3.13).

Consider the grading of $\mathfrak{n}^+ = \bigoplus_{0 < i \leq r} \mathfrak{n}_i^+$ defined by the action of \mathbb{G}_m via the cocharacter ϖ_5^\vee . Then the fibre \mathfrak{n}_i^+ of \mathfrak{n}_i^+ is the direct sum \mathfrak{g}_α of root spaces where

α runs over the set $I_5(i)$ of positive roots such that α_5 has coefficient exactly i in α , and r is the multiplicity of α_5 in the biggest root.

Note that \mathfrak{n}_i^+ is a representation of Λ_5 , so that $\sum_{\alpha \in I_5(i)} \alpha$ is trivial on the derived subgroup of Λ_5 . Therefore $\sum_{\alpha \in I_5(i)} \alpha = n_i \varpi_5$, where n_i depends on i and G .

Put $d_i = \deg \mathfrak{n}_i^+$, so that $d_i = (\sum_{\alpha \in I_5(i)} \alpha) \cdot \alpha_5^\vee = n_i$. But $n_i(\varpi_5 \cdot \varpi_5^\vee) = \sum_{\alpha \in I_5(i)} (\alpha \cdot \varpi_5^\vee) = i \# I_5(i)$, and then an inspection of Bourbaki's planches shows that r and the d_i are given as follows:

$$\begin{aligned} E_8 : r = 5, & \quad (d_1, \dots, d_5) = (2, 3, 3, 2, 1) \\ E_7 : r = 3, & \quad (d_1, d_2, d_3) = (4, 4, 2) \\ E_6 : r = 2, & \quad (d_1, d_2) = (6, 3) \\ D_5 : r = 1, & \quad d_1 = 8. \end{aligned}$$

From the description above of \mathfrak{g} , we know that $H^1(E, \mathfrak{n}^+) = \oplus_i H^1(E, \mathfrak{n}_i^+) = 0$, so by Riemann–Roch $\dim H^0(E, \mathfrak{n}_i^+) = \chi(E, \mathfrak{n}_i^+) = d_i$, and as $\zeta(\Lambda_5) = \mathbb{G}_m$ acts with weight i on the bundle \mathfrak{n}_i^+ and so on $H^0(E, \mathfrak{n}_i^+)$, the lemma is proved. \square

Remark: Note that $H^0(E, \mathfrak{g}_5) = H^0(E, \mathfrak{l}_5) \oplus H^0(E, \mathfrak{n}^+)$ and $\dim H^0(E, \mathfrak{l}_5) = 1$, since λ_5 is determined by a $GL_p \times GL_q \times GL_s$ bundle (η_p, η_q, η_s) with $\det \eta_p = \det \eta_q = \det \eta_s$ of degree 1 and $(p, q, s) = (1, 5, l-4)$. This provides another proof that $\dim H^0(E, \mathfrak{g}_5) = l+4$.

Lemma 6.21 *If $f : A \rightarrow B$ is an injective k -algebra homomorphism of positively graded domains such that the corresponding morphism $F : \mathbf{Spec} B \rightarrow \mathbf{Spec} A$ takes \mathbb{G}_m -orbits to \mathbb{G}_m -orbits, then there is an integer $e = e(f) > 0$ such that $f(A_n) \subset B_{en}$ for all n . That is, f multiplies degrees by e .*

PROOF: That F preserves \mathbb{G}_m -orbits means that for every n , there exists n' such that $f(A_n) \subset B_{n'}$. Suppose that $x \in A_m$ and $y \in A_n$; then

$$n \deg f(x) = \deg f(x^n) = \deg f(y^m) = m \deg f(y),$$

so that $\deg f(x)/\deg f(y) = m/n = \deg x/\deg y$, which proves the lemma. \square

The next result is crucial. To prove it we must use the fact that has now been proved, that X_0 is a simply elliptic LCI singularity and so has embedding dimension at most 4.

Proposition 6.22 *For each group $G = E_8, \dots, D_5$ the integer e is equal to 1.*

PROOF: We check the cases separately. For E_8 the tables above show that A, B are weighted polynomial rings, as follows: $A = k[1, 2^2, 3^2, 4^2, 5, 6]$ and $B = k[1^2, 2^3, 3^3, 4^2, 5]$. Then the fact that the fibre X_0 over the origin of $\mathbf{Spec} B \rightarrow \mathbf{Spec} A$ has embedding dimension at most 4 at its unique singular point forces $e = 1$ in this case, and $e = 1$ in all the other cases for exactly similar reasons. \square

Remark: It is possible instead to give a case-free proof of this proposition via the interpretation of the line bundle \mathcal{L} on \mathcal{Y} as a determinant line bundle; we plan to return to this approach [GS].

Therefore X_0 is embedded \mathbb{G}_m -equivariantly as a complete intersection in an affine space as follows:

$$\begin{aligned} E_8 : X_0 &= (6) \hookrightarrow \mathbb{A}^3(1, 2, 3) \\ E_7 : X_0 &= (4) \hookrightarrow \mathbb{A}^3(1^2, 2) \\ E_6 : X_0 &= (3) \hookrightarrow \mathbb{A}^3(1^3) \\ D_5 : X_0 &= (2, 2) \hookrightarrow \mathbb{A}^4(1^4). \end{aligned}$$

It follows that X_0 has the correct multiplicity, and the proof of Proposition 6.19 is complete. \square

Lemma 6.23 *The morphisms $\tilde{X} \rightarrow \mathbb{A}_k^1$ and $X \rightarrow \mathbb{A}_k^1$ cover the standard action of \mathbb{G}_m on \mathbb{A}_k^1 .*

PROOF: It is enough to prove the lemma for $X \rightarrow \mathbb{A}_k^1$. However, this is nothing but Proposition 6.22. \square

Next, we recover the result whose proof was sketched by Helmke and Slodowy [HS3] on deformations of simply elliptic singularities. In fact the results of Hirokado [Hi], Corollary 4.3 and his calculation of T^1 (the tangent space to a miniversal deformation space) in the proof of [Hi] Theorem 4.4 permit us to prove this in almost all characteristics.

From now on we make the following assumption: $\text{char } k = 0$ if $G = D_5$ and $\text{char } k \neq 9 - l$ if $G = E_l$ with $l = 6$ or 7 . Note that when $G = E_8$ this excludes nothing.

Hirokado shows that then simply elliptic singularities of degree at most 3 are classified exactly as in characteristic zero (in particular, they are quasi-homogeneous) and that the structure of their miniversal deformation spaces is uniform across characteristics. In particular a miniversal deformation space T of such a singularity has a \mathbb{G}_m -action and there is a hypersurface S in T where the \mathbb{G}_m -action has positive weights. The complementary line is where the singularity remains simply elliptic but the elliptic curve varies.

Theorem 6.24 *The morphism $Z \rightarrow [\hat{Y}/W]$ is the positive weight part of a \mathbb{G}_m -equivariant miniversal deformation of X_0 .*

PROOF: Suppose that $V \rightarrow S$ is the positive weight part of a \mathbb{G}_m -equivariant miniversal deformation of X_0 . Certainly $Z \rightarrow [\hat{Y}/W]$ is \mathbb{G}_m -equivariant and of positive weight, so there is a \mathbb{G}_m -equivariant morphism $a : [\hat{Y}/W] \rightarrow S$ and a Cartesian square

$$\begin{array}{ccc} Z & \longrightarrow & V \\ f \downarrow & & \downarrow \\ [\hat{Y}/W] & \xrightarrow{a} & S. \end{array}$$

The derivative of the morphism $a : [\widehat{Y}/W] \rightarrow S$ at the vertex of $[\widehat{Y}/W]$ is determined uniquely, and then a is determined uniquely by its \mathbb{G}_m -equivariance. Inspection shows that the weights of the \mathbb{G}_m action on the affine spaces $[\widehat{Y}/W]$ and S are the same, so that a is an isomorphism. \square

In particular, the family $X \rightarrow \mathbb{A}^1$ is induced by a unique \mathbb{G}_m -equivariant morphism $\phi : \mathbb{A}^1 \rightarrow S$. Recall that the base \mathbb{A}^1 of $X \rightarrow \mathbb{A}^1$ is the generator in the cone $[\widehat{Y}/W]$ that passes through the origin 0_Y of Y .

Proposition 6.25 (1) *Under the morphism a , the line $\mathbb{A}^1 = \mathbf{Spec} k[t]$ in $[\widehat{Y}/W]$ is mapped to a line $\ell = \mathbf{Spec} k[x]$ in S where for $x \neq 0$ the surface V_x has a du Val singularity of type G .*

(2) *For $t \neq 0$, \widetilde{X}_t is the minimal resolution of X_t ; it contains a (-2) -configuration Γ_t of type G .*

PROOF: Suppose first that $k = \mathbb{C}$. Then, by Theorem (7.7) of [L2], the monodromy around V_s is the full Weyl group. That is, take a small neighbourhood U of s in S and consider the complement $U - \Delta$ of the discriminant; then $\pi_1(U - \Delta)$ acts on the cohomology of the generic fibre as the full Weyl group. Since V_s has at most du Val singularities, (1) is proved in this case. Specialization then proves it for all fields.

(2) is a corollary of (1) and Theorem 3.11. \square

Remark: It would be desirable to have a conceptual proof of this proposition, maybe as follows. The copy of $\mathbb{A}^1 = \mathbf{Spec} k[t]$ that appears is the line through the origin 0_Y of \mathcal{Y} so that the bundles parametrized by $\mathcal{Z}|_{t \neq 0}$ are unipotent (and semi-stable).

On the other hand, general considerations of deformation theory show that the automorphism group of each of these bundles has dimension strictly less than that of Ξ_5 , which is $l + 4$. Therefore they are subregular or regular; if we could prove that subregular bundles do arise over points of $\mathbb{A}^1 - \{0\}$ (that is, if we could prove directly that Ξ_5 deforms to a subregular unipotent semi-stable bundle) then Theorem 3.11 could be applied, assuming that E is uniformizable; the result would follow for all E by specialization. As it is, we only know how to deduce the existence of such a subregular deformation from Proposition 6.25.

Proposition 6.26 $r = 1$.

PROOF: Since $K_{\widetilde{X}} \sim \sum Q_i$, the divisors D_1, \dots, D_r appear with discrepancy zero in the resolution \widetilde{X} of the 3-fold X . They also dominate the closed point 0 that represents ξ . So it is enough to exhibit a partial resolution $X' \rightarrow X$ which has c-dV singularities and in which there is only one divisor of discrepancy zero that dominates the origin.

For this, note that from Lemma 6.18 and the proof of Proposition 6.17 it

follows that the Cartesian diagram

$$\begin{array}{ccc} X & \hookrightarrow & Z \\ \downarrow & & \downarrow \\ \mathbb{A}_k^1 & \hookrightarrow & \mathbb{A}_k^{l+1} \end{array}$$

exhibits X as a homogeneous complete intersection of codimension l in some affine space.

For the groups E_8, E_7, E_6 and D_5 , X is, respectively, a homogeneous sextic in $\mathbf{Spec} k[1^2, 2, 3] = \mathbb{A}_k^4$, a quartic in $\mathbf{Spec} k[1^2, 2]$, a cubic in $\mathbf{Spec} k[1^4]$ and a complete intersection of two quadrics in $\mathbf{Spec} k[1^5]$. We know that for $t \neq 0$ the fibre X_t has du Val singularities, so that in each case the subvariety $\text{Proj } \mathcal{O}_X$ in the appropriate weighted projective space has only du Val singularities and is a del Pezzo surface of degree $9 - l$. Therefore the weighted blow-up $X' = \text{Proj } \mathcal{O}_X[t] \rightarrow X$ centred at 0 has a unique exceptional divisor, namely $\text{Proj } \mathcal{O}_X$. Since this divisor is Cartier and has du Val singularities, it follows that X' has only c-dV singularities and is a partial resolution of the kind that we sought.

□

So $\tilde{X}_0 = D_0 + D_1 + Q$ where D_1 is rational and $A = D_0 \cap D_1$ is an effective anticanonical divisor on each of D_0, D_1 . Say $\beta = Q \cap D_0$, $\gamma = Q \cap D_1$. Then each of β, γ is a (-1) -curve.

We define a $(-1, -2)$ -chain of length s on D_0 to be an s -tuple $(\delta_1, \dots, \delta_s)$ of smooth rational curves on D_0 where $\delta_1^2 = -1$, $\delta_i^2 = -2$ for $i \geq 2$, $\delta_1 \cdot A = 1$, $\delta_i \cdot A = 0$ for $i \geq 2$ and the configuration $A, \delta_1, \dots, \delta_s$ is a chain.

Lemma 6.27 *The only complete curves on D_0 are A and those curves that occur in a $(-1, -2)$ -chains. Exactly one of these chains contains β .*

PROOF: This follows from the adjunction formula, the fact that $A \sim -K_{D_0}$ and the fact that any configuration of complete curves on D_0 is negative definite. □

It follows that there is a birational contraction $\tilde{X} \rightarrow \tilde{X}^+$ where Q is contracted onto the curve γ in D_1 and where the images in \tilde{X}^+ of the curves in the $(-1, -2)$ -chains in D_0 can be flopped successively onto D_1 , say via $\tilde{X}^+ \rightarrow \tilde{X}^-$, to give $\tilde{X}^- \rightarrow \mathbb{A}^1$ where $\tilde{X}_0^- = D_0^- + D_1^-$ is semi-stable, $K_{\tilde{X}^-} \sim 0$, D_0^- is the strict transform of D_0 and is the minimal resolution of X_0 , D_1^- is the strict transform of D_1 and is a weak del Pezzo surface on which A is an anti-canonical curve and the flopped curves on D_1^- form a configuration of $(-1, -2)$ -chains which is isomorphic to the configuration of $(-1, -2)$ -chains on D_0 .

Moreover, these birational transformations are \mathbb{G}_m -equivariant, so that \tilde{X}^- has a \mathbb{G}_m -action that covers the standard action on \mathbb{A}_k^1 and is free on $D_0^- - A^-$, where A^- is the strict transform of A . The action covers the standard action on \mathbb{A}_k^1 and so is free on $\tilde{X}^- - D_1^-$.

Since $(A_{D_1^-}^-)^2 = -(A_{D_0^-}^-)^2$, it follows that D_1^- is a weak del Pezzo surface of degree $d = 9 - l$.

It is easy to see that if \mathbb{G}_m acts on a del Pezzo surface S , weak or not, and preserves a smooth member of $| -K_S |$, then the action on S is trivial. So \mathbb{G}_m acts freely on $\tilde{X}^- - D_1^-$ and trivially on the Cartier divisor D_1^- .

The next lemma and its proof are taken almost without change from a paper by Bass and Haboush [BH].

Lemma 6.28 *Assume that S is a normal k -variety and that D is a complete relatively LCI closed subscheme of S that contains every complete subscheme of S . Assume also that there is a \mathbb{G}_m -action on S that is trivial on D and free on $S - D$ and that every closed \mathbb{G}_m -invariant subscheme of S meets D . Then there is a \mathbb{G}_m -equivariant isomorphism $S \rightarrow N_{D/S}$ that identifies D with the zero section of $N_{D/S}$.*

PROOF: The aim is to reduce this to a situation where the arguments of [BH] can be applied.

We need to show that there is a \mathbb{G}_m -linearized locally free sheaf \mathcal{A} on D such that $S \cong \mathbf{Spec} \operatorname{Sym}^* \mathcal{A}$.

Choose an open cover $S = \cup_{i \in I} S_i$ by affine \mathbb{G}_m -invariant open subschemes $S^{(i)} = \mathbf{Spec} R^{(i)}$ such that each $D^{(i)} = D \cap S^{(i)}$ is, if non-empty, defined by a regular sequence of length $r = \operatorname{codim}(D, S)$ in $R^{(i)}$. Say $J = \{j \in I \mid D^{(j)} \neq \emptyset\}$. Then $S - \cup_{j \in J} S^{(j)}$ is a closed \mathbb{G}_m -invariant subscheme of S that is disjoint from D , so is empty. So we can suppose that each $D^{(i)}$ is non-empty.

The existence of the \mathbb{G}_m -action is equivalent to a \mathbb{Z} -grading $R^{(i)} = \bigoplus_{n \in \mathbb{Z}} R_n^{(i)}$. Since every closed \mathbb{G}_m -invariant subscheme of S meets D , the modules $R_n^{(i)}$ vanish for $n < 0$. and the ideal of $D^{(i)}$ in $S^{(i)}$ is $R_+^{(i)} = \bigoplus_{n > 0} R_n^{(i)}$. So $D^{(i)} = \mathbf{Spec} R_0^{(i)}$, where we regard $R_0^{(i)}$ both as a subring of $R^{(i)}$ and as the quotient $R^{(i)}/R_+^{(i)}$.

That is, there is a unique \mathbb{G}_m -equivariant retraction $r_i : S^{(i)} \rightarrow D^{(i)}$. Since they are unique, these retractions glue to a \mathbb{G}_m -equivariant retraction $r : S \rightarrow D$. Note that, as a subscheme of S , D contains all the closed \mathbb{G}_m -orbits in S .

Now we follow [BH], pp. 474 *et seq.* We have $S = \mathbf{Spec} \mathcal{R}$, $\mathcal{R} = \bigoplus_{n \geq 0} \mathcal{R}_n$, a graded sheaf of \mathcal{O}_D -algebras. Set $\mathcal{I} = \bigoplus_{n > 0} \mathcal{R}_n$, the ideal sheaf of the subscheme D of S , $\mathcal{N}^\vee = \mathcal{I}/\mathcal{I}^2$ and $Y = \mathbf{Spec} \operatorname{Sym}^* \mathcal{N}^\vee$, the normal bundle. The tautological \mathbb{G}_m -equivariant morphism $u : S \rightarrow Y$ fits into a $\mathbb{G}_{m,D}$ -equivariant commutative diagram

$$\begin{array}{ccc} S & \xrightarrow{u} & Y \\ r \downarrow & \searrow q & \\ D & & \end{array}$$

Then the argument of [BH], p. 474, applies directly to show that u is étale.

Let $D_0 \hookrightarrow Y$ be the zero section of q .

We now claim that u is finite. To see this, take \tilde{S} to be the normalization of Y in the function field of S . Then there is a \mathbb{G}_m -equivariant open embedding

$S \hookrightarrow \tilde{S}$ and a finite dominant morphism $\nu : \tilde{S} \rightarrow Y$ extending u . Say $V = \tilde{S} - S$; this is closed and \mathbb{G}_m -invariant in Y . Since $D \subset S$ and D is complete, D is disjoint from V . Assume that $V \neq \emptyset$; then, over any affine chart of Y , there is a \mathbb{G}_m -invariant function f on \tilde{S} such that $f|_D = 0$ and $f|_V = 1$. However, $\mathcal{O}_{\tilde{S}}^{\mathbb{G}_m} = \mathcal{O}_D$, so that $V = \emptyset$ and the claim is established.

Then $u^{-1}(D_0) = D \amalg D'$ with D' finite and étale over D_0 . The argument just given, to show that $V = \emptyset$, can now be applied to show that $D' = \emptyset$. So u has degree 1 over D , and so is of degree 1 everywhere. This proves Lemma 6.28. \square

Corollary 6.29 (1) *There are \mathbb{G}_m -equivariant isomorphisms $\tilde{X}^- \rightarrow N_{D_1^-/\tilde{X}^-}$ and $N_{D_1^-/\tilde{X}^-} \rightarrow \omega_{D_1^-}$.*

(2) *X is the cone over the anti-canonical model of D_1^- .*

PROOF: The first isomorphism of (1) is a special case of the lemma and the second is a consequence of the adjunction formula. (2) is an immediate consequence.

\square

Proposition 6.30 *Suppose that D is a weak del Pezzo surface of degree $9 - l$ and contains a (-2) -configuration Δ of type G . Then*

- (1) *D contains a unique line L ;*
- (2) *if $\delta_1, \dots, \delta_l$ are the irreducible components of Δ , numbered as in Bourbaki's planches, then L meets Δ in δ_l and in no other component;*
- (3) *$\{\delta_1, \dots, \delta_l, L\}$ is a \mathbb{Z} -basis of $\text{NS}(D)$;*
- (4) *given a smooth member A of $|-K_D|$, D is obtained by embedding A in \mathbb{P}^2 via $|3[0_A]|$ and then making l successive blow-ups, with the centre of each blow-up being the origin 0_A on the strict transform of A .*

PROOF: This is well known, but we include a proof for lack of a convenient reference.

Existence of a line: D is a specialization of a smooth del Pezzo D_t where $-K_{D_t}$ is ample, which certainly contains lines. So D does also.

Uniqueness of the line: since $\text{rank NS}(D) = \text{rank } \Delta + 1$, $\text{NS}(D)_{\mathbb{Q}}$ is spanned by L and Δ for any line L . Moreover, $L \cdot \delta \geq 1$ for any positive root δ with $\text{Supp } \delta = \Delta$. Put $H = -K_D$ and consider cases separately.

$d = 1$. Then every simple root has multiplicity ≥ 2 in the biggest root δ_{max} , so $L \cdot \delta_{max} \geq 2$. Then $H^2 = 1 = H \cdot (L + \delta_{max})$ and $(L + \delta_{max})^2 \geq 1$, so that, by the index theorem, $L + \delta_{max} \sim H \sim M + \delta_{max}$ for any lines L, M . So $L \sim M$ and then $L = M$.

$d \geq 2$. Then $(L + \delta)^2 \leq 0$ for any line L and any effective root δ , by the index theorem, since $H \cdot (L + \delta) = 1$ and $H^2 \geq 2$. Suppose that δ, ϵ are effective roots with $\text{Supp } \delta = \text{Supp } \epsilon = \Delta$ and that L, M are distinct lines; then $L \cdot \delta, M \cdot \delta \geq 1$ and $(L + M + \delta)^2 \geq 2$. But $H \cdot (L + M + \delta) = 2 \leq H^2$, so that (index) $L + M + \delta \sim H \sim L + M + \epsilon$ and $\delta \sim \epsilon$, which is absurd, and uniqueness is established.

Moreover, the line L meets Δ in a simple root δ of minimal multiplicity (2 if $l = 8$, 1 otherwise). Then successive contraction of $L, \delta_l, \dots, \delta_4, \delta_3, \delta_1$, in that order, is the inverse of the blow-up described in (4), and also proves (3). \square

Now we can complete the proof of Theorem 6.7.

At this point we know, by Proposition 6.19 and Corollary 6.7 2, that D_1^- is a weak del Pezzo surface of degree $d = 9 - l$ and that it contains a (-2) -configuration $\Delta = \{\delta_1, \dots, \delta_l\}$ of type $G = E_l$. So the configuration on D_1^- formed by the strict transform γ^- of γ and the flopped curves (see the paragraph following Lemma 6.7 5) form part of the configuration $L, \delta_1, \dots, \delta_l$ given by Proposition 6.30 and so form a single $(-1, -2)$ -chain. Therefore, by Lemma 6.27, the flopping curves on D_0 form a unique $(-1, -2)$ -chain on D_0 .

This completes the proof of Theorem 6.7. \square

7 Universal families of del Pezzo surfaces

Theorem 6.7 refers to the line in the line bundle $\mathcal{L} \rightarrow \mathcal{Y}$ that is the fibre over the origin 0_Y , which is in turn defined over an algebraically closed field. However, this extends over the whole of \mathcal{L} as follows.

Recall that \mathcal{Z} is obtained from a chart of $\mathcal{G}_{\mathcal{E}}$ by turning off $\text{Pic}^1(\mathcal{E})$ and that $\tilde{\mathcal{Z}} = \mathcal{Z} \times_{\mathcal{G}_{\mathcal{E}}} \tilde{\mathcal{G}}_{\mathcal{E}}$. There is a morphism $\rho : \tilde{\mathcal{Z}} \rightarrow \mathcal{L}$ such that the composite $\tilde{\mathcal{Z}} \rightarrow \mathcal{Y}$ is smooth and $\rho^{-1}(0_{\mathcal{L}})$ coincides with the inverse image of $\Delta \times 0_{\mathcal{Y}}$, where Δ is the discriminant divisor in \mathcal{C}_{pre} , under the smooth morphism $\tilde{\mathcal{Z}} \rightarrow \mathcal{C}_{pre} \times_{\mathcal{M}_{Ell}} \mathcal{Y}$. So $\rho^{-1}(0_{\mathcal{L}}) = \mathcal{Q} + \mathcal{D}_0 + \mathcal{D}_1$.

Observe that there is a blowing-down morphism $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{Z}}^+$ where \mathcal{Q} is contracted onto a curve in \mathcal{D}_1 , because \mathcal{Q} is a family of quadrics with normal bundle $\mathcal{O}(-1, -1)$.

Recall the stack $\tilde{\mathcal{G}}^+$ from Section 5.

Lemma 7.1 $\tilde{\mathcal{Z}}^+$ is isomorphic to $\mathcal{Z} \times_{\mathcal{G}} \tilde{\mathcal{G}}^+$.

PROOF: $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{Z}}^+$ is the contraction of a $\mathbb{P}^1 \times \mathbb{P}^1$ -bundle \mathcal{Q} over \mathcal{Y} to a \mathbb{P}^1 -bundle γ . Fibre by fibre, this is the projection $q : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ given by taking the family of $(1, 1)$ -curves in SL_3/B that pass through a fixed point and projecting to the family of lines in \mathbb{P}^2 through a fixed point. Generically, these $(1, 1)$ -curves arise as the rational tail ϕ in a pre-stable curve $C = E \cup \phi$. From the construction of $\tilde{\mathcal{G}}^+$ and the morphism $\psi : \tilde{\mathcal{G}} \rightarrow \tilde{\mathcal{G}}^+$, this projection is exactly achieved by ψ over \mathcal{Z} . \square

Proposition 7.2 (1) There are sequences of flops $\tilde{\mathcal{Z}}^{+-} \rightarrow \tilde{\mathcal{Z}}^-$ that extend the sequence of flops $\tilde{X}^{+-} \rightarrow \tilde{X}^-$.

(2) They fit into a commutative diagram

$$\begin{array}{ccccc}
 & & \tilde{\mathcal{Z}} & & \\
 & \swarrow & \vdots & \searrow \rho & \\
 \tilde{\mathcal{Z}}^+ & \cdots \rightarrow & \tilde{\mathcal{Z}}^- & \xrightarrow{\sigma} & \mathcal{L} \longrightarrow \mathcal{Y} \\
 & \searrow & \downarrow & \square \text{ 1} & \downarrow \\
 & & \mathcal{Z} & \longrightarrow & [\hat{\mathcal{Y}}/W]
 \end{array}$$

where the square marked $\square 1$ is birationally Cartesian.

(3) $\tilde{\mathcal{Z}}, \tilde{\mathcal{Z}}^+$ and $\tilde{\mathcal{Z}}^-$ are semi-stable over \mathcal{L} . Their restrictions to $\mathcal{L} - 0_{\mathcal{L}}$ are isomorphic and smooth over $\mathcal{L} - 0_{\mathcal{L}}$.

(4) $\tilde{\mathcal{Z}}|_{0_{\mathcal{L}}} = \mathcal{D}_0 + \mathcal{D}_1 + \mathcal{Q}$, $\tilde{\mathcal{Z}}^+|_{0_{\mathcal{L}}} = \mathcal{D}_0^+ + \mathcal{D}_1^+$, $\tilde{\mathcal{Z}}^-|_{0_{\mathcal{L}}} = \mathcal{D}_0^- + \mathcal{D}_1^-$, and $\mathcal{D}_i, \mathcal{D}_i^\pm$ and \mathcal{Q} are smooth over $\mathcal{Y} = 0_{\mathcal{L}}$.

(5) Each of $\mathcal{D}_0 \cap \mathcal{D}_1$ and $\mathcal{D}_0^\pm \cap \mathcal{D}_1^\pm$ is isomorphic over \mathcal{Y} to $\mathcal{E} \times_{\mathcal{M}_{Eu}} \mathcal{Y}$.

(6) Each fibre of $\mathcal{D}_0^- \rightarrow \mathcal{Y}$ is the minimal resolution of the minimally elliptic singularity of degree d that belongs to the corresponding elliptic curve.

PROOF: To prove the existence of the flops there is a local problem and a global problem to be solved.

The local problem is to show that under deformation of the ambient 3-fold the curves to be flopped do not disappear and that under specialization they do not break up.

For each successive flop, the irreducible curve δ to be flopped has normal bundle $\mathcal{N}_\delta = \mathcal{O}(-1)^{\oplus 2}$. So $H^1(\delta, \mathcal{N}_\delta) = 0$ and then δ survives under deformation.

After δ has been flopped, the number of irreducible curves to be considered has diminished, and then induction shows that all the curves to be flopped survive under deformation.

A similar argument shows that they do not break up under specialization.

The global problem is to show that there is no monodromy acting on the lattice generated by the classes $\beta, \epsilon_1, \dots, \epsilon_t$ in the Chow group of the generic fibre of the family $\mathcal{D}_0 \rightarrow \mathcal{Y}$. This will follow from the next result.

Proposition 7.3 \mathcal{Y} is simply connected.

PROOF: We check first that \mathcal{E} is simply connected.

Suppose that $\pi : \tilde{\mathcal{E}} \rightarrow \mathcal{E}$ is finite and étale and that $\tilde{\mathcal{E}}$ is connected.

Since \mathcal{M}_{Eu} is simply connected, by Theorem 6.12, the origin $0_{\mathcal{E}}$ lifts to $\tilde{\mathcal{E}}$ and

we have a 2-commutative diagram

$$\begin{array}{ccc}
 0_{\mathcal{E}} & \xrightarrow{i} & \mathcal{E} \\
 \searrow j & & \downarrow \pi \\
 & & \tilde{\mathcal{E}} \\
 \cong \searrow & & \downarrow \alpha \\
 & & \mathcal{M}_{ELL}
 \end{array}$$

Then $\beta : \tilde{\mathcal{E}} \rightarrow \mathcal{M}_{ELL}$ is an elliptic curve with identity $0_{\tilde{\mathcal{E}}} = j(0_{\mathcal{E}})$. This is induced by a morphism $\gamma : \mathcal{M}_{ELL} \rightarrow \mathcal{M}_{ell}$ and we get a commutative diagram

$$\begin{array}{ccccc}
 0_{\mathcal{E}} & \xrightarrow{i} & & & \\
 \searrow j & & & & \\
 & & \mathcal{E} \times_{\mathcal{M}_{ELL}, \alpha} \mathcal{M}_{ELL} & \xrightarrow{\pi} & \tilde{\mathcal{E}} \\
 \cong \searrow & & \downarrow \beta & & \downarrow \alpha \\
 & & \mathcal{M}_{ELL} & \xrightarrow{\gamma} & \mathcal{M}_{ELL}
 \end{array}$$

Then γ is an isomorphism, and so π is an isomorphism and \mathcal{E} is simply connected.

Since \mathcal{Y} is a fibre product of copies of \mathcal{E} it is enough to check that, if $\mathcal{U} \rightarrow \mathcal{M}$ and $\mathcal{V} \rightarrow \mathcal{M}$ have sections $0_{\mathcal{U}}$ and $0_{\mathcal{V}}$ and if \mathcal{M}, \mathcal{U} and \mathcal{V} are simply connected, then so is $\mathcal{U} \times_{\mathcal{M}} \mathcal{V} = \mathcal{W}$, say.

Suppose that $\tilde{\mathcal{W}} \rightarrow \mathcal{W}$ is finite and étale. The sections $0_{\mathcal{U}}$ and $0_{\mathcal{V}}$ give sections $\tilde{0}_{\mathcal{U}}$ and $\tilde{0}_{\mathcal{V}}$ of $\mathcal{W} \rightarrow \mathcal{U}$ and of $\mathcal{W} \rightarrow \mathcal{V}$, which then lift to sections $\sigma : \mathcal{U} \rightarrow \tilde{\mathcal{W}}$ and $\tau : \mathcal{V} \rightarrow \tilde{\mathcal{W}}$. This gives a section $(\sigma, \tau) : \mathcal{U} \times_{\mathcal{M}} \mathcal{V} \rightarrow \tilde{\mathcal{W}}$, as required. \square

This completes the proof of Proposition 7.2. \square

Corollary 6.29 generalizes straightforwardly to this situation.

Proposition 7.4 (1) *Locally on \mathcal{M}_{ELL} there are \mathcal{Y} -isomorphisms $\tilde{\mathcal{Z}}^- \xrightarrow{\cong} N_{\mathcal{D}_1^-/\tilde{\mathcal{Z}}^-}$ and $N_{\mathcal{D}_1^-/\tilde{\mathcal{Z}}^-} \xrightarrow{\cong} \omega_{\mathcal{D}_1^-/\mathcal{Y}}$.*

(2) *Locally on \mathcal{M}_{ELL} there is an isomorphism, relative to \mathcal{Y} , from \mathcal{Z} to a family of 3-fold cones, namely, to the cone over the anticanonical model of the family $\mathcal{D}_1^- \rightarrow \mathcal{Y}$ of weak del Pezzo surfaces.*

Most of this section is devoted to proving that $\mathcal{D}_1^- \rightarrow \mathcal{Y}$ is close to being universal.

Let $\tau : \tilde{\mathcal{Z}}^- \rightarrow \mathcal{D}_1^-$ be the projection of $N_{\mathcal{D}_1^-/\tilde{\mathcal{Z}}^-}$ to its base. By construction, there is a smooth morphism $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{G}}$; this restricts to give a morphism

$$\tilde{\mathcal{Z}}^0 := \tilde{\mathcal{Z}} - \rho^{-1}(0_{\mathcal{L}}) = \tilde{\mathcal{Z}}^- - \sigma^{-1}(0_{\mathcal{L}}) \rightarrow \tilde{\mathcal{G}}^{ss}.$$

We identify \mathcal{D}_1^- with the zero section of $N_{\mathcal{D}_1^-/\tilde{\mathcal{Z}}^-}$ and \mathcal{D}_0^- with $\tau^{-1}(\mathcal{A})$.

Proposition 7.5 *There is an embedding $\text{Pic}_F^G \hookrightarrow \text{Pic}_{\tilde{\mathcal{G}}_\varepsilon^{ss}}$.*

PROOF: A point P of $\tilde{\mathcal{G}}_\varepsilon^{ss}$ consists of a G -bundle Ξ over an elliptic curve E and a section σ of $\mathcal{F}_\Xi \rightarrow E$ whose cocharacter $[\sigma]$ vanishes. On the other hand, an element ϖ of Pic_F^G gives a line bundle \mathcal{L}_ϖ on \mathcal{F}_Ξ ; evaluating \mathcal{L}_ϖ at the origin 0_σ of σ gives a line, so a line bundle \mathcal{M}_ϖ on $\tilde{\mathcal{G}}_\varepsilon^{ss}$ with $\mathcal{M}_\varpi(P) = \mathcal{L}_\varpi(0_\sigma)$. This gives the embedding that was asserted. \square

Pull back to $\text{Pic}(\tilde{\mathcal{Z}}^- - \sigma^{-1}(0_{\mathcal{L}}))$ via the morphism $\tilde{\mathcal{Z}}^- - \sigma^{-1}(0_{\mathcal{L}}) \rightarrow \tilde{\mathcal{G}}^{ss}$; we get a homomorphism $\lambda : \text{Pic}_F^G \rightarrow \text{Pic}(\tilde{\mathcal{Z}}^- - \sigma^{-1}(0_{\mathcal{L}}))$. Now fix an elliptic curve E over a field k and return to the restriction $\tilde{X}^- \rightarrow \mathbb{A}_k^1$ of $\tilde{\mathcal{Z}}^-$ to the line \mathbb{A}_k^1 in L over the origin 0_Y . For $t \neq 0$, the fibre \tilde{X}_t^- contains a (-2) -configuration Δ_t of type $G = E_i$; since $\tilde{\mathcal{Z}}^- \cong N_{D_1^-/\tilde{\mathcal{Z}}^-}$, Δ_t specializes to an isomorphic such configuration Δ_0 on D_1^- .

Regard \tilde{X}_t^- as a surface contained in $\tilde{\mathcal{Z}}^- - \sigma^{-1}(0_{\mathcal{L}})$. By Theorem 3.8, Δ_t is identified with a subregular unipotent Springer fibre associated to G and, for any $\varpi \in \text{Pic}_F^G$, $\lambda(\varpi)$ is identified with ϖ .

We are led to the following result.

Proposition 7.6 *There are subgroups \tilde{H} of $\text{Pic}_{\tilde{\mathcal{Z}}}$ and H^- of $\text{Pic}_{\tilde{\mathcal{Z}}^-}$ and a commutative diagram with exact rows*

$$\begin{array}{ccccccc} \mathbb{Z}\{[\mathcal{D}_0], [\mathcal{D}_1], [\mathcal{Q}]\} & \longrightarrow & \tilde{H} & \longrightarrow & \text{Pic}_F^G & \longrightarrow & 0 \\ & & \downarrow & & \downarrow = & & \\ \mathbb{Z}\{[\mathcal{D}_0^-], [\mathcal{D}_1^-]\} & \longrightarrow & H^- & \longrightarrow & \text{Pic}_F^G & \longrightarrow & 0 \end{array}$$

where the two leftmost vertical arrows are induced by the blowing-down $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{Z}}^+$ and the flop $\tilde{\mathcal{Z}}^+ \rightarrow \tilde{\mathcal{Z}}^-$. In particular, $[\mathcal{Q}] \mapsto 0$.

PROOF: The only remaining point is to check that the obvious homomorphisms $\tilde{H} \rightarrow \text{Pic}_{\tilde{\mathcal{Z}}}$ and $H^- \rightarrow \text{Pic}_{\tilde{\mathcal{Z}}^-}$ are injective. However, this follows from restricting to one of the surfaces \tilde{Z}_t , as just described. \square

Restrict from $\tilde{\mathcal{Z}}^-$ to \mathcal{D}_1^- . We get a subgroup $H \subset \text{Pic}_{\mathcal{D}_1^-}$ and, since $\mathcal{A} = \mathcal{D}_0^-|_{\mathcal{D}_1^-} \sim \omega_{\mathcal{D}_1^-/\mathcal{Y}}^{-1}$, an exact sequence

$$\mathbb{Z}[\mathcal{A}] \oplus \mathbb{Z}[\gamma] \longrightarrow H \xrightarrow{\chi} \text{Pic}_F^G \longrightarrow 0$$

such that, for any $\Lambda \in H$ and component δ_i of Δ , $(\Lambda.\delta_i)_{\mathcal{D}_1^-} = (\chi(\Lambda).\alpha_i^\vee)$, where α_i^\vee is the simple coroot corresponding to δ_i .

Corollary 7.7 *The natural homomorphism $H \rightarrow \text{NS}(\mathcal{D}_1^-/\mathcal{Y})$ of sheaves of commutative groups on \mathcal{Y} is surjective, and $\text{NS}(\mathcal{D}_1^-/\mathcal{Y})$ is constant.*

PROOF: It is enough to prove surjectivity for one (geometric) weak del Pezzo surface D_1^- . Choose D_1^- lying over 0_Y ; this surface contains a configuration Δ_0

of type G and a line γ , and the result now follows from Proposition 6.30 and the fact that \mathcal{Y} is simply connected. \square

Take D_1^- as in the proof just given. Let $I_{1,l}$ denote the \mathbb{Z} -lattice with \mathbb{Z} -basis $\{\delta_1, \dots, \delta_l, \gamma\}$ and inner product given by the intersection numbers on D_1^- ; then $I_{1,l}$ is isomorphic to the standard odd unimodular hyperbolic lattice of rank $1+l$ and the basis just given describes an isometry $\phi : I_{1,l} \rightarrow \text{NS}(D_1^-)$. Under ϕ^{-1} the exceptional curves of the blow-up $D_1 \rightarrow \mathbb{P}^2$ described in Proposition 6.30 are $\gamma, \gamma + \delta_l, \dots, \gamma + \delta_l + \dots + \delta_3$ and $\gamma + \delta_l + \dots + \delta_3 + \delta_1$, while δ_2 is the strict transform of a line in \mathbb{P}^2 .

Since $\text{NS}(\mathcal{D}_1^-/\mathcal{Y})$ is constant and $\text{Pic}(\mathcal{D}_1^-/\mathcal{Y}) \rightarrow \text{NS}(\mathcal{D}_1^-/\mathcal{Y})$ is an isomorphism, ϕ extends to an isometry $\phi : I_{1,l} \rightarrow \text{Pic}(\mathcal{D}_1^-/\mathcal{Y})$. Also, \mathcal{D}_1^- contains an anti-canonical divisor \mathcal{A} that is a copy of $\mathcal{E} \times_{\mathcal{M}_{EII}} \mathcal{Y}$; the isomorphism $\mathcal{A} \rightarrow \mathcal{E} \times_{\mathcal{M}_{EII}} \mathcal{Y}$ is provided by the base point $\gamma \cap \mathcal{A}$.

Restricting to \mathcal{A} provides a homomorphism

$$\psi : I_{1,l} \rightarrow \text{Pic}(\mathcal{A}/\mathcal{Y}) \cong \text{Pic}(\mathcal{E}) = \coprod_{n \in \mathbb{Z}} \text{Pic}^n(\mathcal{E}).$$

Define $\kappa_l = \phi^{-1}[\omega_{\mathcal{D}_1^-/\mathcal{Y}}^{-1}]$, the anti-canonical class. Then

$$\kappa_l = \delta_1 + \delta_2 + 2\delta_3 + 3 \sum_{i \geq 4} \delta_i + 3\gamma.$$

Note that $I_{1,l}/\mathbb{Z}\kappa_l$ is the weight lattice P , while $\oplus \mathbb{Z}\delta_i$ is the root lattice Q . Consider the positive cone C_+ in $I_{1,l} \otimes \mathbb{R}$ defined by the conditions $(v, v) \geq 0, v \cdot \kappa_l \geq 0$; this inherits a finite decomposition into chambers from the decomposition of $P \otimes \mathbb{R}$ into fundamental domains for the W -action. One such is the chamber C_0 defined by the inequalities $(v, \delta_i) \geq 0$ for all i .

Lemma 7.8 $\omega_{\mathcal{D}_1^-/\mathcal{Y}}^{-1}|_{\mathcal{A}}$ is isomorphic to the pullback of $\mathcal{O}_E(d[O_E])$ to $\mathcal{A} = \mathcal{E} \times_{\mathcal{M}_{EII}} \mathcal{Y}$.

PROOF: \mathcal{D}_0^- is the minimal resolution of a singular normal surface that is defined over the stack \mathcal{M}_{EII} of elliptic curves, and so $\mathcal{N}_{\mathcal{A}/\mathcal{D}_0^-}$ is defined over \mathcal{M}_{EII} . So $\mathcal{N}_{\mathcal{A}/\mathcal{D}_0^-} \cong \mathcal{O}_E(-d[O_E])$. Since $\mathcal{N}_{\mathcal{A}/\mathcal{D}_0^-} \cong \mathcal{N}_{\mathcal{A}/\mathcal{D}_1^-}^\vee$, the result follows from the adjunction formula and the triviality of $\omega_{\mathcal{Z}^-/\mathcal{L}}$. \square

By Lemma 7.8, $\psi(\kappa_l) = d[O_E]$, so there is a homomorphism $\bar{\psi} : P \rightarrow \text{Pic}_E^0$ defined by $\bar{\psi}(L) = \psi(L) - \deg \psi(L)[O_E]$.

Now consider the stack \mathcal{MdP}_d of *marked* weak del Pezzo surfaces of degree d ; by definition, the objects consist of:

- (1) a family $f : X \rightarrow S$ of weak del Pezzo surfaces of degree d ;
- (2) an embedding $i : E \times S \hookrightarrow X$ such that the class of the image \mathcal{A} equals the class $[\omega_{X/S}^{-1}]$ and $i^*(\omega_{X/S}^{-1})$ is linearly equivalent to $d[O_E]$;

(3) an isometry $\phi : I_{1,l} \rightarrow \text{Pic}(X/S)$ such that $\phi(\gamma)|_{\mathcal{A}}$ is linearly equivalent to $[0_{\mathcal{A}}]$ and $\phi(\delta_j)$ has degree 0 for all j .

In this language, $(\mathcal{D}_1^- \rightarrow \mathcal{Y}, \mathcal{A} \hookrightarrow \mathcal{D}_1^-, \phi)$ is an object of $\mathcal{M}d\mathcal{P}_d$, so defines a morphism $H : \mathcal{Y} \rightarrow \mathcal{M}d\mathcal{P}_d$.

There is also a morphism $F : \mathcal{M}d\mathcal{P} \rightarrow \mathcal{Y}$ defined by $(f, i, \phi) \mapsto \psi'$, where ψ' is constructed exactly as above, and a morphism $G : \mathcal{Y} \rightarrow \mathcal{M}d\mathcal{P}$, as follows:

Given $\psi' : P \rightarrow E$, construct $\psi : I_{1,l} \rightarrow \text{Pic}(E)$ by

$$\psi(\gamma) = [0_E], \quad \psi(\delta_i) = \psi'(\delta_i) \pmod{\mathbb{Z}\kappa_l}.$$

Then $\psi(\kappa_l)$ has degree 3. Embed $E \hookrightarrow \mathbb{P}^2$ via $|\psi(\kappa_l)|$ and then make l blow-ups \mathbb{P}^2 along the points on E defined by the degree 1 classes

$$\psi(\gamma + \delta_l + \cdots + \delta_3 + \delta_1), \psi(\gamma + \delta_l + \cdots + \delta_3), \dots, \psi(\gamma + \delta_l), \psi(\gamma)$$

in that order.

The next result is due to Mérindol [M]. Its proof is an immediate consequence of the definitions of F and G .

Proposition 7.9 $F \circ G = 1_{\mathcal{Y}}$.

Corollary 7.10 F exhibits \mathcal{Y} as the maximal separated quotient of $\mathcal{M}d\mathcal{P}$.

PROOF: Immediate. □

That is, $\mathcal{M}d\mathcal{P}$ is obtained by glueing together copies of \mathcal{Y} along open subvarieties. On the locus of del Pezzo surfaces that have no (-2) -curves the stacks $\mathcal{M}d\mathcal{P}$ and \mathcal{Y} are isomorphic but $\mathcal{M}d\mathcal{P}$ fails to be separated when (-2) curves appear.

Proposition 7.11 $F \circ H = 1_{\mathcal{Y}}$.

PROOF: $F \circ H$ takes $0_{\mathcal{Y}}$ to $0_{\mathcal{Y}}$, so is a homomorphism of abelian schemes. Fix a geometric elliptic curve E ; then over $0_{\mathcal{Y}}$ the fibre of \mathcal{D}_1 contains a (-2) -configuration of type G . Consider the subdiagram

$$\begin{array}{ccccc} \tilde{\mathcal{Z}}^- & \xrightarrow{\sigma} & \mathcal{L} & \longrightarrow & \mathcal{Y} \\ \downarrow \epsilon & & \downarrow & & \\ \mathcal{Z} & \longrightarrow & [\widehat{\mathcal{Y}}/W] & & \end{array}$$

of the diagram in Proposition 7.2. It follows from Theorem 6.24 that $\mathcal{Z} \rightarrow [\widehat{\mathcal{Y}}/W]$, which is a family of affine surfaces, has du Val singularities of type G exactly over $\mathbb{A}^1 - \{0\}$, where \mathbb{A}^1 is the line in the cone $[\widehat{\mathcal{Y}}/W]$ that lies over the image of $0_{\mathcal{Y}}$ in $[\widehat{\mathcal{Y}}/W]$. Therefore $0_{\mathcal{Y}}$ is isolated in the fibre $(F \circ H)^{-1}(0_{\mathcal{Y}})$ and then $F \circ H$ is an isomorphism. Since $F \circ H$ is W -equivariant it is then ± 1 .

Suppose that $F \circ H = -1$. Put $S = \mathcal{U} \times_{\mathcal{M}d\mathcal{P}, G} \mathcal{Y}$, where $\mathcal{U} \rightarrow \mathcal{M}d\mathcal{P}$ is universal, and consider the \mathcal{Y} -isomorphism $\mathcal{D}_1 \rightarrow S \times_{\mathcal{Y}, F \circ H} \mathcal{Y}$. Since $F \circ H = -1$,

this takes the ample cone on \mathcal{D}_1 to the negative of the ample cone on S . This is impossible, so $F \circ H = 1$. \square

The modular interpretation of \mathcal{Y} as an open substack of \mathcal{MdP} is given as follows.

Suppose that X is a weak del Pezzo surface. Define a *line* in X to be a class m in $\text{NS}(X)$ such that $m^2 = -1$ and $m \cdot (-K_X) = 1$ and a *root* in X to be a class δ in $\text{NS}(X)$ such that $\delta^2 = -2$ and $\delta \cdot K_X = 0$. Let Λ denote the set of lines and Δ the set of roots on X . Then Δ is a finite root system and reflections in the roots generate a finite Weyl group W , as usual. A root δ is *effective* if $H^0(X, \delta) \neq 0$.

Define the *positive cone* $C^+ = C^+(X)$ by $C^+ = \{x \in \text{NS}(X)_{\mathbb{R}} \mid x^2 > 0, x \cdot (-K_X) > 0\}$. It is well known that a class x in $C^+ \cap \text{NS}(X)$ is nef if and only if $x \cdot m > 0$ for all $m \in \Lambda$ and $x \cdot \delta \geq 0$ for all effective roots δ on X . Put $\mathcal{F} = \{x \in C^+ \mid x \cdot m > 0 \forall m \in \Lambda\}$. Then the roots define walls that tessellate \mathcal{F} into chambers that are permuted simply transitively by W . So, if \mathcal{F}' is one such chamber, then $\mathcal{F} = \cup_{w \in W} w(\mathcal{F}')$.

In the lattice $I_{1,l}$, say $\Lambda_0 = \{m \in I_{1,l} \mid m^2 = -1, m \cdot \kappa = 1\}$, $\Delta_0 = \{\delta \in I_{1,l} \mid \delta^2 = -2, \delta \cdot \kappa = 0\}$, $C_0^+ = \{x \in I_{1,l} \otimes \mathbb{R} \mid x^2 > 0, x \cdot \kappa > 0\}$ and $\mathcal{F}_0 = \{x \in C_0^+ \mid x \cdot m > 0 \forall m \in \Lambda_0\}$.

Pick a chamber \mathcal{F}'_0 in the tessellation of \mathcal{F}_0 defined by the roots in Δ_0 . Let \mathcal{MdP}^+ be the open substack of \mathcal{MdP} whose objects are triples $(X \rightarrow S, i, \phi)$ such that $\phi_{\mathbb{R}}(\mathcal{F}'_0)$ is contained in the nef cone of every geometric fibre of $X \rightarrow S$. Then \mathcal{MdP}^+ is isomorphic to \mathcal{Y} , and \mathcal{MdP} is the union of these copies of \mathcal{Y} , one for each chamber \mathcal{F}'_0 in \mathcal{F}_0 .

This brings us to the main result of the paper.

Theorem 7.12 *Suppose that $G = E_l$ for $l = 5, 6, 7, 8$.*

(1) *The family $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$ has simply elliptic singularities over the vertex of the cone $[\hat{\mathcal{Y}}/W]$.*

(2) *Over $\mathbf{Spec} \mathbb{Z}[1/(9-l)]$ this family is a miniversal deformation of these singularities.*

(3) *The commutative diagram*

$$\begin{array}{ccc} \tilde{\mathcal{Z}}^- & \longrightarrow & \mathcal{Z} \\ \downarrow & & \downarrow \\ \mathcal{L} & \longrightarrow & [\hat{\mathcal{Y}}/W] \end{array}$$

is a simultaneous log resolution of $\mathcal{Z} \rightarrow [\hat{\mathcal{Y}}/W]$.

(4) *The exceptional divisor \mathcal{D}_1^- in $\tilde{\mathcal{Z}}^-$ is the restriction to \mathcal{Y} of the universal marked del Pezzo surface of degree $d = 9 - l$.*

PROOF: (1)-(3) summarize the results of Section 6. (4) follows at once from Corollary 7.10 and the construction of $\tilde{\mathcal{Z}}^-$. \square

Remark: Suppose that the base is $\mathbf{Spec} \mathbb{C}$. The affine del Pezzo surface obtained by deleting the elliptic curve E has a mixed Hodge structure on H^2 . These mixed Hodge structures are naturally parametrized by the Looijenga variety \mathcal{Y} . The above theorem shows, when restricted to the exceptional divisor \mathcal{D}_1^- in \mathcal{Z}^- , that the period map for these affine del Pezzo surfaces is the same as the map from \mathcal{D}_1^- to \mathcal{Y} that we have constructed in terms of group theory.

8 del Pezzo surfaces and the unipotent singularity of E_8 in characteristics 2, 3 and 5

Consider again the family $\tilde{X}^- \rightarrow \mathbb{A}^1$ where \mathbb{A}^1 is the line in the line bundle $L \rightarrow Y$ lying over 0_Y . The closed fibre of \tilde{X}_0^- is $\tilde{X}_0^- = D_0^- + D_1^-$ and for $t \neq 0$ the fibre $\tilde{X}_t^- = \tilde{X}_t$ contains a configuration Γ_t of (-2) -curves of the same combinatorial type as G . When $t = 0$ this configuration specializes to an isomorphic configuration Γ_0 which lies in D_1^- and is disjoint from the double curve $A^- = D_0^- \cap D_1^-$. Recall that A^- is isomorphic to E .

Let $D_1^- \rightarrow D_1^b$ be the contraction of Γ_0 . Then D_1^b has a singularity of combinatorial type G that does not lie on the curve A^b which is the (isomorphic) image of A^- .

Proposition 8.1 (1) \tilde{X}^- is isomorphic to the line bundle $\omega_{D_1^-}$.

(2) There is a commutative diagram

$$\begin{array}{ccc} \tilde{X}^- & \longrightarrow & \tilde{X}^b \\ & \searrow & \downarrow \\ & & \mathbb{A}_k^1 \end{array}$$

where $\tilde{X}^- \rightarrow \tilde{X}^b$ contracts each Γ_t to a du Val singularity, $\cup_t \Gamma_t$ is contracted to a section of $\tilde{X}^b \rightarrow \mathbb{A}_k^1$ and $\tilde{X}_0^b = D_0^b + D_1^b$ where $D_0^+ \rightarrow D_0^b$ is an isomorphism.

(3) \tilde{X}^b is isomorphic to the line bundle $\omega_{D_1^b}$.

(4) The singular affine surface \tilde{X}_t is isomorphic to the affine del Pezzo surface $D_1^b - A^b$ for all $t \neq 0$.

PROOF: The existence of the contraction $X^+ \rightarrow X^b$ is a special case of the well known fact that a family of (-2) -configurations lying in the relative smooth locus of a family of surfaces can be simultaneously contracted. The rest follows from Lemma 6.28. \square

As has been already recalled, if $Uni(G)$ is the unipotent variety of a simply connected simple group G of type A , D or E , then the singular locus of $Uni(G)$ is the subregular locus of $Uni(G)$ and the strict localization of $Uni(G)$ at the geometric

generic point of its subregular locus (we shall refer to this as the unipotent singularity) has a du Val singularity of the same combinatorial type as G . However, in low characteristics the combinatorial type does not specify the singularity, even over an algebraically closed field; for example, the E_8 singularities are classified up to formal isomorphism on p. 270 of [Li] (and up to henselian equivalence in [Ar]). The local equation of each is

$$t^2 + z^3 + y^5 = \phi$$

where $\phi \in \{0, y^3t, y^3zt, y^2zt, yzt\}$ in characteristic 2, $\phi \in \{0, y^3z^2, y^2z^2\}$ in characteristic 3 and $\phi \in \{0, y^4z\}$ in characteristic 5. We refer to them as $E_8^{(p,\phi)}$ accordingly. In Artin's list [Ar] they are listed instead as E_8^r where $0 \leq r \leq 4$ or 2 or 1 in these characteristics and $\dim T^1 = 16 - 2r$ or $12 - 2r$ or $10 - 2r$.

For the rest of this section the base field k will be a complete valued algebraically closed field of characteristic $p > 0$ and E will be a uniformizable elliptic curve over k .

Theorem 8.2 *The singularity on D_1^b is isomorphic to the unipotent singularity of the group G .*

PROOF: By Theorem 3.11 the singularity of $X_{t \neq 0}$ is isomorphic to the unipotent singularity. The theorem follows from Proposition 8.1. \square

Recall that in characteristic p the j -invariant of a uniformizable elliptic curve E is not algebraic over \mathbb{F}_p ; in particular, it is non-zero.

Until further notice the group G will be E_8 .

Recall that the anticanonical model of a del Pezzo surface S of degree 1 over a field k is a sextic hypersurface in weighted projective space $\mathbb{P}(1, 1, 2, 3) = \text{Proj } k[X_1, Y_1, Z_2, T_3]$.

The next result is a counterpart to Proposition 6.30.

Proposition 8.3 *Suppose that S is a del Pezzo surface of degree 1 over a separably closed field k with a singular point P of type E_8 .*

(1) *Up to projective equivalence, the defining equation of S in $\mathbb{P}(1, 1, 2, 3)$ is*

$$T^2 + Z^3 + XY^5 = \Phi$$

where $\Phi \in \{ZY^4, 0\}$ if $\text{char } k \neq 2, 3$, $\Phi \in \{Z^2Y^2, ZY^4, 0\}$ if $\text{char } k = 3$ and $\Phi \in \{TYZ, TY^3, \gamma ZY^4\}$ if $\text{char } k = 2$, where $\gamma = \lambda^4 + c$, so defines a class in $H_{flat}^1(k, \mu_4)$.

(2) *Suppose that C is a smooth member of $|-K_S|$. Then, whatever the characteristic, $j(C)$ is never zero for the first value of Φ and is always zero for the second. (In characteristics 2 and 3 the pencil $|-K_S|$ is quasi-elliptic for the third value, namely, when $\Phi = 0$ or γZY^4 .)*

(3) *In the first case of each characteristic the singularity (S, P) is isomorphic to $E_8^{(5, y^4z)} = E_8^1$ or $E_8^{(3, y^2z^2)} = E_8^2$ or $E_8^{(2, yzt)} = E_8^4$.*

PROOF: This depends upon the following recognition principle for E_8 singularities. It goes back to du Val.


Lemma 8.4 (1) *A du Val singularity (S, P) is of combinatorial type E_8 if and only if the blow-up $\tilde{S} = \text{Bl}_P S$ has a singular point of type E .*

(2) *A du Val singularity whose defining equation can be written as*

$$t^2 - G(y, z) \in t(y, z, t)^2$$

is of type E if and only if the leading term of G is proportional to a perfect cube.

PROOF: (1) Suppose that $S^* \rightarrow S$ is the minimal resolution. Then, according to du Val and Artin, this factors through \tilde{S} and the fundamental cycle Z on S^* is a divisor and the sheaf of ideals that defines it is $\mathfrak{m}_P \cdot \mathcal{O}_{S^*}$. So, for any component C_i of the exceptional locus in S^* , $Z \cdot C_i \leq 0$, and $Z \cdot C_i < 0$ if and only if C_i appears on \tilde{S} . Moreover, Z is the biggest root; in the case of E_8 this is the following linear combination of simple roots:



Therefore the curves contracted by $S^* \rightarrow \tilde{S}$ form an E_7 configuration. Similarly an E_7 (or E_6) singularity has a D_6 (or A_5) singularity on its first blow-up.

(2) This is to be found on p. 268 of [Li]. □

Beyond this lemma the proof of Proposition 8.3 consists of a straightforward manipulation of polynomials and a calculation of j -invariants using the standard formulae to be found on, for example, p. 46 of [Si]. We omit the details. □

Remark: In particular, over an algebraically closed field of characteristic 2 there are five E_8 singularities but only three of them can lie on a del Pezzo surface. We do not know what to make of this.

Corollary 8.5 *The unipotent singularity of the group E_8 in characteristics 2, 3 and 5 is isomorphic to $E_8^{(2,yzt)} = E_8^4$, $E_8^{(3,y^2z^2)} = E_8^2$ and $E_8^{(5,y^4z)} = E_8^1$, respectively.*

PROOF: The unipotent singularity is realized on a del Pezzo surface that contains a uniformizable curve. The j -invariant of a uniformizable curve E is transcendental over \mathbb{F}_p , so that in particular $j(E) \neq 0$, and we can now apply Proposition 8.3. □

Corollary 8.6 *The BGSS construction gives a versal deformation of the unipotent singularity of E_8 in all characteristics.*

PROOF: For each du Val singularity Artin computed [Ar] the dimension of T^1 , the tangent space to a miniversal deformation space of the singularity. From his

list and Corollary 8.5 we can see that $\dim T^1 = 8$ for the unipotent singularity of E_8 . Now the argument of [SB01] goes through unchanged, as it does whenever $\dim T^1$ equals the rank of the group. \square

Corollary 8.7 *If E is supersingular, $G = E_8$ and $\text{char } k = 2, 3$ or 5 then $\tilde{\mathcal{G}}^{ss} \rightarrow \mathcal{G}^{ss}$ is not smoothly equivalent to $\tilde{G} \rightarrow G$ in any neighbourhood of the identity of G . That is, the conclusion of Theorem 3.11 fails in these cases.*

PROOF: If it were true, then the morphism $\tilde{\mathcal{G}}^{ss} \rightarrow \mathcal{G}^{ss}$ would be smoothly equivalent to $\tilde{G} \rightarrow G$ and then the unipotent singularity of E_8 would be realized on a del Pezzo surface S on which E is an anticanonical curve. Then, according to Proposition 8.3, S can be defined by the equation

$$T^2 + Z^3 + XY^5 = \Phi$$

where $\Phi = 0, ZY^4$ or TY^3 when $\text{char } k = 5, 3$ or 2 , respectively, since in these characteristics an elliptic curve is supersingular if and only if its j -invariant is 0 . But then S has a singularity of type $E_8^{(5,0)}$ or $E_8^{(3,y^4z)}$ or $E_8^{(2,y^3t)}$ and it remains only to check that $E_8^{(3,y^4z)}$ is not isomorphic to $E_8^{(3,y^2z^2)}$. This can be done by calculating $\dim T^1$ for $E_8^{(3,y^4z)}$; the result is $\dim T^1 = 10$, so that inspection of Artin's list [Ar] shows that $E_8^{(3,y^4z)}$ is in fact isomorphic to $E_8^{(3,y^3z^2)}$. \square

For E_6 and E_7 similar considerations involving del Pezzo surfaces of degrees 3 and 2 work to describe the unipotent singularity in low characteristics: in the notation of [Ar] they are E_6^1 and E_7^3 when $p = 2$ and E_6^1 and E_7^1 when $p = 3$. In all cases the unipotent singularity is the one in its combinatorial class where $\dim T^1$ is minimal, so is the “most general” one.

References

- [Ar] M. Artin, *Coverings of rational double points in characteristic p* , in *Complex Analysis*, W. Baily and T. Shioda, eds.
- [AS] J. Armstrong and N. Shepherd-Barron, *A_g^P has terminal singularities for $g \geq 6$* , arXiv
- [BG] A. Braverman and D. Gaitsgory, *Geometric Eisenstein Series*, arXiv:math/9912097v2
- [BH] H. Bass and W. Haboush, *Linearizing certain reductive group actions*, Trans. AMS **292** (1985), 463–482.
- [BGR] S. Bosch, U. Güntzer and R. Remmert, *Non-Archimedean Analysis*, Springer, 1984.

- [Br] E. Brieskorn, *On the unfolding of exceptional singularities*, Nova Acta Leopoldina **52**, 1981.
- [Brue] G. Bruechert, *Trace class elements and cross-sections on Kac-Moody groups*, Canad. J. Math. **50** (1998), 972–1006.
- [D] D. Davis, thesis, in preparation.
- [FM] R. Friedman and J. Morgan, *Holomorphic Principal Bundles Over Elliptic Curves II: The Parabolic Construction*, J. Diff. Geom. **56** (2000), 301–379.
- [GS] I. Grojnowski and N. Shepherd-Barron, *Springer fibres revisited*, in preparation.
- [HS1] S. Helmke and P. Slodowy, *Loop groups, elliptic singularities and principal bundles over elliptic curves*, Banach Center Publications, Volume 62, 2004.
- [HS2] ———, *On Unstable Principal Bundles over Elliptic Curves*, Publ. RIMS, Kyoto Univ. **37** (2001), 349–395
- [HS3] ———, *Singular elements of affine Kac-Moody groups*, Proc. ECM Stockholm (2004).
- [Hi] M. Hirokado, *Deformations of rational double points and simple elliptic singularities in characteristic p* , Osaka J. Math **41** (2004), 605–616.
- [Li] J. Lipman, *Rational singularities with applications to algebraic surfaces and unique factorization*, Publ. Math. IHES **36** (1969), 195–279.
- [L] E. Looijenga, *Homogeneous spaces associated to certain semi-universal deformations*, Proc. ICM (Helsinki, 1978), pp. 529–536.
- [L1] ———, *Root systems and elliptic curves*, Invent. Math. **38** (1976), 17–32.
- [L2] ———, *Semi-universal deformation of a simple elliptic singularity, II*, Topology **17** (1978), 23–40.
- [M] J.Y. Mérindol, *Les singularités simples elliptiques, leurs déformations, les surfaces de del Pezzo et les transformations quadratiques*, Ann. Sci. ENS **15** (1982), 17–44.
- [MOSW] R. Muñoz, G. Occhetta, L.E. Solá Conde and K. Watanabe, *Rational curves, Dynkin diagrams and Fano manifolds with nef tangent bundle*, Math. Ann. **361** (2015), 583–609.

- [P] H. Pinkham, *Simple elliptic singularities, Del Pezzo surfaces and Cremona transformations*, Proc. Sympos. Pure Math., Vol. XXX, Part 1, 69–71.
- [SGA3] *Schémas en groupes*, Springer Lecture Notes **151-3**
- [SB01] N.I. Shepherd-Barron, *On simple groups and simple singularities*, Israel J. Math. **123** (2001), 179–188.
- [SB16] ———, *An exceptional locus in the perfect compactification of A_g* , arXiv:
- [Si] J. Silverman, *The arithmetic of elliptic curves*, Springer, 1986.
- [Sl] P. Slodowy, *Simple singularities and simple algebraic groups*, Springer Lecture Notes **815**.
- [SSS] *Séminaire sur les singularités des surfaces*, Springer Lecture Notes **777**.

D.P.M.M.S., CAMBRIDGE UNIVERSITY, CAMBRIDGE CB2 1SB, U.K.
E-mail address: groj@dpms.cam.ac.uk

KING'S COLLEGE LONDON, STRAND, LONDON WC2R 2LS, U.K.
E-mail address: Nicholas.Shepherd-Barron@kcl.ac.uk