



King's Research Portal

DOI:

[10.1017/S0033291719001983](https://doi.org/10.1017/S0033291719001983)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Crush, E., Arseneault, L., Danese, A., Jaffee, S. R., & Fisher, H. (2019). Using discordant twin methods to investigate an environmentally mediated pathway between social support and the reduced likelihood of adolescent psychotic experiences. *Psychological Medicine*, 1898-1905. Advance online publication. <https://doi.org/10.1017/S0033291719001983>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Original Article

Cite this article: Crush E, Arseneault L, Danese A, Jaffee SR, Fisher HL (2019). Using discordant twin methods to investigate an environmentally mediated pathway between social support and the reduced likelihood of adolescent psychotic experiences. *Psychological Medicine* 1–8. <https://doi.org/10.1017/S0033291719001983>

Received: 13 November 2018

Revised: 22 March 2019

Accepted: 16 July 2019

Key words:


genetic risk; psychosis; resilience; trauma; twins; victimisation

Author for correspondence:

Helen Fisher, E-mail: helen.2.fisher@kcl.ac.uk

Using discordant twin methods to investigate an environmentally mediated pathway between social support and the reduced likelihood of adolescent psychotic experiences

Eloise Crush¹, Louise Arseneault¹, Andrea Danese^{1–3}, Sara R. Jaffee⁴ and

Helen L. Fisher¹ 

¹King's College London, Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, London, UK; ²Department of Child & Adolescent Psychiatry, King's College London, Institute of Psychiatry, Psychology & Neuroscience, London, UK; ³National & Specialist CAMHS Clinic for Trauma, Anxiety, and Depression, South London and Maudsley NHS Foundation Trust, London, UK and ⁴Department of Psychology, University of Pennsylvania, Philadelphia, PA, USA

Abstract

Background. Social support has been shown to be associated with a reduced likelihood of developing psychotic experiences in the general population and even amongst those at high risk due to exposure to multiple forms of victimisation (poly-victimised). However, it is unclear whether this association is merely due to the confounding effects of shared environmental and genetic influences, or reverse causality. Therefore, we investigated whether social support has a unique environmentally mediated effect on adolescent psychotic experiences after accounting for familial factors, including genetic factors, and also prior psychopathology. **Methods.** Participants were from the Environmental Risk (E-Risk) Longitudinal Twin Study, a nationally-representative cohort of 2232 UK-born twins. Adolescents were interviewed at age 18 about psychotic experiences and victimisation exposure since age 12, and their perceptions of social support. Prior childhood mental health problems and psychotic symptoms were assessed at age 12. The discordant twin method was used to disentangle the relative family-wide and unique-environmental effects of social support on psychotic experiences in the general population and among poly-victimised adolescents.

Results. Perceived social support, particularly from friends, was found to have a unique environmentally mediated buffering effect on adolescent psychotic experiences in the whole sample and in the high-risk poly-victimised group.

Conclusions. The protective effects of social support on adolescent psychotic experiences cannot be accounted for by shared environmental or genetic factors, nor by earlier psychopathology. Our findings suggest that early intervention programmes focused on increasing perceptions of social support have the potential to prevent the emergence of psychotic experiences amongst adolescents.

Introduction

It is widely acknowledged that psychotic experiences such as hearing voices and feeling very paranoid, occur amongst individuals in the general population (McGrath *et al.*, 2015). They are relatively common, with prevalence rates around 17% during childhood and 8% during adolescence (Kelleher *et al.*, 2012a). Poly-victimisation (exposure to two or more types of victimisation) has been found to be a major risk factor for the emergence of psychotic phenomena with the odds of reporting such phenomena around five times higher than in the general population (Arseneault *et al.*, 2011; Crush *et al.*, 2018a). However, a large proportion of children and adolescents in the general population will not develop psychotic phenomena, even in the context of poly-victimisation (Janssen *et al.*, 2004; Shevlin *et al.*, 2008; Arseneault *et al.*, 2011). Research focused on those who do not develop psychotic phenomena, despite being at high risk (poly-victimised), may provide valuable insights into which factors reduce the likelihood of psychotic experiences emerging and thus inform preventive interventions. Preventing early psychotic phenomena is crucial not only because these experiences are extremely distressing for adolescents (Kelleher *et al.*, 2015) but also because they have been shown to predict suicidal behaviours (Kelleher *et al.*, 2012b) and major psychiatric disorders (Fisher *et al.*, 2013) in later life.

In a recent paper, we found social support particularly from friends and family to be protective in relation to age-18 psychotic experiences in the general population and amongst a high-risk group of poly-victimised adolescents after controlling for gender, family

© The Author(s) 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

socioeconomic status, family psychiatric history, and childhood mental health problems including psychotic symptoms (Crush *et al.*, 2018a). We concluded that these findings provide evidence for an independent protective effect of adolescent social support in relation to psychotic phenomena in a nationally-representative longitudinal cohort study. This paper aims to further interrogate the potentially causal nature of this association by taking advantage of our longitudinal twin sample to control for all unmeasured shared family-wide environmental and genetic factors that could be confounding this association, and also accounting for earlier mental health problems to limit the possibility of reverse causality.

An association which may be causal in nature would be inferred if social support was found to have a direct, environmentally-mediated protective effect on adolescent psychotic phenomena. An environmentally-mediated pathway would suggest that, to some extent, the effect of perceived social support on psychotic phenomena is distinctly environmental in nature. Ultimately, evidence for this would indicate that interventions to increase social support, or perceptions of social support, could be effective in preventing the development of psychotic phenomena in adolescence. In order to investigate this, it is necessary to limit the possibility of alternative explanations, including potential confounding by shared family-wide environmental and genetic factors, and also considering whether reverse causality is operating, all of which are considered herein and discussed below.

There are several non-causal or indirect explanations for why higher levels of social support are associated with a reduced likelihood of developing psychotic experiences during adolescence. Firstly, environments or experiences shared by family members, including the home and community environment could influence perceptions of social support and also protect against the onset of psychotic phenomena. For example, children who grow up in a warm, nurturing home environment are less likely to report psychotic phenomena (Crush *et al.*, 2018b) and may also perceive others to be more supportive, perhaps due to secure attachment formation (Blain *et al.*, 1993). Similarly, individuals who grow up in a neighbourhood with high social cohesion may have more access to social support within their community, while low neighbourhood social cohesion has been found to be associated with the emergence of psychotic phenomena (Newbury *et al.*, 2016). Therefore, shared family-wide environmental factors might be confounding the protective effect of social support on psychotic experiences.

Secondly, genetic factors may also explain the association by influencing both perceptions of social support and the propensity to develop psychotic experiences. Indeed, there are modest to high heritability estimates for the emergence of psychotic phenomena during childhood and adolescence (Polanczyk *et al.*, 2010; Zavos *et al.*, 2014). Additionally, despite intuitively social support being considered an environmental exposure, it is also influenced by genetic factors (Kendler, 1997), with a moderate heritability of 40% found in the current cohort (Matthews *et al.*, 2016). Given that social support and psychotic phenomena are both influenced by genetic factors, it is possible that genes may confound the association between them. For example, it is plausible that individuals with paranoia or suspicious thoughts, that may arise from a genetic predisposition towards psychotic experiences, could have problems maintaining relationships with friends (Claes, 1994) and family (Riggio and Kwong, 2011), and be less appealing to new potential friends or partners. Conversely, those without such genetic vulnerability may be more likely to elicit social support and also be protected from developing psychotic phenomena.

Relatedly, the potential for reverse causality should be considered in the association between social support and psychotic phenomena. It is possible that early manifestations of psychosis or other mental health problems in childhood might reduce affected individuals' social networks, and the resulting social isolation may increase the likelihood of psychotic phenomena developing or persisting. Indeed, individuals with early signs of psychosis have been shown to have limited social networks (Gayer-Anderson and Morgan, 2013), and psychotic symptoms and other mental health problems in childhood have also been shown to increase risk for adolescent psychotic phenomena (Polanczyk *et al.*, 2010; Zammit *et al.*, 2013). Those without mental health problems in childhood might therefore be more likely to have greater social support and less likely to develop psychotic phenomena in adolescence. Thus, it is important to take into account prior psychopathology to improve understanding of the temporal association between social support and adolescent psychotic phenomena.

This study aims to utilise the discordant twin design (Vitaro *et al.*, 2009; Pingault *et al.*, 2018) in a longitudinal cohort to consider the relative family-wide *v.* unique environmental effects of social support on adolescent psychotic experiences and control for earlier psychopathology. This approach capitalises on the fact that twins reared together share the same family environment and the same genes (100% for monozygotic [MZ] twins; 50% for dizygotic twins [DZ]). To estimate family-wide effects of social support we will firstly consider between-twin effects, thus testing whether twin pairs with higher social support also have a reduced likelihood of psychotic phenomena relative to other twin pairs with lower levels of social support. Next to estimate the unique environmental effects of social support we will consider the within-twin effects, i.e. whether twins with higher levels of social support than their co-twin also have a reduced likelihood of adolescent psychotic phenomena relative to their co-twin. Modelling these effects together allows us to ascertain the unique environmental effects of social support on psychotic phenomena relative to shared family-wide effects. We will additionally conduct analyses restricted to MZ twins to fully rule out genetic confounding, and also control for psychotic symptoms and other mental health problems in childhood to limit reverse causality. Analyses will be run in the full general population sample and also in the subgroup of adolescents who have been poly-victimised to test whether a unique environmental pathway is still evident in this high-risk group.

Methods

Study cohort

Participants were members of the Environmental Risk (E-Risk) Longitudinal Twin Study, which tracks the development of a nationally-representative birth cohort of 2232 British twin children. The sample was drawn from a larger cohort of twins born in England and Wales in 1994–1995 (Trouton *et al.*, 2002). Full details about the sample are reported elsewhere (Moffitt and E-Risk Study Team, 2002). Briefly, the E-Risk sample was constructed in 1999–2000, when 1116 families with same-sex 5-year-old twins (93% of those eligible) participated in home-visit assessments. Families were recruited to represent the UK population of families with newborns in the 1990s, based on residential location throughout England and Wales and mothers' age. Teenaged mothers with twins were over-selected to replace high-risk families who were selectively lost to the register through

non-response. Older mothers having twins via assisted reproduction were under-selected to avoid an excess of well-educated older mothers. E-Risk families are representative of UK households across the spectrum of neighbourhood-level deprivation: 25.6% of E-Risk families live in 'wealthy achiever' neighbourhoods compared to 25.3% of households nation-wide; 5.3% *v.* 11.6% live in 'urban prosperity' neighbourhoods; 29.6% *v.* 26.9% live in 'comfortably off' neighbourhoods; 13.4% *v.* 13.9% live in 'moderate means' neighbourhoods; and 26.1% *v.* 20.7% live in 'hard-pressed' neighbourhoods (Caspi *et al.*, 2000; CACI, 2006). E-Risk families under-represent 'urban prosperity' neighbourhoods because such households are likely to be childless. The sample comprised 56% MZ and 44% DZ twin pairs, and sex was evenly distributed within zygosity (49% male). All families were English speaking, and the majority (93.7%) were White.

Follow-up home visits were conducted when children were 7 years (98% of the 1116 E-Risk Study families participated), 10 years (96% participation), 12 years (96% participation) and 18 years (93% participation). Home visits at ages 5, 7, 10, and 12 years included assessments with participants as well as their mother (or primary caretaker); the home visit at age 18 included interviews only with the participants. Each twin participant was assessed by a different interviewer. The average age of the 2066 twins at the time of the age 18 assessment was 18.4 years (s.d. = 0.36); all interviews were conducted after the 18th birthday. There were no differences between those who did and did not take part at age 18 in terms of socioeconomic status (SES) assessed when the cohort was initially defined ($\chi^2 = 0.86$, $p = 0.65$), age-5 IQ scores ($t = 0.98$, $p = 0.33$), or age-5 internalising or externalising behaviour problems ($t = 0.40$, $p = 0.69$ and $t = 0.41$, $p = 0.68$, respectively). The Joint South London and Maudsley and the Institute of Psychiatry Research Ethics Committee approved each phase of the study. Parents gave informed consent, and participants gave assent at ages 5–12 and informed consent at age 18.

Measures

Adolescent psychotic experiences

At age 18, each E-Risk participant was privately interviewed by a trained research worker about 13 psychotic experiences occurring since age 12. Seven items pertained to delusions and hallucinations, such as 'have you ever thought you were being watched, followed or spied on?' and 'do you hear voices that others cannot?'. This interview has been described in detail previously (Polanczyk *et al.*, 2010). Interviewers probed responses to these items to ensure that participants understood the questions and obtained descriptions of their experiences. The other six items pertained to unusual experiences which drew on item pools since formalised in prodromal psychosis instruments including the PRIME-screen and SIPS (Loewy *et al.*, 2011). These included 'I worry that my food may be poisoned' and 'My thinking is unusual or frightening'. Further information on this measure is provided in the online Supplementary Materials. Interviewers coded each item 0, 1, 2 indicating respectively 'not present', 'probably present', and 'definitely present'. All 13 items were summed to create a psychotic experiences scale (range = 0–18, $M = 1.19$, s.d. = 2.58). Just over 30% of participants had at least one psychotic experience between ages 12 and 18 ($n = 623$, 30.2%). This is similar to the prevalence of self-reported psychotic experiences in other community samples of teenagers and young adults (Yoshizumi *et al.*, 2004; Kelleher *et al.*, 2012c). Almost all of the twins who

Table 1. Demographic characteristics of the Environmental Risk study cohort with psychotic experiences data available at age 18

Characteristic	Whole sample ($N = 2063$) n (%)
Gender	
Male	980 (47.5)
Female	1083 (52.5)
Zygosity	
Monozygotic	1164 (56.4)
Dizygotic	899 (43.6)
Family socioeconomic status	
Low	691 (33.5)
Intermediate	683 (33.1)
High	689 (33.4)
	Mean (standard deviation)
Age at interview	18.4 years (0.36)

participated in the age-18 assessment reported on their psychotic experiences ($N = 2063$, 99.9%). An overview of the demographics of the sample with psychotic experiences data available is provided in Table 1.

Social support

Social support was assessed using the Multidimensional Scale of Perceived Social Support (MSPSS), which assesses individuals' access to supportive relationships with family, friends and significant others (Zimet *et al.*, 1988). The 12 items in the MSPSS consist of statements such as 'There is a special person who is around when I am in need' and 'I can count on my friends when things go wrong'. At age 18, participants rated these statements as 'not true' (0), 'somewhat true' (1), or 'very true' (2). We summed the scores to produce an overall social support scale with higher scores reflecting greater social support (internal consistency: $\alpha = 0.88$). In addition, each of the three sub-scales (social support from either family, friends or significant others) were utilised separately to examine whether the findings were consistent across these different sources of social support.

Adolescent poly-victimisation

At age 18, participants were interviewed about exposure to a range of adverse experiences between 12–18 years using the Juvenile Victimization Questionnaire, 2nd revision (JVQ-R2) (Finkelhor *et al.*, 2011) adapted as a clinical interview, which has been outlined in a previous paper (Fisher *et al.*, 2015) and in the online Supplementary Materials. Each twin was interviewed by a different research worker, and each JVQ question was asked for the period 'since you were 12'. Age 12 is a salient age for our participants because it is the age when British children leave primary school to enter secondary school. Our adapted JVQ comprised 45 questions covering 7 different forms of victimisation: maltreatment, neglect, sexual victimisation, family violence, peer/sibling victimisation, internet/mobile phone victimisation, and crime victimisation. The worst experience (according to the participant) for each victimisation type was rated by trained coders using a 6-point scale: 0 = not exposed, then 1–5 for increasing levels of severity. The adolescent poly-victimisation variable was derived

by summing all victimisation experiences that received a code of '4' or '5' (i.e. severe exposure). Due to small numbers in some of the groups, we collapsed this variable into '0' not victimised (64.6%), '1' experienced 1 type of severe victimisation (19.2%), and '2' poly-victimised (16.2%; experienced 2 or more types of severe victimisation).

Potential confounders

Family SES was measured via a composite of parental income (total household), education (highest for mother/father), and occupation (highest for mother/father) when children were aged 5 (Trzesniewski *et al.*, 2006), and was categorised into tertiles (i.e. low-, medium-, and high-SES). Childhood psychotic symptoms pertaining to seven delusions and hallucinations were measured when children were aged 12 during private interviews. Items and interviewer notes were assessed by a psychiatrist expert in schizophrenia, a psychologist expert in interviewing children, and a child and adolescent psychiatrist to verify the validity of the symptoms (Polanczyk *et al.*, 2010). A total of 5.9% of children reported experiencing at least one definite psychotic symptom at age 12 ($N = 125$). In order to capture earlier psychopathology more broadly, a variable was also created for the presence *v.* absence of any childhood mental health problems that captured children who met criteria for extreme anxiety, clinically-relevant depression symptoms, attention deficit hyperactivity disorder (ADHD), or conduct disorder by age 12 (see online Supplementary Materials).

Statistical methods

Analyses were conducted in STATA 15 (Stata-Corp, College Station, TX). We applied Generalised Estimating Equations (GEE) with binomial function specified (logistic regression) and an exchangeable correlation structure to account for familial clustering in order to simultaneously estimate the family-wide (between-twin pair) and unique (within-twin pair) effects of social support on age-18 psychotic experiences (Carlin *et al.*, 2005; Vitaro *et al.*, 2009). The between-twin pair analysis considers whether pairs of twins with higher social support than other twin pairs are also less likely to have psychotic experiences. In contrast, the within twin-pair analysis considers whether the twin with higher social support than his or her co-twin is also less likely to have psychotic experiences than his or her co-twin (Carlin *et al.*, 2005). Because co-twins share their rearing environment as well as half (dizygotic twins) or all (monozygotic twins) their genes, significant *within-twin pair effects* would indicate that social support is associated with a reduced likelihood of psychotic experiences independent of latent, family-wide factors, thus suggesting a unique environmental effect of social support. Further restricting analyses to MZ twins fully rules out genetic influences, and additionally controlling for age-12 psychotic symptoms and other childhood mental health problems accounts for the possibility of reverse causality. All analyses were conducted in the full general population sample and also in the sub-group of poly-victimised adolescents where both twins were poly-victimised ($n = 158$).

Results

We have previously shown that social support was associated with a reduced likelihood of age-18 psychotic experiences amongst adolescents in the whole sample (OR = 0.91, 95% CI 0.89–0.93,

$p < 0.001$) and amongst a high-risk group exposed to poly-victimisation (OR = 0.93; 95% CI 0.88–0.98, $p = 0.009$) after controlling for gender, family SES, age-12 psychotic symptoms and other mental health problems at age 12 (Crush *et al.*, 2018a). There was a reasonable level of discordance among both MZ (30.2%) and DZ (32.7%) twin pairs for age-18 psychotic experiences, and the intraclass correlation for the total social support score at age 18 was greater than 0 but less than 1 for both MZ (ICC = 0.46, 95% CI 0.40–0.52) and DZ (ICC = 0.15, 95% CI 0.06–0.24) twins indicating that there was both within- and between-twin pair variation in these measures.

Is the association between increased social support and a reduced likelihood of psychotic experiences during adolescence accounted for by shared environmental and genetic factors?

Using discordant twin analyses in the whole sample, we considered the association between overall social support, and each subtype of social support, with a reduced likelihood of adolescent psychotic experiences at age 18. We found that these associations were explained by family-wide effects of social support shared by twin pairs (Table 2), therefore showing that twin pairs with higher social support were less likely to report psychotic experiences relative to twin pairs who reported lower social support. Notably, we also found evidence of a unique environmental effect, whereby higher perceived social support by one twin relative to their co-twin was associated with a reduced likelihood of psychotic experiences (Table 2). When analyses were repeated for MZ twins only (to fully control for genetic influences), we again found both family-wide influences and the unique environment to be implicated in the associations for total social support and for support from both family and friends (Table 2), although the findings were inconclusive in relation to social support from significant others. The fact that the results showed a significant within-twin-pair association for total social support, and also social support from friends and family, with a reduced likelihood of psychotic experiences in the whole sample and MZ group provides support for the presence of a unique environmentally-mediated protective pathway for these social support types.

Similarly, amongst twins exposed to poly-victimisation, we found evidence of a unique environmental pathway between social support from friends and a reduced likelihood of adolescent psychotic experiences (Table 2). When analyses were restricted to MZ twins only who had both been exposed to poly-victimisation, we found similar trends (Table 2). However, these effects were not significant, possibly due to the relatively small number of twins ($N = 96$) within this sub-group.

Is the association between increased social support and a reduced likelihood of psychotic experiences accounted for by childhood psychopathology?

Next, in order to investigate any potential reverse causality between social support and psychotic experiences, we controlled for psychotic symptoms and other mental health problems at age 12 to support the interpretation of the directionality of the association. We found that the unique environmental effect of total social support on the reduced likelihood of adolescent psychotic experiences remained significant when accounting for earlier psychopathology within the full sample and when analyses were restricted to MZ twins (Table 3), thus suggesting this type of

Table 2. Family-wide and unique environmental effects of social support on age-18 psychotic experiences

Social support scale	Whole sample				Both twins poly-victimised			
	All twins N = 2058		MZ twins N = 1162		All twins N = 158		MZ twins N = 96	
	Adjusted OR ^a (95% CI)	p Value	Adjusted OR ^a (95% CI)	p Value	Adjusted OR ^a (95% CI)	p Value	Adjusted OR ^a (95% CI)	p Value
Total								
Family-wide	0.89 (0.86–0.92)	<0.001	0.91 (0.88–0.95)	<0.001	0.94 (0.84–1.04)	0.237	0.98 (0.85–1.11)	0.710
Unique	0.93 (0.90–0.96)	<0.001	0.93 (0.89–0.97)	<0.001	0.91 (0.82–1.00)	0.052	0.92 (0.81–1.04)	0.193
Family								
Family-wide	0.75 (0.70–0.80)	<0.001	0.77 (0.70–0.84)	<0.001	0.85 (0.70–1.02)	0.077	0.84 (0.66–1.06)	0.143
Unique	0.85 (0.78–0.92)	<0.001	0.86 (0.77–0.96)	0.007	0.81 (0.61–1.08)	0.140	0.80 (0.61–1.05)	0.114
Friends								
Family-wide	0.78 (0.73–0.84)	<0.001	0.82 (0.76–0.90)	<0.001	0.90 (0.73–1.10)	0.303	1.03 (0.80–1.33)	0.831
Unique	0.88 (0.82–0.94)	<0.001	0.88 (0.80–0.96)	0.004	0.81 (0.69–0.96)	0.012	0.85 (0.69–1.04)	0.121
Significant Others								
Family-wide	0.90 (0.84–0.97)	0.004	0.97 (0.89–1.07)	0.535	1.01 (0.79–1.30)	0.914	1.15 (0.87–1.51)	0.321
Unique	0.92 (0.86–0.99)	0.023	0.92 (0.84–1.02)	0.120	0.92 (0.68–1.24)	0.592	1.00 (0.62–1.63)	0.984

CI, confidence interval; MZ, monozygotic; OR, odds ratio.

Bold text indicates $p < 0.05$. Family-wide indicates between-twin pair difference; unique, within-twin pair difference.

^aAdjusted for child's gender.

Table 3. Family-wide and unique environmental effects of social support on age-18 psychotic experiences controlling for childhood mental health problems

Social support scale	Whole sample				Both twins poly-victimised			
	All twins N = 2058		MZ twins N = 1162		All twins N = 158		MZ twins N = 96	
	Adjusted OR ^a (95% CI)	p Value	Adjusted OR ^a (95% CI)	p Value	Adjusted OR ^a (95% CI)	p Value	Adjusted OR ^a (95% CI)	p Value
Total								
Family-wide	0.90 (0.87–0.93)	<0.001	0.92 (0.88–0.96)	<0.001	0.93 (0.83–1.04)	0.191	0.97 (0.83–1.13)	0.674
Unique	0.93 (0.90–0.96)	<0.001	0.92 (0.88–0.97)	0.001	0.90 (0.82–1.00)	0.045	0.91 (0.80–1.04)	0.163
Family								
Family-wide	0.77 (0.72–0.83)	<0.001	0.79 (0.72–0.87)	<0.001	0.83 (0.68–1.01)	0.058	0.80 (0.63–1.03)	0.083
Unique	0.86 (0.79–0.93)	<0.001	0.86 (0.76–0.97)	0.012	0.85 (0.64–1.13)	0.256	0.81 (0.60–1.08)	0.144
Friends								
Family-wide	0.80 (0.75–0.85)	<0.001	0.84 (0.77–0.91)	<0.001	0.91 (0.74–1.11)	0.339	1.08 (0.84–1.39)	0.531
Unique	0.87 (0.81–0.93)	<0.001	0.86 (0.78–0.96)	0.004	0.78 (0.66–0.92)	0.003	0.82 (0.65–1.03)	0.092
Significant Others								
Family-wide	0.91 (0.85–0.98)	0.011	0.97 (0.88–1.07)	0.596	0.99 (0.74–1.32)	0.928	1.13 (0.79–1.64)	0.502
Unique	0.92 (0.85–0.99)	0.020	0.91 (0.82–1.01)	0.092	0.89 (0.65–1.20)	0.440	1.00 (0.59–1.67)	0.989

CI, confidence interval; MZ, monozygotic; OR, odds ratio.

Bold text indicates $p < 0.05$. Family-wide indicates between-twin pair difference; unique, within-twin pair difference.

^aAdjusted for child's gender, age-12 psychotic symptoms, and other mental health problems at age 12.

reverse causality was unlikely to be operating. The buffering effects of support from friends and family were also robust to adjustment for childhood psychopathology and shared family factors (Table 3).

A similar pattern of results was found after controlling for earlier mental health problems in the poly-victimised group, with total social support and support from friends continuing to have a significant unique environmental effect on the reduced

likelihood of adolescent psychotic experiences (Table 3). When restricting analyses to MZ twins, only non-significant trends were found (Table 3) but again this was probably due to the small number in this high-risk sub-group ($N = 96$).

Discussion

This is the first study to consider whether social support has an environmentally-mediated effect on psychotic phenomena amongst adolescents in the general population and amongst those at high risk by virtue of having been exposed to poly-victimisation. Our findings indicate a unique environmental buffering effect of higher perceived social support on psychotic phenomena amongst our general population sample. Whilst it is important to note that effects sizes were relatively modest in size, these effects were apparent in relation to overall social support and also separately for social support from friends and family. These results held after accounting for shared family-wide environmental and genetic influences as well as for earlier psychopathology thus limiting these potentially confounding effects and reverse causality explanations. We also found evidence for a direct environmentally-mediated protective effect of perceived social support on adolescent psychotic experiences even within the high-risk poly-victimised group. Collectively, these findings provide initial evidence for a possible causal association between higher perceived social support and the reduced likelihood of adolescent psychotic experiences and therefore add weight to the importance of social support as a potential candidate for preventive interventions focused upon adolescent psychotic phenomena.

No prior research has addressed this particular question and thus it is not possible to draw comparisons directly with previous research. Nonetheless, one study conducted in adults diagnosed with psychotic disorders reported a similar buffering effect of social support following exposure to specific types of childhood victimisation (Gayer-Anderson *et al.*, 2015). Moreover, in this cohort we have previously reported that warm parent-child relationships had an environmentally mediated protective effect on children's behavioural adjustment following bullying victimisation (Bowes *et al.*, 2010). Although these findings relate to a different type of psychopathology, they suggest that supportive relationships may promote resilience to a range of mental health outcomes following victimisation exposure in childhood. Given the paucity of research in this area, more studies are needed and our findings require replication in other large population-based cohorts.

Overall our findings have practical implications as they suggest that early prevention efforts focusing upon improving perceived social support – through greater availability of supportive figures or enhancing perceptions of existing social support, could be effective in protecting against the development of psychotic phenomena in adolescence. Interventions aimed at improving social support from family and peers have previously been found to be effective amongst individuals who already have psychosis (Pilling *et al.*, 2002; Norman *et al.*, 2005; Castelein *et al.*, 2008). Whilst family interventions have been most widely applied, recently there has been increased support for the use of peer interventions (Harrop *et al.*, 2015; Morin *et al.*, 2017) which our findings also indicate might be helpful. Given resources for interventions are limited, it is possible that internet-based peer support networks could represent a promising solution (Álvarez-Jiménez *et al.*, 2012; Naslund *et al.*, 2016). Our findings support targeting perceptions of social support amongst

adolescents exposed to poly-victimisation. This is practically relevant for clinicians given it has been suggested that individuals exposed to complex trauma may need specific types of treatment or interventions compared to those not exposed (Cook *et al.*, 2005).

There are several mechanisms through which social support may exert protective influences in relation to psychotic experiences in both the general population and amongst poly-victimised individuals. For instance, it is possible that social support may facilitate stress reduction (Cohen and Wills, 1985), improve self-esteem (Dumont and Provost, 1999; Turner *et al.*, 2015), and may also reduce feelings of loneliness (Sündermann *et al.*, 2014), which have all been implicated in the development of psychotic phenomena (Corcoran *et al.*, 2003; Smith *et al.*, 2006; Pruessner *et al.*, 2011; Lim *et al.*, 2018).

Limitations

Some limitations warrant consideration. Firstly, both social support and psychotic experiences were measured at age 18 which has implications for interpreting the directionality of the association between them. We did, however, control for a broad range of earlier mental health problems at age 12, including age-12 psychotic symptoms, to account for this as far as possible and are thus able to largely rule out reverse causality. Research around the consistency of social support during adolescence has not suggested any fundamental shifts in total social support levels over time, whilst there are trends for family support being replaced by peer support, relatively stable levels of support appear to be maintained (Levitt *et al.*, 1993; Cantin and Boivin, 2004). Relatedly, as the social support scale used is a subjective measure reflecting individuals' perceptions of support from friends, family and significant others, it is possible that adolescents who develop psychotic experiences perceive their social support levels to be lower than they actually are. Therefore, we welcome replication of our findings amongst cohorts with co-informant measures of social support. Additionally, due to the number of poly-victimised adolescents being relatively low ($N = 334$), our ability to detect some associations may have been affected by this. Moreover, we defined poly-victimisation as reported exposure to two or more severe types of victimisation in adolescence which, although it is a commonly used approach, means individuals in this group may have experienced quite different combinations of victimisation. Other approaches, such as latent class analysis, have been used to create more homogeneous groupings of victimisation types (e.g. Ford *et al.*, 2010), and it is possible that utilisation of such methods might have led to different findings. Our analyses thus warrant replication in even larger twin cohorts and with different methods of conceptualising poly-victimisation. Finally, our measure of psychotic experiences was a self-report measure and it is possible that it captured some genuine experiences (e.g. being followed by someone), as well as psychotic phenomena (e.g. being followed by a fictional character). Nonetheless, we have previously shown that higher levels of social support were associated with a reduced likelihood of clinically-verified adolescent psychotic symptoms (Crush *et al.*, 2018a), indicating that this is unlikely to have substantially affected our findings.

Conclusion

The association between greater perceived social support and a reduced likelihood of psychotic experiences in adolescence

appears to be extremely robust, even in the context of poly-victimisation, as it was not fully accounted for by family-wide environmental or genetic factors nor was there any evidence of reverse causation. These findings suggest that early intervention programmes focused on increasing perceptions of social support, particularly from friends, have the potential to prevent the emergence of psychotic experiences amongst adolescents. Given that there are finite resources for interventions, efforts might be most efficiently targeted at adolescents exposed to multiple forms of victimisation who are at high risk of developing psychotic phenomena.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291719001983>.

Acknowledgements. We are grateful to the study mothers and fathers, the twins, and the twins' teachers for their participation. Our thanks to Terrie E. Moffitt and Avshalom Caspi the founders of the E-Risk Study, CACI, Inc., to Laurie Hannigan for statistical advice, and to members of the E-Risk team for their dedication, hard work and insights.

Financial support. The E-Risk Study is funded by the UK Medical Research Council (G1002190). Additional support was provided by the National Institute of Child Health and Human Development (HD077482); the Jacobs Foundation; the Avielle Foundation; a research grant from the National Society for Prevention of Cruelty to Children (NSPCC) and Economic and Social Research Council (ESRC); a Medical Research Council Studentship to EC; and an MQ Fellows Award (MQ14F40) to HLF. LA is the Mental Health Leadership Fellow for the UK ESRC.

Conflict of interest. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Álvarez-Jiménez M, Gleeson JF, Bendall S, Lederman R, Wadley G, Killackey E and McGorry PD (2012) Internet-based interventions for psychosis. A sneak-peek into the future. *Psychiatric Clinics of North America* **35**, 735–747.
- Arseneault L, Cannon M, Fisher HL, Polanczyk G, Moffitt TE and Caspi A (2011) Childhood trauma and children's emerging psychotic symptoms: a genetically sensitive longitudinal cohort study. *American Journal of Psychiatry* **168**, 65–72.
- Blain MD, Thompson JM and Whiffen VE (1993) Attachment and perceived social support in late adolescence. *Journal of Adolescent Research* **8**, 226–241.
- Bowes L, Maughan B, Caspi A, Moffitt TE and Arseneault L (2010) Families promote emotional and behavioural resilience to bullying: evidence of an environmental effect. *Journal of Child Psychology & Psychiatry* **51**, 809–817.
- CACI (2006) *ACORN User Guide*. London, UK: CACI Information Services.
- Cantin S and Boivin M (2004) Change and stability in children's social network and self-perceptions during transition from elementary to junior high school. *International Journal of Behavioral Development* **28**, 561–570.
- Carlin JB, Gurrin LC, Sterne JAC, Morley R and Dwyer T (2005) Regression models for twin studies: a critical review. *International Journal of Epidemiology* **34**, 1089–1099.
- Caspi A, Taylor A, Moffitt TE and Plomin R (2000) Neighborhood deprivation affects children's mental health: environmental risks identified in a genetic design. *Psychological Science* **11**, 338–342.
- Castelein S, Bruggeman R, Van Busschbach JT, Van Der Gaag M, Stant AD, Knegtering H and Wiersma D (2008) The effectiveness of peer support groups in psychosis: a randomized controlled trial. *Acta Psychiatrica Scandinavica* **118**, 64–72.
- Claes M (1994) Friendship characteristics of adolescents referred for psychiatric treatment. *Journal of Adolescent Research* **9**, 180–192.
- Cohen S and Wills TA (1985) Stress, social support, and the buffering hypothesis. *Psychological Bulletin* **98**, 310–357.
- Cook A, Spinazzola J, Ford J, Lanktree C, Blaustein M, Cloitre M, DeRosa R, Hubbard R, Kagan R, Liautaud J, Mallah K, Olafson E and van der Kolk B (2005) Complex trauma in children and adolescents. *Psychiatric Annals* **35**, 390–398.
- Corcoran C, Walker E, Huot R, Mittal V, Tessner K, Kestler L and Malaspina D (2003) The stress cascade and schizophrenia: etiology and onset. *Schizophrenia Bulletin* **29**, 671–692.
- Crush E, Arseneault A, Moffitt TE, Danese A, Caspi A, Jaffee SR, Matthews T and Fisher HL (2018a) Protective factors for psychotic experiences amongst adolescents exposed to multiple forms of victimization. *Journal of Psychiatric Research* **104**, 32–28.
- Crush E, Arseneault A, Jaffee SR, Danese A and Fisher HL (2018b) Protective factors for psychotic symptoms among poly-victimized children. *Schizophrenia Bulletin* **44**, 691–700.
- Dumont M and Provost MA (1999) Resilience in adolescents: protective role of social support, coping strategies, self-esteem, and social activities on experience of stress and depression. *Journal of Youth and Adolescence* **28**, 343–363.
- Finkelhor D, Hamby SL, Turner H and Ormrod R (2011) *The Juvenile Victimization Questionnaire: 2nd Revision (JVQ-R2)*. Durham, NH: Crimes Against Children Research Center.
- Fisher HL, Caspi A, Poulton R, Meier MH, Harrington H, Arseneault L and Moffitt TE (2013) Specificity of childhood psychotic symptoms for predicting schizophrenia by 38 years of age: a birth cohort study. *Psychological Medicine* **43**, 2077–2086.
- Fisher HL, Caspi A, Moffitt TE, Wertz J, Gray R, Newbury J, Ambler A, Zavos H, Danese A, Mill J, Odgers CL, Pariante C, Wong CC and Arseneault L (2015) Measuring adolescents' exposure to victimization: the Environmental Risk (E-Risk) Longitudinal Twin Study. *Development and Psychopathology* **27**, 1399–1416.
- Ford JD, Elhai JD, Connor DF and Frueh BC (2010) Poly-victimization and risk of posttraumatic, depressive, and substance use disorders and involvement in delinquency in a national sample of adolescents. *Journal of Adolescent Health* **46**, 545–552.
- Gayer-Anderson C and Morgan C (2013) Social networks, support and early psychosis: a systematic review. *Epidemiology and Psychiatric Sciences* **22**, 131–146.
- Gayer-Anderson C, Fisher HL, Fearon P, Hutchinson G, Morgan K, Dazzan P, Boydell J, Doody GA, Jones PB, Murray RM, Craig TK and Morgan C (2015) Gender differences in the association between childhood physical and sexual abuse, social support and psychosis. *Social Psychiatry and Psychiatric Epidemiology* **50**, 1489–1500.
- Harrop C, Ellett L, Brand R and Lobban F (2015) Friends interventions in psychosis: a narrative review and call to action. *Early Intervention in Psychiatry* **9**, 269–278.
- Janssen I, Krabbendam L, Bak M, Hanssen M, Vollebergh W, de Graaf R and van Os J (2004) Childhood abuse as a risk factor for psychotic experiences. *Acta Psychiatrica Scandinavica* **109**, 38–45.
- Kelleher I, Connor D, Clarke MC, Devlin N, Harley M and Cannon M (2012a) Prevalence of psychotic symptoms in childhood and adolescence: a systematic review and meta-analysis of population-based studies. *Psychological Medicine* **42**, 1857–1863.
- Kelleher I, Lynch F, Harley M, Molloy C, Roddy S, Fitzpatrick C and Cannon M (2012b) Psychotic symptoms in adolescence index risk for suicidal behavior: findings from 2 population-based case-control clinical interview studies. *Archives of General Psychiatry* **69**, 1277–1283.
- Kelleher I, Keeley H, Corcoran P, Lynch F, Fitzpatrick C, Devlin N, Molloy C, Roddy S, Clarke MC, Harley M, Arseneault L, Wasserman C, Carli V, Sarchiapone M, Hoven C, Wasserman D and Cannon M (2012c) Clinicopathological significance of psychotic experiences in non-psychotic young people: evidence from four population-based studies. *British Journal of Psychiatry* **201**, 26–32.
- Kelleher I, Wigman JTW, Harley M, O'Hanlon E, Coughlan H, Rawdon C, Murphy J, Power E, Higgins NM and Cannon M (2015) Psychotic experiences in the population: association with functioning and mental distress. *Schizophrenia Research* **165**, 9–14.

- Kendler KS** (1997) Social support: a genetic-epidemiologic analysis. *American Journal of Psychiatry* **154**, 1398–1404.
- Levitt MJ, Guacci-Franco N and Levitt JL** (1993) Convoys of social support in childhood and early adolescence: structure and function. *Developmental Psychology* **29**, 811–818.
- Lim MH, Gleeson M, Alvarez-Jimenez M and Penn DL** (2018) Loneliness in psychosis: a systematic review. *Social Psychiatry and Psychiatric Epidemiology* **53**, 221–238.
- Loewy RL, Pearson R, Vinogradov S, Bearden CE and Cannon TD** (2011) Psychosis risk screening with the Prodromal Questionnaire – brief version (PQ-B). *Schizophrenia Research* **129**, 42–46.
- Matthews T, Danese A, Wertz J, Odgers CL, Ambler A, Moffitt TE and Arseneault L** (2016) Social isolation, loneliness and depression in young adulthood: a behavioural genetic analysis. *Social Psychiatry and Psychiatric Epidemiology* **51**, 339–348.
- McGrath JJ, Sukanta S, Al-Hamzawi A, Alonso J, Bromet EJ, Bruffaerts R, Caldas-De-Almeida JM, Chiu WT, De Jonge P, Fayyad J, Florescu S, Gureje O, Haro JM, Hu C, Kovess-Masfety V, Lepine JP, Lim CCW, Mora MEM, Navarro-Mateu F, Ochoa S, Sampson N, Scott K, Viana MC and Kessler RC** (2015) Psychotic experiences in the general population: a cross-national analysis based on 31 261 respondents from 18 countries. *JAMA Psychiatry* **72**, 697–705.
- Moffitt TE and E-Risk Study Team** (2002) Teen-aged mothers in contemporary Britain. *Journal of Child Psychology and Psychiatry* **43**, 727–742.
- Morin F, Dhir A, Mitchell E and Jones A** (2017) Social support: a useful tool in the management of psychotic disorders. *University of British Columbia Medical Journal* **8**, 10–12.
- Naslund JA, Aschbrenner KA, Marsch LA and Bartels SJ** (2016) The future of mental health care: Peer-To-peer support and social media. *Epidemiology and Psychiatric Sciences* **25**, 113–122.
- Newbury J, Arseneault L, Caspi A, Moffitt TE, Odgers C and Fisher HL** (2016) Why are children in urban neighborhoods at increased risk for psychotic symptoms? Findings from a UK longitudinal cohort study. *Schizophrenia Bulletin* **42**, 1372–1383.
- Norman RMG, Malla AK, Manchanda R, Harricharan R, Takhar J and Northcott S** (2005) Social support and three-year symptom and admission outcomes for first episode psychosis. *Schizophrenia Research* **80**, 227–234.
- Pilling S, Bebbington P, Kuipers E, Garety P, Geddes J, Orbach G and Morgan C** (2002) Psychological treatments in schizophrenia: I. Meta-analysis of family intervention and cognitive behaviour therapy. *Psychological Medicine* **32**, 763–782.
- Pingault JB, O'Reilly PF, Schoeler T, Plouhidis GB, Rijdsdijk F and Dudbridge F** (2018) Using genetic data to strengthen causal inference in observational research. *Nature Reviews Genetics* **19**, 566–580.
- Polanczyk G, Moffitt T, Arseneault L, Cannon M, Ambler A, Keefe RS, Houts R, Odgers CL and Caspi A** (2010) Etiological and clinical features of childhood psychotic symptoms. *Archives of General Psychiatry* **67**, 328–338.
- Pruessner M, Iyer S, Faridi K, Joober R and Malla AK** (2011) Stress and protective factors in individuals at ultra-high risk for psychosis, first episode psychosis and healthy controls. *Schizophrenia Research* **129**, 29–35.
- Riggio HR and Kwong WY** (2011) Paranoid thinking, quality of relationships with parents, and social outcomes among young adults. *Journal of Family Issues* **32**, 1030–1049.
- Shevlin M, Houston JE, Dorahy MJ and Adamson G** (2008) Cumulative traumas and psychosis: an analysis of the national comorbidity survey and the British Psychiatric Morbidity Survey. *Schizophrenia Bulletin* **34**, 193–199.
- Smith B, Fowler DG, Freeman D, Bebbington P, Bashforth H, Garety P, Dunn G and Kuipers E** (2006) Emotion and psychosis: links between depression, self-esteem, negative schematic beliefs and delusions and hallucinations. *Schizophrenia Research* **86**, 181–188.
- Sündermann O, Onwumere P, Bebbington P and Kuipers E** (2014) Social networks and support in first-episode psychosis: exploring the role of loneliness and anxiety. *Social Psychiatry and Psychiatric Epidemiology* **49**, 359–366.
- Trouton A, Spinath FM and Plomin R** (2002) Twins Early Development Study (TEDS): a multivariate, longitudinal genetic investigation of language, cognition and behavior problems in childhood. *Twin Research* **5**, 444–448.
- Trzesniewski TH, Moffitt TE, Caspi A, Taylor A and Maughan B** (2006) Revisiting the association between reading achievement and antisocial behavior: new evidence of an environmental explanation from a twin study. *Child Development* **77**, 72–88.
- Turner HA, Shattuck A, Finkelhor D and Hamby S** (2015) Effects of poly-victimization on adolescent social support, self-concept, and psychological distress. *Journal of Interpersonal Violence* **32**, 1–26.
- Vitaro F, Brendgen M and Arseneault L** (2009) The discordant MZ-twin method: one step closer to the holy grail of causality. *International Journal of Behavioral Development* **33**, 376–382.
- Yoshizumi T, Murase S, Honjo S, Kaneko H and Murakami T** (2004) Hallucinatory experiences in a community sample of Japanese children. *Journal of the American Academy of Child and Adolescent Psychiatry* **43**, 1030–1036.
- Zammit S, Kounali D, Cannon M, David A, Gunnell D, Heron J, Jones PB, Lewis S, Sullivan S, Wolke D and Lewis G** (2013) Psychotic experiences and psychotic disorders at age 18 in relation to psychotic experiences at age 12 in a longitudinal population-based cohort study. *American Journal of Psychiatry* **170**, 742–750.
- Zavos HMS, Freeman D, Haworth CMA, McGuire P, Plomin R, Cardno AG and Ronald A** (2014) Consistent etiology of severe, frequent psychotic experiences and milder, less frequent manifestations. *JAMA Psychiatry* **71**, 1049–1057.
- Zimet G, Powell S, Farley G, Werkman S and Berkoff K** (1988) Psychometric characteristics of the multidimensional scale of perceived social support. *Journal of Personality Assessment* **55**, 610–617.