



## King's Research Portal

*Document Version*  
Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Bourni, T., Langford, M., & Tinaglia, G. (Accepted/In press). A collapsing ancient solution of mean curvature flow in  $\mathbb{R}^3$ . *JOURNAL OF DIFFERENTIAL GEOMETRY*.

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

Report on the paper

## **Collapsing ancient solutions of mean curvature flow**

by T. BOURNI, M. LANGFORD AND G. TINAGLIA

This paper deals with ancient solutions of the mean curvature flow, i.e. solutions which are defined for arbitrary negative times. Ancient solutions have attracted much attention in recent years because they are relevant in the analysis of singularities of the flow. The simplest example is given by the shrinking sphere. Few examples are also known of nonspherical convex compact ancient solutions, which become more and more eccentric as they evolve backwards in time. The oldest one is a solution of curve shortening flow in the plane, known as the Angenent oval. Higher dimensional examples have been constructed in [12,19]. While the Angenent oval is contained in a strip for all times, the higher dimensional examples of [12,19] sweep the whole space and are noncollapsed in the sense of [1]. For curves in the plane, it is known that the shrinking sphere and the Angenent oval are the only compact ancient solutions. In higher dimensions, a classification of ancient solution seems much harder to obtain, although the results of [14] show that this class enjoys strong rigidity results.

This paper focuses the attention on ancient solutions which can be considered a higher dimensional analogue of the Angenent oval. The authors first prove the existence of a rotationally symmetric ancient solution which resembles a rotated Angenent oval: in particular, the solution is contained in a slab and the asymptotic profile is given by the grim reaper curve, in any direction orthogonal to the axis of rotation. In contrast with the examples of [12,19], this solution is collapsed for large negative times. The existence result is proved together with a detailed asymptotic analysis of the solution. The second part of the paper gives a uniqueness result, showing that any other compact, convex, rotationally symmetric ancient solution which is contained in the same slab must coincide with the one constructed here.

The results of the paper are new and interesting. The existence of an ancient solution looking like a rotated Angenent oval was commonly believed to be true in the community of mean curvature flow. However, the only available result in this direction was due to X.-J. Wang in [18], who proved the existence of ancient solutions contained in a slab. In the present paper, the authors use a more explicit method and provide a description of the asymptotic profile which was not given in [18]. The statement about uniqueness is probably even more interesting, since it is one of the few available results until now on the general structure of ancient solutions in higher dimensions.

In my opinion, the results of the paper are an important and original contribution to the understanding of ancient solutions of mean curvature flow. The paper analyzes the class of collapsed solutions, which had not been treated in detail before, and gives a comprehensive existence and uniqueness result in the rotationally symmetric case. The authors cannot follow the technique of the noncollapsed case as in [12], and give a direct

proof based on simple and explicit computations, but which often involves subtle arguments and estimations, especially in the uniqueness part. I think that the results of this paper are of great interest for the general theory of mean curvature flow, and I believe that they fully meet the quality standards of JDG.

The paper is very clearly written and well organized, and I have not detected misprints or passages where corrections are needed, except for one spot: it seems to me that statement (1) of Theorem 1.1 should hold for  $\lambda \rightarrow +\infty$  rather than  $\lambda \rightarrow 0$ . Except for this minor remark, my suggestion is that the paper can be accepted in its present form.