



King's Research Portal

DOI:

[10.3390/nu14051062](https://doi.org/10.3390/nu14051062)

Document Version

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Terblanche, E., Hills, J., Russell, E., Lewis, R., & Rose, L. (2022). Dietetic-Led Nutrition Interventions in Patients with COVID-19 during Intensive Care and Ward-Based Rehabilitation: A Single-Center Observational Study. *Nutrients*, 14(5), Article 1062. Advance online publication. <https://doi.org/10.3390/nu14051062>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Dietetic-led nutrition interventions in patients with COVID-19 during intensive care and ward-based rehabilitation: single-centre observational study

Ella Terblanche ^{1,2,*}, Jessica Hills ¹, Edie Russell ¹, Rhiannon Lewis ¹ and Louise Rose ^{2,3}

¹ Dietetics Department, St Georges University Hospitals NHS Foundation Trust, London, UK; jessica.hills@uon.edu.au (J.H.), edie.russell@stgeorges.nhs.uk (E.R.), r.lewis3@rbht.nhs.uk (R.L.)

² Florence Nightingale Faculty of Nursing, Midwifery and Palliative Care, King's College London; louise.rose@kcl.ac.uk

³ Adult Critical Care, Guy's and St Thomas' NHS Foundation Trust

* Correspondence: ella.terblanche@kcl.ac.uk

Abstract: Background: This study described dietitian-led nutrition interventions for patients with COVID-19 during ICU and ward-based rehabilitation. As knowledge of COVID-19 and its medical treatments were evolving, dietetic-led interventions between surge 1 (S1) and surge 2 (S2) were compared. Methods: A prospective observational study was conducted of patients admitted to the ICU service in a large academic hospital (London, UK). Clinical and nutrition data were collected during 1st (March-June 2020) N= 200 and 2nd (November 2020-March 2021) N= 253 COVID-19 surges. Results: 453 patients were recruited. All required individualized dietetic-led interventions during ICU admission as the ICU nutrition protocol didn't meet nutritional needs. Feed adjustments for deranged renal function (p=0.001) and propofol calories (p=0.001) were more common in S1 whereas adjustment for gastrointestinal dysfunction was more common in S2 (p=0.001). One third of all patients were malnourished on ICU admission, all lost weight in ICU; mean (SD) total percentage loss of 8.8% (6.9%) Further weight loss was prevented over the remaining hospital stay with continued dietetic-led interventions. Conclusions: COVID-19 patients have complex nutritional needs due to malnutrition on admission and ongoing weight loss. Disease complexity and evolving nature of medical management required multifaceted dietetic-led nutritional strategies which differed between surges.

Citation: Lastname, F.; Lastname, F.; Lastname, F. Title. *Nutrients* **2022**, *14*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Lastname

Received: date
Accepted: date
Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: COVID-19; Intensive Care; Dietitian/Dietician; Nutrition; Malnutrition; weight loss

1. Introduction

In March 2020, the exponential increase in intensive care unit (ICU) admissions in the United Kingdom (UK) due to the COVID-19 pandemic required significant planning and restructuring of dietetic services to ensure safe and effective nutrition provision [1]. ICU dietitians were faced with multiple challenges including how to best provide nutrition support for patients with an unknown disease; rapidly train redeployed dietitians inexperienced in ICU nutrition; prioritize dietetic-led nutrition interventions; and manage the logistics of shortages of enteral feed, feed pumps, and ancillaries. Critically ill patients with COVID-19 appeared nutritionally complex with no international consensus on optimal nutritional management.

COVID-19 patients frequently present with malnutrition [2, 3]. This is due to pre-existing chronic disease associated with underlying poor nutritional intake, combined with further decline due to common COVID-19 symptoms including gastrointestinal dysfunction and loss of taste and smell [2]. Patients with severe COVID-19 pneumonia exhibit a marked hyperdynamic state with persistent pyrexia leading to hypermetabolism and

protein catabolism [4, 5]. Furthermore, enteral nutrition (EN) intolerance was reported due to gastrointestinal symptoms, refractory hypoxemia requiring prone positioning, hypotension or shock requiring the use of vasopressors, and the progression of multiple organ failure [1, 6-8].

Nutritional guidelines were written rapidly based on experiential learning and knowledge gained from dietitians in other countries managing critically ill patients with COVID-19 in addition to prior dietetic knowledge of managing patients with severe respiratory failure [1, 6-8]. Feeding protocols were devised to simplify nutrition delivery and ensure consistency of nutrition interventions [6-9] at a time when EN was perceived difficult to achieve and not a medical priority [9].

The aim of this study was to describe dietitian-led nutrition interventions for patients with COVID-19 during ICU admission and ward-based rehabilitation in a large tertiary ICU service in a single center. As our knowledge of COVID-19 and its medical treatments were evolving, we compared dietetic-led nutrition interventions between surge one (S1) and surge two (S2) in the UK.

2. Materials and Methods

2.1. Study design, setting and sample

A prospective observational study was conducted enrolling critically ill patients with COVID-19 admitted to an ICU (66 beds outside of pandemic conditions, 148 maximum bed number during pandemic) in a large academic hospital in London, UK. Data were collected during S1 (March-June 2020) and S2 (November 2020-March 2021).

All adult (≥ 16 years) patients with COVID-19 were included who received advanced respiratory support defined as invasive ventilation via endotracheal or tracheostomy tube and required EN or parenteral nutrition (PN) for longer than 48 hours in ICU. Data were collected from ICU admission to hospital discharge.

2.2. Nutritional Interventions

To support increased ICU patient numbers, dietitians were redeployed from other clinical areas, equating to one dietitian to 25 ICU patients. The primary treatment aim was to commence nutrition using the newly devised local hospital COVID-19 ICU feeding protocol with EN within 48 hours of admission. The protocol advised feeding to start with a high protein EN product, at 30ml/hr. for six hours. After which time a gastric residual volume (GRV) should be obtained. If the volume is less than 500ml (or 300ml if in the prone position), the rate of feeding is increased to meet a target rate based on the patient's weight, (Actual or ideal if BMI > 25kg/m²). Ideal body weight (IBW) was based on the patient's height calculated to BMI of 25kg/m² [10]. If EN was not tolerated sufficiently as defined in the protocol, PN was commenced. As per usual practice prior to COVID-19, all patients were screened by a dietitian each morning and only patients whose nutritional needs were not met using the ICU feeding protocol underwent dietetic assessment within 48 hours of commencing nutrition. Treating dietitians set energy and protein targets as per European Society for Parenteral and Enteral Nutrition (ESPEN) guidelines [7, 10], with target energy recommended at 20kcal/kg/day to avoid overfeeding in the early phase and increased to 30kcal/kg/day or more to facilitate rehabilitation after the acute phase of critical illness. Actual weight was used for patients with a BMI less than 25 kg/m², IBW for those BMI 25-30 kg/m² and an adjusted body weight for those with BMI more than 30kg / m². Indirect calorimetry was used where appropriate to aid prediction of energy needs during the recovery phase on ICU. Nutritional data was collected on type of EN, use of protein supplementation, reasons for dietetic-led nutrition interventions during the ICU admission, on ICU discharge, during ward admission and at hospital discharge.

2.3. Data Collection

Baseline demographics were collected from the medical record including pre-existing comorbidities, severity of illness using the Acute Physiology and Chronic Health Evaluation (APACHE) II score, anthropometric measurements, and malnutrition risk as

defined by Global Leadership Initiative on Malnutrition [11] and European ICU criteria [10]. Outcome data included ICU mortality, and length of stay in ICU and hospital.

Anthropometric measurements included Body Mass Index (BMI) calculated using ICU admission body weight (kg) and height (m). When a recent accurate weight was not available from medical records, we measured patient weight using a hoist or patient transfer scales or obtained a reported weight from a family member. When height was not available, it was estimated from ulnar length measurement and converted into an estimated height [12]. ICU survivors were weighed at ICU and hospital discharge using sitting, standing or hoist scales. Change in weight during ICU and hospital stay was determined in kilograms and percentage total weight loss. Percentage weight loss $\geq 5\%$ was considered clinically significant and was used to diagnose malnutrition [11].

2.4. Data Analysis

Categorical data is presented as frequencies and proportions and continuous data as means and standard deviations (SD). Proportions were compared using chi-squared or Fishers exact tests (depending on cell size) and continuous data using 2-sample independent t-test or the Wilcoxon rank-sum (Mann-Whitney) test. All tests were 2-tailed with $p \leq 0.05$ considered statistically significant. Statistical analyses were done using Stata version 17. The STROBE reporting guidelines for observational studies were followed

2.5. Ethical Considerations

The study was approved as a service evaluation by the National Health Service Health Research Authority and the Research and Development Service at St George's University Hospitals NHS Foundation Trust, London, UK. Registration number AUDI000637. Consent was waived as the project was approved as a service evaluation.

3. Results

There were 453 critically ill patients with COVID-19 included 200 patients were admitted during S1 (March-June 2020) and 253 during S2 (November 2020-March 2021). Baseline characteristics are shown in Table 1. Figure 1 presents patient numbers from admission to discharge.

Table 1. Baseline characteristics and clinical outcomes (whole cohort and comparison between patients in S1 and S2).

Patient characteristics	All patients N = 453	S1 n = 200	S2 n = 253	p value
Age (years), mean (SD)	61 (12.4)	57.9 (12.7)	62.5 (12.0)	0.001
Male gender, n (%)	315 (70)	135 (67)	180 (71)	0.44
Weight (kg), mean (SD)	84 (20)	86 (21)	84 (19.4)	0.32
BMI (kg/m ²), mean (SD)	29 (6.3)	29 (6.5)	29 (6.1)	0.82
Ethnicity, n (%)				
White	114 (25)	46 (23)	68 (27)	0.34
Asian	96 (21)	41 (21)	55 (21)	0.85
Black	71 (16)	36 (18)	35 (14)	0.22
Other ethnic groups	48 (11)	23 (12)	25 (10)	0.57
Not stated	124 (27)	54 (27)	70 (28)	0.87
Comorbidities, n (%)				
Nil or 1	90 (20)	50 (25)	40 (16)	0.02
More than 2	362 (80)	150 (75)	212 (84)	
APACHE II, mean (SD)	15.1 (6.8)	15.5 (7.3)	14.8 (6.5)	0.32
ICU mortality, n (%)	253 (55)	107 (54)	146 (57)	0.37
ICU length of stay (days), mean (SD)	20 (18)	18.1 (14.4)	22.3 (21.2)	0.02

Total hospital stay (days), mean (SD)	35.6 (21)	33.8 (19.7)	37.4 (22.85)	0.2
---------------------------------------	-----------	-------------	--------------	-----

Acute Physiology and Chronic Health Evaluation II score (APACHE). All values are mean (SD).

125
126
127




Figure 1. Patient flow diagram Key- S1 = surge one, S2 = surge two.

128
129

3.1. Patient characteristics

130

S1 patients were younger than in S2: mean (SD) 57.9 (12.7) compared to 62.5 (12.0) years ($p=0.001$) and had fewer comorbidities ($p=0.02$). Length of ICU stay was longer in S2 ($p=0.019$) (Table 1). Of the 453 patients, 167 (37%) were malnourished on ICU admission; 64 (32%) in S1 and 103 (40%) in S2.

3.2. Dietitian-led nutrition interventions

Individualized dietetic-led nutrition interventions were required by all 453 (100%) patients during ICU admission as nutritional needs were not met using the standardized ICU nutrition protocol. Mean (SD) time to first dietetic-led nutrition interventions was 2.3 (1.2) days in S1 and 2.9 (2.3) days in S2. Patients received a mean (SD) of 5.2 (4.5) dietetic interventions during the ICU stay. Patients had a similar number of dietetic-led nutrition interventions during their ICU stay in S1 and S2 (mean 5.2 (4.2) compared to 5.3 (4.8) times). PN was only required for six (1%) patients during S2 and none in S1.

More patients required feed adjustment for calories derived from propofol, impaired renal function or changes in fluid or electrolyte status in S1 whereas more patients needed adjustments for gastrointestinal dysfunction in S2 (Table 2).

Table 2. Percentage of patients requiring specific dietetic-led nutrition interventions in ICU.

Intervention	N = 453 n (%)	S1 N = 200 n (%)	S2 N = 253 n (%)	p value
Feed adjustment to meet energy needs for the different metabolic phases	337 (74)	141 (71)	196 (78)	0.09
Feed adjustment to account for calories derived from propofol sedation >15ml/hr (360kCals/24hrs)	248 (55)	144 (72)	104 (41)	0.001
Feed adjustment for gastrointestinal dysfunction	154 (34)	47 (24)	107 (42)	0.001
Transition from EN to oral diet	146 (32)	62 (31)	84 (33)	0.62
Feed adjustment due to changes in renal function, fluid status, or electrolyte balance	120 (26)	73 (37)	47 (19)	0.001
Feed adjustment to allow feed interruption for drug absorption of medication given via the enteral route	18 (4)	6 (3)	12 (5)	0.34

All values are patient numbers and (%).

Most patients required high-protein enteral feeds during ICU admission; with similar proportions in S1 and S2 (184, 92% vs 236, 93%). There was no difference in the proportion of patients requiring concentrated feeds (39, 20% in S1 vs 45, 18% in S2). More peptide feeds were used in S1 (26, 13%) compared to S2 (17, 7%) ($p=0.03$). Protein supplementation also increased from S1 (98, 49%) to S2 (174, 69%) ($p=0.001$). During S1, first choice of feed was not available due to supplier production shortages in 61 (30%) patients, shortages were not experienced during S2.

3.3. Weight loss

Mean (SD) weight loss over the ICU admission was 7.9kg (6.8kg), equivalent to a mean (SD) total percentage loss of 8.8% (6.9%) indicating clinically significant malnutrition. Mean (SD) weight loss over the total hospital stay was 7.5kg (6.6kg) suggesting no further weight loss occurred after ICU discharge. Details of weight loss according to surge are shown in Table 3.

Table 3. Weight loss.

	All N = 160	S1 N = 78	S2 N = 82	p value
ICU admission weight (kg)	85 (20.1)	86 (21)	84 (19.4)	0.32
ICU admission BMI (kg/m ²)	29 (6.3)	29 (6.5)	29 (6.1)	0.82
ICU weight loss (kg)	7.9kg (6.8)	7.8 (7.8)	8.1 (5.9)	0.80
ICU weight loss %	8.8 (6.9)	8.5 (7.7)	9.0 (6.3)	0.65
Percentage ICU weight loss				
N (%)				
< 5 %	58 (35)	30 (41)	36 (40)	0.22
5-10%	57 (34)	27 (36)	27 (30)	
> 10%	53 (32)	22 (30)	26 (29)	
Total weight loss (kg) from ICU admission to hospital discharge	7.5 (6.6)	8.0 (7.4)	7.3 (6.1)	0.70

All values are mean (SD).

3.4. Dietetic-led nutrition interventions after ICU discharge

Ward dietitians received a dietetic handover for all patients discharged alive from ICU ($n = 177$). Most patients (160, 90%) received ward-based dietetic-led nutrition interventions with similar numbers in S1 (78, 89%) and S2 (82, 92%). The 17 patients not reviewed were discharged home before the ward dietitians were able to assess.

Upon discharge to the ward, patients were reviewed earlier during S1 compared to S2, within a mean (SD) of 1.9 (1.3) vs. 2.4 (1.8) days ($p=0.04$); but at similar frequencies (2.8 (2.5) times over 9.9 (8.0) days vs. 3.3 (2.4) times over 13 (8.5) days).

Table 4 details the type of ward-based nutrition support received. Exclusive or supplementary (to oral diet) EN was prescribed for 106 patients (52 (67%) in S1 and 54 (66%) in S2) for a mean of 6 (5.5) days (similar duration in both surges). More high-protein feed was used on the ward in S2 (33, 61%) compared to S1 (9, 17%) ($p=0.0001$), whereas more high-energy feed was used in S1 (17, 33% vs. 7, 13%, $p=0.007$). Of the 106 patients receiving EN, nasogastric feeding was ceased on the ward without prior dietetic review on 58 occasions (54%). During ward admission, 117 (73%) patients required prescribed ready-to-drink oral nutritional supplements; 52 (66%) in S1 and 65 (79%) in S2. The most commonly prescribed product was a compact high-protein milkshake-style; 32 (62%) in S1 and 40 (62%) in S2.

Table 4. Nutrition support received on ward admission.

Nutrition intervention	N = 160	S1 N =78	S2 N = 82	p value
------------------------	---------	----------	-----------	---------

	n (%)	n (%)	n (%)	
Exclusive EN	36 (23)	14 (18)	22 (27)	0.17
Supplementary EN	70 (44)	38 (49)	32 (39)	0.21
Exclusive and supplementary EN combined	106 (66)	52 (67)	54 (66)	0.91
ONS	68 (43)	34 (44)	34 (42)	0.78
Texture modification	34 (21)	23 (29)	11 (13)	0.01
Diet alone	33 (21)	21 (27)	12 (14)	0.05

All values are patient numbers and (%). Enteral Nutrition (EN), Oral Nutrition Supplement Drinks (ONS).

Of the 453 patients, 175 (39%) survived to hospital discharge, of which 128 (73%) needed community dietetic follow-up. This was consistent across both surges with 63 (72%) respectively in S1 vs. 65 (74%) in S2. Community referrals reasons for each surge were continued EN (15, 12%), oral nutritional supplements (70, 55%), and healthy eating advice (43, 34%).

4. Discussion

In this large cohort of patients with COVID-19, a standardized feeding protocol was insufficient to meet nutritional needs with individualized dietetic-led nutrition interventions required for all patients during ICU admission. Significant differences were observed in dietetic-led nutrition interventions in S1 and S2. More patients required feed adjustment for calories derived from propofol, impaired renal function, and fluid and electrolyte adjustments in S1, whereas patients more commonly needed adjustments for gastrointestinal dysfunction in S2. Over one third of patients were malnourished on ICU admission and patients lost an average of 7.9kg over the ICU stay. Further weight loss was prevented over the remaining hospital stay with continued dietetic-led nutrition interventions. Only 39% of the cohort survived to hospital discharge, with 73% of these requiring further dietetic interventions in the community.

In our COVID-19 cohort, all required individualized dietetic-led nutrition interventions during ICU admission suggesting critically ill patients with COVID-19 are nutritionally complex. This proportion is markedly higher than non-COVID patients for whom the need for individualized dietetic led nutrition interventions is reported to range from 50 to 70% [13, 14]. Differences in dietetic led nutrition interventions required in pandemic S1 and S2 reflect changes in medical management as knowledge of the disease evolved. Furthermore, in our institution, propofol shortages in S2 requiring alternate opioid sedation agents obviating the need for EN manipulation to avoid overfeeding were experienced. Fluid restriction commonly employed in S1 was not used in S2 as evidence emerged that it was not beneficial [15]. Increased gastrointestinal dysfunction and EN intolerance in S2 might be attributed to two changes in clinical practice. Firstly, there was an increased use of prone positioning which can present unique feeding challenges due to large gastric residual volumes, vomiting, and aspiration of gastric contents [16]. Secondly, increased use of opioid-based sedation due to propofol shortages observed in this study may have contributed to the cycles of constipation and laxative induced diarrhoea observed [3, 17].

In our patient cohort a mean weight loss of 7.9kg over the ICU stay was identified. Two-thirds of patients lost more than 5% of body weight; one-third lost more than 10% indicating severe malnutrition. Disease severity and prolonged ICU stay were likely contributing factors. Weight loss and malnutrition contribute to worse functional ability following ICU discharge in patients with COVID-19 [2, 3]. A substantial proportion of patients surviving to hospital discharge required further dietetic led nutrition interventions in the community was also found.

Despite all patients experiencing weight loss in ICU, no further weight loss occurred before hospital discharge which may be attributed to ongoing dietetic led nutrition

interventions. Of the patients transferred to the ward, 90% received ward-based dietitian-prescribed individualized nutrition interventions. Continuity of nutritional interventions can be problematic during transition from ICU to the ward [18]. Structured dietetic hand-over between ICU and ward dietitians to improve transfer of nutritional history and meet nutritional needs for recovery were used [1]. Additionally, more patients received high-protein feeds on the ward during S2 as ward dietitians became more familiar with treating COVID-19 patients and as guidelines became available. A high protein diet is recommended due to the catabolism experienced during critical illness and thought to aid recovery [1].

To our knowledge, this is the largest prospective cohort study describing dietetic-led nutritional interventions provided to patients with COVID-19 from ICU admission to ward discharge. Our study has limitations. First, a single-center limits generalizability. Second, the observational design means no assumptions about causality can be made and therefore cannot determine if dietetic led nutrition interventions influenced clinical outcomes. Third, it was not possible to make comparisons to dietetic-led nutritional interventions for patients without COVID-19 at this time as there were very few patients admitted without COVID-19. Finally, due to pandemic dietetic working conditions data on amounts of energy and protein received could not be captured.

5. Conclusions

In this large prospective cohort of patients with COVID-19, all patients required individualized dietetic led nutrition interventions during ICU admission as the standardized ICU nutrition protocol did not meet nutritional needs. The complexity and evolving nature of medical management necessitated multifaceted dietetic led nutrition interventions which differed between surges. Over one third of patients were malnourished on ICU admission and all patients lost weight in the ICU. Most patients were nutritionally compromised at ICU discharge and required ongoing dietitian-led individualized nutrition interventions in the ward which prevented further weight loss over the remaining hospital stay. Based on our results future studies are needed to determine if individualized nutrition support provision led by the dietitian, compared to standard care, increases the adequacy of nutritional delivery, and improves clinical outcomes.

Author Contributions: Study conception and design ET. Statistical analysis and interpretation of data, E.T., J.H., E.R., R.L., L.R. Drafting the manuscript, reviews and revisions, E.T., J.H., E.R., R.L., L.R. Study supervision LR.

Funding: This research received no external funding

Institutional Review Board Statement: The study was approved as a service evaluation by the National Health Service Health Research Authority and the Research and Development Service at St George's University Hospitals NHS Foundation Trust, London, UK. Registration number AUDI000637.

Informed Consent Statement: Consent was waived.

Data Availability Statement: Not applicable.

Acknowledgments: We wish to thank Meg Skelton, Jen Wetherden and James Jackson for their clinical care and assistance in collecting data for this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. BDA. British Dietetic Association: Critical Care Specialist Group. Guidance on management of nutrition and dietetic services during the COVID-19 pandemic. 2020. <https://www.bda.uk.com/resource/critical-care-dietetics-guidance-covid-19.html> [accessed 23.10.21].
2. Bedock D, Bel Lassen P, Mathian A, Moreau P, Couffignal J, Ciangura C, et al. Prevalence and severity of malnutrition in hospitalized COVID-19 patients. *Clin Nutr.* 2020;40:214-9.

3. Eden T, McAuliffe S. Critical care nutrition and COVID-19: a cause of malnutrition not to be underestimated. *BMJ Nutr Prev Health*. 2021;4(1):342-7. 278
279
4. Niederer LE, Miller H, Haines KL, Molinger J, Whittle J, MacLeod DB, et al. Prolonged progressive hypermetabolism during COVID-19 hospitalization undetected by common predictive energy equations. *Clin Nutr*. 2021;45:341-350 280
281
282
5. Whittle J, Molinger J, MacLeod D, Haines K, Wischmeyer PE, Group L-CS. Persistent hypermetabolism and longitudinal energy expenditure in critically ill patients with COVID-19. *Crit Care*. 2020;24(1):1-4 283
284
285
6. Chapple LS, Fetterplace K, Asrani V, Burrell A, Cheng AC, Collins P, et al. Nutrition management for critically and acutely unwell hospitalised patients with coronavirus disease 2019 (COVID-19) in Australia and New Zealand. *Nutr Diet*. 2020;77(4):426-36. 286
287
288
7. Barazzoni R, Bischoff SC, Breda J, Wickramasinghe K, Krznaric Z, Nitzan D, et al. ESPEN expert statements and practical guidance for nutritional management of individuals with SARS-CoV-2 infection. *Clin Nutr*. 2020;39(6):1631-8. 289
290
291
8. Martindale R, Patel JJ, Taylor B, Arabi YM, Warren M, McClave SA. Nutrition Therapy in Critically Ill Patients With Coronavirus Disease 2019. *J Parenter Enteral Nutr*. 2020;44(7):1174-84. 292
293
9. de Watteville A, Montalbano F, Wozniak H, Collet T-H, Jaksic C, Le Terrier C, et al. Impact of nutritional therapy during the first wave of the COVID-19 pandemic in intensive care patients: a retrospective observational study. *Clin Nutr*. 2021. <https://doi.org/10.1016/j.clnu.2021.05.024> 294
295
296
10. Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr*. 2019;38(1):48-79. 297
298
11. Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical nutrition community. *J Cachexia Sarcopenia Muscle*. 2019;10(1):207-17. 299
300
301
12. BAPEN. British Association of Parenteral and Enteral Nutrition. Malnutrition Universal Screening Tool (MUST) http://www.bapen.org.uk/pdfs/must/must_full.pdf. [accessed 23.10.21]. 302
303
13. Rothpletz-Puglia P, Mena MD. Emergency preparedness in dietetics during a pandemic: Lessons learned from an ICU dietitian during Covid-19. *Internet Journal of Allied Health Sciences and Practice*. 2021;19(3):12. 304
305
306
14. Minnelli N, Gibbs L, Larrivee J, Sahu KK. Challenges of maintaining optimal nutrition status in COVID-19 patients in intensive care settings. *J Parenter Enteral Nutr*. 2020;44(8):1439-46. 307
308
15. Jha AK, Kulkarni SG. Evolution of COVID-19 management in critical care: review and perspective from a hospital in the United Kingdom. *Acute Crit Care*. 2021;36(1):1-14. 309
310
16. Behrens S, Kozeniecki M, Knapp N, Martindale RG. Nutrition support during prone positioning: An old technique reawakened by COVID-19. *Nutrition in Clinical Practice*. 2021;36(1):105-9. 311
312
313
17. Chappell D, Rehm M, Conzen P. Opioid-induced constipation in intensive care patients: relief in sight? *Crit Care*. 2008;12(4):1-2. 314
315
18. Ghorbanzadeh K, Ebadi A, Hosseini M, Madah SSB, Khankeh H. Challenges of the patient transition process from the intensive care unit: a qualitative study. *Acute and Critical Care*. 2021;36(2):133. 316
317
318
319
320
321