



## King's Research Portal

### *Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

### *Citation for published version (APA):*

Ide, M., Karimova, M., & Setterfield, J. (2022). Oral health, antimicrobials and care for patients with chronic oral diseases – a review of knowledge and treatment strategies. *Frontiers in Oral health*, 3, Article 866695. <https://www.frontiersin.org/articles/10.3389/froh.2022.866695/full>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Oral Health, Antimicrobials and Care for Patients With Chronic Oral Diseases – A Review of Knowledge and Treatment Strategies

Mark Ide\*, Malika Karimova and Jane Setterfield

Centre for Host-Microbiome Interactions, Faculty of Dentistry, Oral and Craniofacial Sciences, King's College London, London, United Kingdom

## OPEN ACCESS

### Edited by:

Hatice Hasturk,  
The Forsyth Institute, United States

### Reviewed by:

Mia Rakic,  
Complutense University of  
Madrid, Spain  
Matthew Mason,  
University of North Carolina at Chapel  
Hill, United States

Darko Bozic,  
University of Zagreb, Croatia

### \*Correspondence:

Mark Ide  
mark.ide@kcl.ac.uk

### Specialty section:

This article was submitted to  
Oral Infections and Microbes,  
a section of the journal  
Frontiers in Oral Health

**Received:** 31 January 2022

**Accepted:** 04 April 2022

**Published:** 07 June 2022

### Citation:

Ide M, Karimova M and Setterfield J  
(2022) Oral Health, Antimicrobials and  
Care for Patients With Chronic Oral  
Diseases – A Review of Knowledge  
and Treatment Strategies.  
*Front. Oral. Health* 3:866695.  
doi: 10.3389/froh.2022.866695

Periodontal and chronic oral mucosal diseases are significant life impacting conditions which may co-exist and synergistically act to cause more severe and widespread oral pathology with enhanced challenges in effective management. Clinicians regularly observe these effects and struggle to effectively manage both problems in many patients. There is limited understanding of many basic and applied scientific elements underpinning potentially shared aetiopathological features and management. Recent developments in translational science provide an opportunity to greater improve knowledge and subsequently care for patients with these problems.

**Keywords:** oral health, periodontitis, gingivitis, microbiome, pemphigus vulgaris, lichen planus, mucous membrane pemphigoid, mucosal disease

## INTRODUCTION

Periodontal diseases and chronic oral mucosal diseases (OMD) both have an impact on comfort, function, and quality of life [1–11]. Beyond this they may co-exist in some patients and although there is limited knowledge of the interactions between these conditions, there are good reasons to believe that they may act in a bidirectional manner to both exacerbate their severity and impact and to complicate management.

## Periodontal Diseases and Management

Periodontal diseases are processes driven by the patient's inflammatory response to the presence and accumulation of bacterial plaque and form a major part of the global health burden [12–15]. They are known to result in significant impacts on function, wellbeing, other disease states and Quality of Life for the majority of populations in many states. Management of these problems is generally simple and based around control of local and systemic risk factors, with good home care the cornerstone for success, and supportive professional care where appropriate [16]. Outcomes from treatment are further enhanced by management of associated secondary risk factors including aspects of health and behavioral factors such as control of diabetes, smoking (and possibly vaping) cessation and, where appropriate, modification of other drug prescriptions to eliminate side effects [17].

Periodontal diseases can be subcategorised based on the nature and degree of changes seen and damage caused. The majority of disease falls largely into one of two groups [18], namely gingivitis and periodontitis.

Gingivitis is a condition in which there is inflammation of the marginal soft tissues around the neck of the tooth. It is completely reversible but is also extremely common. There are associated shifts in the local microflora in the presence of gingivitis [19–21]. There is no irreversible damage to supporting tissues, however, it is possible for this condition to subsequently progress to periodontitis [22]. Gingivitis can be controlled by simple home care measures, supported by professional interventions at an early phase with occasional review [23, 24].

Periodontitis is a more advanced condition in which there is (frequently) irreversible destruction of supporting soft and hard tissues around teeth. This can lead to mobility, acute infection, pain and ultimately tooth loss. Again there are recognized shifts in microflora in periodontitis, which may be further affected by recognized systemic factors [25]. Periodontitis management is primarily focussed on stopping progression, with clinically significant regeneration of lost tissues only being predictable in an increasing but still limited number of scenarios [16, 26].

Although periodontitis may be expected in all adult humans in the absence of oral hygiene, the onset, extent and rate of development is known to vary between individuals [27, 28], and this is equally true even if “normal” oral hygiene is practiced [29]. Patients with particularly higher levels of susceptibility to periodontitis are required to deliver a very high standard of plaque control supported by regular professional maintenance in order to prevent further ongoing inflammation and tissue damage [30–33]. However in the presence of good maintenance, even questionable teeth can be maintained for a long period of time [30, 34–37].

Likewise, regeneration of tissues is presently only feasible in patients with exemplary oral hygiene where other risk factors have also been managed [16, 38, 39].

## Oral Mucosal Conditions

Oral mucosal diseases form a group of chronic predominantly immune mediated disease, such as lichen planus (OLP), pemphigus vulgaris (PV), bullous pemphigoid (BP), dermatitis herpetiformis, Epidermolysis bullosa acquisita (EBA), mucous membrane pemphigoid and linear IgA bullous disease [40]. The pathogenesis of these diseases are not completely clear, but autoantibodies play critical roles in their pathogenesis [41–43]. The majority of these diseases are rare. Clinically they are characterized by formation of blisters and erosions in oral mucosa, often presenting as desquamative gingivitis. Erosive gingival lesions can be the source for the microbial accumulation and may hence trigger periodontal tissue breakdown [44]. They often clinically overlap with each other; therefore, immunological and histopathologic tests are requested for differential diagnosis.

## The Microbiome and Aspects of Periodontal Disease and Management

The role of plaque as an aetiological factor in, and its removal in management of, periodontal diseases has been known for hundreds of years. However, in the last 40 years with the advent of new technologies our understanding the role of the oral microflora has expanded significantly. Regardless there are still large gaps in our knowledge, such that although we understand

which simple measures are likely to work in many patients, it is still not fully clear why, or why their efficacy varies between individuals. Individual variation in disease susceptibility and rate of disease progression has been attributed to genetic or epigenetic variations in host inflammatory response, including neutrophil function, cytokine production and macrophage activity [45–50], to specific microbial infection (such as with clones of *Aggregatibacter actinomycetemcomitans*) [51] or to a combination of these, whereby one leads to the other with debate around which happens first. Current concepts of the microbiology of periodontal disease, and more specifically periodontitis, relate to the presence of groups of putative periodontal pathogens described as “complexes” (based on their pathogenic potential, association with each and presence in diseased sites). These concepts first arose with the advent of early nonculture-based means of microbial assessment such as “checkerboard” DNA-DNA hybridisation [52, 53] and have been supplemented and superseded by more sophisticated methods such as qt-PCR [54] and high throughput gene sequencing and complex analysis [55]. Supplemented by metabolomic analysis and other approaches, a massive volume of potential information is available to better understand the pathobiology of both periodontal and mucosal diseases.

More recently the concept of dysbiosis induced by various environmental shifts and “keystone pathogens” has gained support [56–58]. However, this does not fully explain all aspects of what is without doubt a complex series of interactions between the environment created by the host (and potential variations) combined with those within the microbiome [59, 60]. This may lead to patient or even site-specific disruptions responsible for disease activation or progression, with no universal pattern seen across all hosts [61]. It has also been suggested that such modified dysbiotic microbiota can form stable communities that may additionally be suppressed but not eliminated by antimicrobial therapies [62]. There is an extensive literature confirming that regardless of OMD status, development and maturation of dental plaque can potentiate local inflammatory change and hence likely periodontal disease progression [63–65].

This picture is further muddled by the complex nature of the oral microbiota [66], and the belief that many microorganisms present have not yet been formally identified or characterized. In addition, the ability to transfer resistance genes and evade antimicrobials of organisms within a plaque biofilm also is likely to impact on future strategies for patient management especially if these agents are used for other purposes [67].

A further central aspect of periodontitis management relates to modifiable risk factors that further influence susceptibility to disease by changing individual host responses. These include smoking and possibly vaping, uncontrolled diabetes and possibly obesity, medications that can change cellular activity in gingival tissues or drugs which result in oral dryness or xerostomia [16, 68]. Changes in systemic health, lifestyle or medications, or variation in host response can therefore relate to and influence the oral microbiome, leading to specific individual patterns [69].

Current consensus for the management of periodontal diseases is based on a foundation of effective daily self-performed plaque removal and/or biofilm disruption combined with

professional interventions to complement or facilitate this [68]. The effectiveness of this essentially non-specific approach has been confirmed by national and international guidelines for care [16, 70–73].

Whilst chemical means of plaque control are supported in certain circumstances, they are not seen as a mainstay of longer-term therapy for the majority of patients with periodontitis due to either limited efficacy and / or the presence of a range of undesirable complications. Even so it has been established that use of common means of chemical plaque control such as chlorhexidine used topically as a mouthwash were associated with further large shifts in microbiome biomass as well as diversity [20, 21, 69, 74].

## The Microbiome and Aspects of Chronic Oral Mucosal Disease

The microbiome of oral mucosal diseases has been investigated for the last couple of decades [75]. Few studies focus on lichen planus, but one describes the increase of periodontal pathogens such as *Porphyromonas gingivalis*, *Aggregatibacter actinomycetemcomitans*, *Treponema denticola*, *Prevotella intermedia* and *Tannerella forsythia* in subgingival plaque of OLP patients [76], and another describes changes in both richness and diversity of subgingival plaque associated with the presence of OLP when co-existing with periodontitis [77]. The role of the microbiome in autoimmune bullous disease (AIBD) is less investigated. One study on skin microbiome of PV and BP patients did not show significant differences in a- and b-diversity in comparison with healthy controls [78]. However, another study of the gut microbiome of PV patients showed increases in *Flavonifractor* genera, which was correlated with increased levels of chronic inflammatory cytokines. It supported the concept that the dysbiosis in the gut may trigger inflammatory processes and cause the development of disease [79]. This corresponds to research suggesting that initially healthy mice showed a reduction in local microbial richness after immunization with EBA antibodies [80]. However, limitations of all current studies include low patient numbers due to the relative rarity of these diseases, supporting the need to investigate these links further.

## Interactions Between Periodontal Diseases and Chronic Oral Mucosal Conditions – Bidirectional?

Soft tissue tenderness, ulcerations and sometimes scarring associated with many OMD cause many patients to struggle to achieve good mechanical oral hygiene. This then leads to more local plaque formation and increased levels of marginal inflammation. This can manifest as localized gingivitis but may also predispose to acceleration of gingival recession, periodontitis and ultimately tooth loss in those who are more susceptible to this condition [44, 81–84].

Whilst we do not yet fully understand the true relationship, it appears feasible that the presence of active periodontitis and associated microorganisms could potentially exacerbate local mucosal disease. This could

occur by locally enhanced inflammation and/or modulation of autoimmune responses, either alone [85] or facilitated by superinfection of mucosal lesions by the proteolytic organisms often identified in periodontally involved sites [86, 87]. These organisms can act to directly or indirectly promote local inflammatory damage and hence could increase the severity and duration of mucosal disease [86–89].

## Management of Periodontal Problems for Patients With Mucosal Disease

Hopefully, adequate management of OMD would reduce the likelihood of periodontal complications for those with OMD. However patients may find themselves in a situation where the co-existence of these conditions may be almost synergistic.

Systematic reviews such as that by Figuero et al. [90] suggest that antiseptic agents including essential oils, chlorhexidine, stannous fluoride, cetylpyridinium chloride, sanguinarine, sodium hexametaphosphate and delmopinol, delivered via toothpastes or mouthwashes, can effectively reduce plaque and gingivitis levels and may as a result offer benefits for tissues in patients with gingivitis or periodontitis. However, these are still recommended as an adjunctive therapy [16] supporting mechanical oral hygiene. Mucosal diseases present challenges in oral hygiene and plaque control in regions with ulceration due to discomfort caused not only by physical brushing but also potentially by toothpaste constituents. It has been suggested that conventional formulations containing certain detergents may act to exacerbate some ulcerative conditions intraorally [91–95]. Unfortunately, chemical methods of plaque are often also not helpful. Those agents widely recognized as having clinical efficacy in management of periodontal diseases are often not well tolerated by patients, due to discomfort and pain potentially caused by various constituents such as flavorings, astringents, essential oils and preservatives. Even if diluted to the lowest effective concentration, many patients still report significant pain. Consequently, many patients will resort to using non-foaming, detergent-free agents and softer brushes, although removal of detergent may not impair plaque control [94, 95].

As a result, it is important to consider management of mucosal inflammation and lesion resolution whilst also trying to simultaneously improve plaque control. However, this can create challenges in compliance for the delivery of active agents by patients as well as risks of drug interactions. A better understanding of the exact relationships between these conditions and their management strategies may be expected to allow development of more effective approaches to therapy.

## Local Agents for Periodontal or OMD Management

Topical steroids are widely used in treatment of all clinical symptoms of OMD, with a prime aim of reducing pain and inflammation. These are generally applied topically, and include 0.1% triamcinolone acetonide in orabase, 0.05%

fluocinolone acetonide, 0.05% clobetasol propionate, or 0.05% halobetasol. A further approach has been intralesional injection of triamcinolone acetonide (20 µg/L) [96]. They often used as a first line drug with rare side effects and low cost [97].

Whilst topically applied agents have been proposed for periodontal management, these have largely been professionally applied, and have been shown to be of questionable efficacy and correspondingly low cost-effectiveness [98, 99]. The lack of high-quality evidence and significant clear clinical benefit has led to no clear recommendation for their use initial or longer term maintenance periodontal therapy [16]. There are no data related to the value of specific agents designed for periodontal management (such as topical slow release antimicrobial agents) in OMD management.

## Systemic Agents for Periodontal or OMD Management

Various systemic adjuncts have been proposed for periodontal management [100, 101], and these have aimed to either act as antimicrobials or to modulate host inflammatory responses, or both. There are limited data from RCTs comparing antimicrobial regimes, but recent network meta-analyses and reviews have suggested that some agents (primarily a combination of amoxicillin and metronidazole) when used at the end of a course of professional mechanical intervention can give enhanced clinical benefit in certain patient groups [100, 102], and this has led to guidelines and recommendations for clinical practice [16, 103] related to dosage, duration, timing and patient based criteria for use. Systemic antimicrobials are generally not considered a valid sole therapy for periodontal therapy in the absence of local biofilm control, a concept supported by *in vivo* studies [62].

Systemic agents are widely used in management of oral mucosal disease. Systemic steroids being used for AIBD because of its immunosuppressive effect to inhibit cytokines, which further prevents T cell activation [43, 104].

Azathioprine and mycophenolate mofetil are another two immunosuppressive drugs which reduce the number of circulating B and T lymphocytes, resulting in decreased immunoglobulin production and reduced IL-2 secretion, as well as inhibit the generation of cytotoxic T cells [105].

Rituximab is a biological agent directed against the B-lymphocyte-specific antigen CD20 which use where the treatment with other systemic agents such as those above has failed.

Even though immunosuppressive treatment is the first line treatment for autoimmune conditions, systemic antibiotics are an alternative treatment for oral mucosal disease. The positive effect of antibiotics can be explained by possible shared pathogenic mechanisms. The cross-talk activation of autoreactive B and T cells is explained by the mechanism of epitope spreading or molecular mimicry, may lead to immune cells are attacking host tissue, confusing them with the infection epitopes. One such antibiotic is dapsone, a sulfone derived antimicrobial agent with anti-inflammatory effects. This is often used as an adjuvant therapy for AIBD, and as a first line medication for LABD and dermatitis herpetiformis [106]. Tetracyclines have been

used in AIBD for their anti-inflammatory action [107, 108]. A recent study supported the hypothesis that oral doxycycline is as effective as oral prednisolone and that doxycycline is less likely to cause severe side effects [109]. Tetracyclines were shown to inhibit the activity of antiphospholipase, scavenge free radicals, and inhibit various matrix metalloproteinases, as well as impair lymphocyte activity [110].

Macrolides form another group of antibiotics described in the literature showing a positive effect on AIBD. Mensing and Krause [111] reported a decrease of hospitalization days, as well as side effects of steroid use, while using erythromycin in combination with methylprednisolone. Later, numerous studies showed that erythromycin alone showed improvement in patients with bullous pemphigoid and linear IgA disease [112, 113].

However, some antibiotics are capable of triggering autoimmune mucosal disease. The literature describes the case of AIBD induced with ciprofloxacin, levofloxacin, and penicillin groups of antibiotics [114], but the exact mechanism and the overall impact compared to benefit unclear. It has also been suggested that thiol/phenol drugs may activate acantholytic enzymes or stimulate keratinocytes to release proinflammatory cytokines and trigger acantholysis through other mechanisms [115, 116].

## Implications for Self Care and for Professionally Delivered Care

All of these challenges mean that OMD patients often require personalized diagnosis, focussed and sometimes more frequent professional follow up and support for effective primary and secondary prevention of periodontal disease, supported by OMD management. This would ideally be delivered by dental hygienists or therapists, although suitably trained oral health educators may have a role [117, 118].

However, studies have shown consistently and conclusively that the main determinant for long term success in periodontal management is effective oral hygiene [23, 24, 119, 120]. Without this, treatments are known to deliver poorer clinical outcomes with corresponding microbiological changes.

As outlined above, this is presently hard to achieve for many patients, but initial studies (with limited patient numbers) are promising in this regard, even without the use of chemical adjuncts to therapy [88, 121–125].

Whilst it is generally accepted that the presence and management of local and systemic risk factors influences both initial periodontal therapy and the frequency of subsequent maintenance, the presence of OMD per se is not part of most routine paradigms. In fact, the literature focusses on local evidence such as inflammation and gingival bleeding, and the presence of residual periodontal pockets, and these have been shown to be effective and form the basis for current treatment protocols [126]. Hence when considering the frequency of professional recall, currently we are driven by the periodontal rather than other signs. It is unclear if this is suitable, based on the limited evidence base, for this group of patients.



## What Do We Need to Know?

Many patients with oral mucosal disease suffer significant problems ranging from discomfort and dietary restriction to tooth loss, loss of function and difficulty undergoing dental rehabilitation. This can occur in the presence or absence of periodontal disease, but where there is ongoing periodontitis and associated complications, management challenges and disease impacts are considerably greater. The most recent World Workshop Classification of Periodontal diseases [127] does not list OMDs as separate mucosal conditions that affect the periodontium, or as risk factors or grade modifiers in making a periodontitis diagnosis [128], and this may be worthy of review.

The advent of genomic approaches to characterize the oral and more specifically periodontal and lesional microbiome offers a novel and exciting way to understanding better not only whether these are linked, but also may allow better understanding of how these factors relate to the extent, severity and persistence of OMD. Beyond pathobiology, a knowledge of how these characteristics may change following the use of drugs or other interventions designed to manage periodontal diseases, OMD or both may allow more efficient care. We are only at the very beginning of this journey but there is much promise and much to be delivered. Topics worthy of further investigation include cross sectional studies informing our understanding of basic associations, and longer term work to better understand the impact of OMP management protocols on both the local and systemic host response, the lesional and oral microbiome and clinical manifestations of both conditions. This could be

further enhanced by further development of disease classification to ensure that the epidemiological aspects of these problems can be better understood.

We are now aiming to deliver personalized, patient-centered care across medical and dental care. However, it is difficult to achieve a reasoned patient centered care pathway for OMD patients without having a better evidence-based understanding of the relationships as well as mechanisms underlying these, and the true nature of any bi-directional relationships, should they be found to exist. As a result there is a clear need for further basic and translational investigation work to better understand microbial and host-related aspects of the etiology and persistence of both conditions as well as related factors such as antimicrobial efficacy and resistance, in order to determine what treatments are most likely to work in which individuals [69]. This would be expected to offer the potential for major improvements in clinical outcomes and quality of life for this often overlooked but large body of patients.

## AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

## FUNDING

This work was supported by internal funds.

## REFERENCES

- Ng SK, Leung WK. Oral health-related quality of life and periodontal status. *Commun Dent Oral Epidemiol.* (2006) 34:114–22. doi: 10.1111/j.1600-0528.2006.00267.x
- Liu LJ, Xiao W, He QB, Jiang W-W. Generic and oral quality of life is affected by oral mucosal diseases. *BMC Oral Health.* (2012) 12:2. doi: 10.1186/1472-6831-12-2
- Brauchle F, Noack M, Reich E. Impact of periodontal disease and periodontal therapy on oral health-related quality of life. *Int Dent J.* (2013) 63:306–311. doi: 10.1111/idj.12042
- Durham J, Fraser HM, McCracken GI, Stone KM, John MT, Preshaw PM. Impact of periodontitis on oral health-related quality of life. *J Dent.* (2013) 41:370–6. doi: 10.1016/j.jdent.2013.01.008
- Tan H, Peres KG, Peres MA. Retention of teeth and oral health-related quality of life. *J Dent Res.* (2016) 95:1350–7. doi: 10.1177/0022034516657992
- Krain RL, Kushner CJ, Tarazi M, Gaffney RG, Yeguez AC, Zamalin DE, et al. Assessing the correlation between disease severity indices and quality of life measurement tools in pemphigus. *Front Immunol.* (2019) 10:2571. doi: 10.3389/fimmu.2019.02571
- Ferreira MC, Dias-Pereira AC, Branco-de-Almeida LS, Martins CC, Paiva SM. Impact of periodontal disease on quality of life: a systematic review. *J Periodont Res.* (2017) 52:651–665. doi: 10.1111/jre.12436
- Sulaiman L, Saub R, Baharuddin NA, Safii SH, Gopal Krishna V, Bartold PM, et al. Impact of severe chronic periodontitis on oral health-related quality of life. *Oral Health Prev Dent.* (2019) 17:365–73. doi: 10.3290/j.ohpd.a42502
- Yiemstan S, Krisdapong, S, Piboonratanakit P. Association between clinical signs of oral lichen planus and oral health-related quality of life: a preliminary study. *Dent J.* (2020) 8:113. doi: 10.3390/dj8040113
- Daume L, Kreis C, Bohner L, Kleinheinz J, Jung S. Does the clinical form of oral lichen planus (OLP) influence the oral health-related quality of life (OHRQoL)? *Int J Environ Res Public Health.* (2020) 17:6633. doi: 10.3390/ijerph17186633
- Yuwanati M, Gondivkar S, Sarode SC, Gadbail A, Sarode GS, Patil S, et al. Impact of oral lichen planus on oral health-related quality of life: a systematic review and meta-analysis. *Clin Pract.* (2021) 11:272–86. doi: 10.3390/clinpract11020040
- Kassebaum NJ, Bernabé E, Dahiya M, Bhandari B, Murray CJL, Marcenes W. Global burden of severe periodontitis in 1990–2010: a systematic review and meta-regression. *J Dent Res.* (2014) 93:1045–53. doi: 10.1177/0022034514552491
- Listl S, Galloway J, Mossey PA, Marcenes W. Global economic impact of dental diseases. *J Dent Res.* (2015) 94:1355–61. doi: 10.1177/0022034515602879
- Tonetti MS, Jepsen S, Jin L, Otomo-Corgel J. Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: a call for global action. *J Clin Periodontol.* (2017) 44:456–462. doi: 10.1111/jcpe.12732
- Chen MX, Zhong YJ, Dong QQ, Wong HM, Wen YF. Global, regional, and national burden of severe periodontitis, 1990–2019: an analysis of the Global Burden of Disease Study 2019. *J Clin Periodontol.* (2021) 48:1165–88. doi: 10.1111/jcpe.13506
- Sanz M, Herrera D, Kebschull M, Chapple IL, Jepsen S, Berglundh T, et al. Treatment of stage I–III periodontitis—The EFP S3 level clinical practice guideline. *J Clin Periodontol.* (2020) 47:4–60. doi: 10.1111/jcpe.13290
- Ramseier CA, Woelber JP, Kitzmann J, Detzen L, Carra MC, Bouchard P. Impact of risk factor control interventions for smoking cessation and promotion of healthy lifestyles in patients with periodontitis: a systematic review. *J Clin Periodontol.* (2020) 47:90–106. doi: 10.1111/jcpe.13240
- Page RC, Kornmann KS. The pathogenesis of human periodontitis: an introduction. *Periodontology.* (2000) 14:9–11. doi: 10.1111/j.1600-0757.1997.tb00189.x

19. Huang S, Yang F, Zeng X, Chen J, Li R, Wen T, et al. Preliminary characterization of the oral microbiota of Chinese adults with and without gingivitis. *BMC Oral Health*. (2011) 11:33. doi: 10.1186/1472-6831-11-33
20. Klukowska M, Goyal CR, Khambe D, Cannon M, Miner M, Gurich N, et al. Response of chronic gingivitis to hygiene therapy and experimental gingivitis. Clinical, microbiological and metabonomic changes. *Am J Dent*. (2015) 28:273–84.
21. Al-Kamel A, Baraniya D, Abdunaser Al-Hajj W, Halboub E, Abdurab S, Chen T, et al. Subgingival microbiome of experimental gingivitis shifts associated with the use of chlorhexidine and N-acetyl cysteine mouthwashes. *J Oral Microbiol*. (2019) 11:1608141. doi: 10.1080/20002297.2019.1608141
22. Schätzle M, Løe H, Bürgin W, Ånerud Å, Boysen H, Lang NP. Clinical course of chronic periodontitis. *J Clin Periodontol*. (2003) 30:887–901. doi: 10.1034/j.1600-051X.2003.00414.x
23. Chapple ILC, Van der Weijden D, Doerfer C, Herrera D, Shapira L, Polak D, et al. Primary prevention of periodontitis: managing gingivitis. *J Clin Periodontol*. (2015) 42:S71–6. doi: 10.1111/jcpe.12366
24. Van der Weijden FA, Slot DE. Efficacy of homecare regimens for mechanical plaque removal in managing gingivitis - a meta review. *J Clin Periodontol*. (2015) 42:S77–S91. doi: 10.1111/jcpe.12359
25. Buduneli N. Environmental factors and periodontal microbiome. *Periodontol*. (2020) 85:112–25. doi: 10.1111/prd.12355
26. Nibali L, Koidou VP, Nieri M, Barbato L, Pagliaro U, Cairo F. Regenerative surgery versus access flap for the treatment of intra-bony periodontal defects: a systematic review and meta-analysis. *J Clin Periodontol*. (2020) 47:320–51. doi: 10.1111/jcpe.13237
27. Løe H, Ånerud Å, Boysen H, Smith M. The natural history of periodontal disease in man. The rate of periodontal destruction before 40 years of age. *J Periodontol*. (1978) 49:607–20. doi: 10.1902/jop.1978.49.12.607
28. Løe H, Ånerud A, Boysen H, Morrison E. Natural history of periodontal disease in man. Rapid, moderate and no loss of attachment in Sri Lankan laborers 14 to 46 years of age. *J Clin Periodontol*. (1986) 13:431–445.
29. Heitz-Mayfield LJA, Schätzle M, Løe H, Bürgin W, Ånerud Å, Boysen H, et al. Clinical course of chronic periodontitis. *J Clin Periodontol*. (2003) 30:902–8. doi: 10.1034/j.1600-051X.2003.00399.x
30. Graetz C, Dorfer CE, Kahl M, Kocher T, Fawzy El-Sayed K, Wiebe J-F, et al. Retention of questionable and hopeless teeth in compliant patients treated for aggressive periodontitis. *J Clin Periodontol*. (2011) 38:707–14. doi: 10.1111/j.1600-051X.2011.01743.x
31. Salvi GE, Mischler DC, Schmidlin K, Matuliene G, Pjetursson BE, Bragger U, et al. Risk factors associated with the longevity of multi-rooted teeth. Long-term outcomes after active and supportive periodontal therapy. *J Clin Periodontol*. (2014) 41:701–7. doi: 10.1111/jcpe.12266
32. Ramseier CA, Nydegger M, Walter C, Fischer G, Sculean A, Lang NP, et al. Time between recall visits and residual probing depths predict long-term stability in patients enrolled in supportive periodontal therapy. *J Clin Periodontol*. (2019) 46:218–30. doi: 10.1111/jcpe.13041
33. Slot DE, Valkenburg C, Van der Weijden GA. Mechanical plaque removal of periodontal maintenance patients: A systematic review and network meta-analysis. *J Clin Periodontol*. (2020) 47:107–24. doi: 10.1111/jcpe.13275
34. Hirschfeld LI, Wasserman B. A long-term survey of tooth loss in 600 treated periodontal patients. *J Periodontol*. (1978) 49:225237.
35. Chace R, Low SB. Survival characteristics of periodontally-involved teeth: a 40-year study. *J Periodontol*. (1993) 64:701–705.
36. Fardal Ø, Johannessen AC, Linden GJ. Tooth loss during maintenance following periodontal treatment in a periodontal practice in Norway. *J Clin Periodontol*. (2004) 31:550–5. doi: 10.1111/j.1600-051X.2004.00519.x
37. Müller Campanile V, Megally A, Campanile G, Gayet-Ageron A, Giannopoulou C, Mombelli A. Risk factors for recurrence of periodontal disease in patients in maintenance care in a private practice. *J Clin Periodontol*. (2019) 46:918–26. doi: 10.1111/jcpe.13165
38. Novaes ABR, Palioto DB, de Andrade PF, Marchesan JT. Regeneration of class II furcation defects: determinants of increased success. *Braz Dent J*. (2005) 16:87–97. doi: 10.1590/S0103-64402005000200001
39. Artzi Z, Sudri S. The impact of supportive periodontal treatment on the outcome of regenerative surgical therapy in aggressive periodontitis patients. *Quintessence Int*. (2021) 52:402–10. doi: 10.3290/j.qi.b912657
40. Sciubba JJ. Autoimmune oral mucosal diseases: clinical, etiologic, diagnostic, and treatment considerations. *Dent Clin North Am*. (2011) 55:89–103. doi: 10.1016/j.cden.2010.08.008
41. Yang M, Wu H, Zhao M, Chang C, Lu Q. The pathogenesis of bullous skin diseases. *Journal of Translational Autoimmunity*. (2019) 2:100014. doi: 10.1016/j.jtauto.2019.100014
42. Carrozzo M, Porter S, Mercadante V, Fedele S. Oral lichen planus: a disease or a spectrum of tissue reactions? Types, causes, diagnostic algorithms, prognosis, management strategies. *Periodontology*. (2019) 80:105–25. doi: 10.1111/prd.12260
43. Buonavoglia A, Leone P, Dammacco R, Di Lernia G, Petrucci M, Bonamonte D, et al. Pemphigus and mucous membrane pemphigoid: an update from diagnosis to therapy. *Autoimmun Rev*. (2019) 18:349–58. doi: 10.1016/j.autrev.2019.02.005
44. Akman A, Kacaroglu H, Yilmaz E, Alpsoy E. Periodontal status in patients with pemphigus vulgaris. *Oral Dis*. (2008) 14:640–3. doi: 10.1111/j.1601-0825.2008.01445.x
45. Anovazzi G, Kim YJ, Viana AC, Curtis KM, Orrico SR, Cirelli JA, et al. Polymorphisms and haplotypes in the interleukin-4 gene are associated with chronic periodontitis in a Brazilian population. *J Periodontol*. (2010) 81:392–402. doi: 10.1902/jop.2009.090392
46. Karimbux NY, Saraiya VM, Elangovan S, Allareddy V, Kinnunen T, Kornman KS, et al. Interleukin-1 gene polymorphisms and chronic periodontitis in adult whites: a systematic review and meta-analysis. *J Periodontol*. (2012) 83:1407–19. doi: 10.1902/jop.2012.110655
47. Chen X, Huang J, Zhong L, Ding C. Quantitative assessment of the associations between interleukin-8 polymorphisms and periodontitis susceptibility. *J Periodontol*. (2015) 86:292–300. doi: 10.1902/jop.2014.140450
48. Schäfer A. Genetics of periodontitis: discovery, biology, and clinical impact. *Periodontol*. (2018) 78:162–173. doi: 10.1111/prd.12232
49. Kurushima Y, Tsai PC, Castillo-Fernandez J, Couto Alves A, El-Sayed Moustafa JS, Le Roy C, et al. Epigenetic findings in periodontitis in UK twins: a cross-sectional study. *Clin Epigenet*. (2019) 11:27. doi: 10.1186/s13148-019-0614-4
50. Hashim A, Alsam A, Payne MA, Aduse-Opoku J, Curtis MA, Joseph S. Loss of neutrophil homing to the periodontal tissues modulates the composition and disease potential of the oral microbiota. *Infect Immun*. (2021) 89:e0030921. doi: 10.1128/IAI.00309-21
51. Haubek D, Ennibi O-K, Poulsen K, Benzarti N, Baelum V. The highly leukotoxic JP2 clone of actinobacillus actinomycetemcomitans and progression of periodontal attachment loss. *J Dent Res*. (2004) 83:767–70. doi: 10.1177/154405910408301006
52. Haffajee AD, Cugini MA, Tanner A, Pollack RP, Smith C, Kent Jr RL, et al. Subgingival microbiota in healthy, well-maintained elder and periodontitis subjects. *J Clin Periodontol*. (1998) 25:346–353. doi: 10.1111/j.1600-051X.1998.tb02454.x
53. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL. Microbial complexes in subgingival plaque. *J Clin Periodontol*. (1998) 25:134–44. doi: 10.1111/j.1600-051X.1998.tb02419.x
54. Jung WR, Joo JY, Lee JY, Kim HJ. Prevalence and abundance of 9 periodontal pathogens in the saliva of periodontally healthy adults and patients undergoing supportive periodontal therapy. *J Periodontal Implant Sci*. (2021) 51:316–28. doi: 10.5051/jpis.2006640332
55. Chen C, Hemme C, Beleno J, Zj S, Ning D, Qin Y, et al. Oral microbiota of periodontal health and disease and their changes after nonsurgical periodontal therapy. *ISME J*. (2018) 12:1210–24. doi: 10.1038/s41396-017-0037-1
56. Hajishengallis G, Liang S, Payne MA, Hashim A, Jotwani R, Eskan MA, et al. Low-abundance biofilm species orchestrates inflammatory periodontal disease through the commensal microbiota and complement. *Cell Host Microbe*. (2011) 10:497–506. doi: 10.1016/j.chom.2011.10.006
57. Jiao Y, Hasegawa M, Inohara N. The role of oral pathobionts in dysbiosis during periodontitis development. *J Dent Res*. (2014) 93:539–46. doi: 10.1177/0022034514528212
58. Hajishengallis G. Periodontitis: from microbial immune subversion to systemic inflammation. *Nat Rev Immunol*. (2015) 15:30–44. doi: 10.1038/nri3785

59. Herrero ER, Fernandes S, Verspecht T, Ugarte-Berzal E, Boon N, Proost P, et al. Dysbiotic biofilms deregulate the periodontal inflammatory response. *J Dent Res.* (2018) 97:547–55. doi: 10.1177/0022034517752675
60. Kumar PS. Microbial dysbiosis: The root cause of periodontal disease. *J Periodontol.* (2021) 92:1079–87. doi: 10.1002/JPER.21-0245
61. Hajishengallis G, Lamont RJ. Polymicrobial communities in periodontal disease: their quasi-organismal nature and dialogue with the host. *Periodontol.* (2021) 86:210–30. doi: 10.1111/prd.12371
62. Payne MA, Hashim A, Alsam A, Joseph S, Aduse-Opoku J, Wade WG, et al. Horizontal and vertical transfer of oral microbial dysbiosis and periodontal disease. *J Dent Res.* (2019) 98:1503–1510. doi: 10.1177/0022034519877150
63. Tatakis DN, Trombelli L. Modulation of clinical expression of plaque-induced gingivitis. *J Clin Periodontol.* (2004) 31:229–38. doi: 10.1111/j.1600-051x.2004.00477.x
64. Murakami S, Mealey BL, Mariotti A, Chapple ILC. Dental plaque-induced gingival conditions. *J Clin Periodontol.* (2018) 45:S17–27. doi: 10.1111/jcpe.12937
65. Trombelli L, Farina R, Silva CO, Tatakis DN. Plaque-induced gingivitis: Case definition and diagnostic considerations. *J Clin Periodontol.* (2018) 45:S44–67. doi: 10.1111/jcpe.12939
66. Ng E, Tay JRH, Balan P, Ong MMA, Bostanci N, Belibasakis GN, et al. Metagenomic sequencing provides new insights into the subgingival bacteriome and aetiopathology of periodontitis. *J Periodontol Res.* (2021) 56:205–18. doi: 10.1111/jre.12811
67. Sukumar S, Roberts AP, Martin FE, Adler CJ. Metagenomic insights into transferable antibiotic resistance in oral bacteria. *J Dent Res.* (2016) 95:969–76. doi: 10.1177/0022034516648944
68. Jepsen S, Caton JG, Alnabdar JM, Bissada NF, Bouchard P, Cortellini P, et al. Periodontal manifestations of systemic diseases and developmental and acquired conditions: Consensus report of workgroup 3 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Clin Periodontol.* (2018) 45:S219–29. doi: 10.1111/jcpe.12951
69. Belibasakis GN, Bostanci N, Marsh PD, Zaura E. Applications of the oral microbiome in personalized dentistry. *Arch Oral Biol.* (2019) 104:7–12. doi: 10.1016/j.archoralbio.2019.05.023
70. Suomi JD, Greene JC, Vermillion JR, Doyle J, Change JJ, Leatherwood EC. The effect of controlled oral hygiene procedures on the progression of periodontal disease in adults: results after third and final year. *J Periodontol.* (1971) 42:152–60. doi: 10.1902/jop.1971.42.3.152
71. Magnusson I, Lindhe J, Yoneyama T, Liljenberg B. Recolonization of a subgingival microbiota following scaling in deep pockets. *J Clin Periodontol.* (1984) 11:193–207. doi: 10.1111/j.1600-051X.1984.tb01323.x
72. Sälzer S, Graetz C, Dörfer CE, Slot DE, Van der Weijden FA. Contemporary practices for mechanical oral hygiene to prevent periodontal disease. *Periodontol.* (2020) 84:35–44. doi: 10.1111/prd.12332
73. Suvan J, Leira Y, Moreno F, Graziani F, Derks J, Tomasi C. Subgingival instrumentation for treatment of periodontitis. A systematic review. *J Clin Periodontol.* (2020) 47:155–75. doi: 10.1111/jcpe.13245
74. Huang S, Li Z, He T, Bo C, Chang J, Li L, et al. Microbiota-based Signature of Gingivitis Treatments: A Randomized Study. *Sci Rep.* (2016) 20:6. doi: 10.1038/srep24705
75. Karimova M, Moyes D, Ide M, Setterfield JF. The human microbiome in immunobullous disorders and lichen planus. *Clin Exp Dermatol.* (2021) 47:522–8. doi: 10.1111/CED.14987
76. Seckin Ertugrul A, Arslan U, Dursun R, Sezgin Hakki S. Periodontopathogen profile of healthy and oral lichen planus patients with gingivitis or periodontitis. *Int J Oral Sci.* (2013) 5:92–7. doi: 10.1038/ijos.2013.30
77. Liu H, Chen H, Liao Y, Li H, Shi L, Deng Y, et al. Comparative analyses of the subgingival microbiome in chronic periodontitis patients with and without gingival erosive oral lichen planus based on 16S rRNA gene sequencing. *Biomed Res Int.* (2021) 2021:9995225. doi: 10.1155/2021/9995225
78. Künstner A, Sommer A, Künzel S, Zillikens D, Gläser R, Baines J, et al. 45th Annual Meeting of the Arbeitsgemeinschaft Dermatologische Forschung (ADF) Zurich, Switzerland, March 7–10, 2018. *Exp Dermatol.* (2018) 27:e2–e106. doi: 10.1111/exd.13486
79. Huang S, Mao J, Zhou L, Xiong X, Deng Y. The imbalance of gut microbiota and its correlation with plasma inflammatory cytokines in pemphigus vulgaris patients. *Scand J Immunol.* (2019) 90. doi: 10.1111/sji.12799
80. Ellebrecht CT, Srinivas G, Bieber K, Banczyk D, Kalies K, Künzel S, et al. Skin microbiota-associated inflammation precedes autoantibody induced tissue damage in experimental epidermolysis bullosa acquisita. *J Autoimmun.* (2016) 68:14–22. doi: 10.1016/j.jaut.2015.08.007
81. Ramón-Fluixá C, Bagán-Sebastián J, Milián-Masanet M, Scully C. Periodontal status in patients with oral lichen planus: a study of 90 cases. *Oral Dis.* (1999) 5:303–306. doi: 10.1111/j.1601-0825.1999.tb00094.x
82. Tricamo MB, Rees TD, Hallmon WW, Wright JM, Cueva MA, Plemons JM. Periodontal status in patients with gingival mucous membrane pemphigoid. *J Periodontol.* (2006) 77:398–405. doi: 10.1902/jop.2006.050113
83. Azizi A, Rezaee M. Comparison of Periodontal Status in Gingival Oral Lichen Planus Patients and Healthy Subjects. *Dermatol Res Pract.* (2012) 2012:561232. doi: 10.1155/2012/561232
84. Jascholt I, Lai O, Zillikens D, Kasperkiewiczabella M. Periodontitis in oral pemphigus and pemphigoid: a systematic review of published studies. *J Am Acad Dermatol.* (2017) 76:975–8. doi: 10.1016/j.jaad.2016.10.028
85. Wang H, Luo Z, Lei L, Sun Z, Zhou M, Dan H, et al. Interaction between oral lichen planus and chronic periodontitis with TH17-associated cytokines in serum. *Inflammation.* (2013) 36:696–704. doi: 10.1007/s10753-013-9594-2
86. Erpenstein H. Periodontal and prosthetic treatment in patients with oral lichen planus. *J Clin Periodontol.* (1985) 12:1041112. doi: 10.1111/j.1600-051X.1985.tb01369.x
87. Choi YS, Kim Y, Yoon HJ, Baek KJ, Alam J, Park HK, et al. The presence of bacteria within tissue provides insights into the pathogenesis of oral lichen planus. *Sci Rep.* (2016) 6:29186. doi: 10.1038/srep29186
88. Holmstrup P, Schiøtz AW, Westergaard J. Effect of dental plaque control on gingival lichen planus. *Oral Surg Oral Med Oral Pathol.* (1990) 69:585–90. doi: 10.1016/0030-4220(90)90241-J
89. Lo Russo L, Gallo C, Pellegrino G, Lo Muzio L, Pizzo G, Campisi G, et al. Periodontal clinical and microbiological data in desquamative gingivitis patients. *Clin Oral Invest.* (2014) 18:917–25. doi: 10.1007/s00784-013-1038-8
90. Figuero E, Roldán S, Serrano J, Escribano M, Martín C, Preshaw PM. Efficacy of adjunctive therapies in patients with gingival inflammation: a systematic review and meta-analysis. *J Clin Periodontol.* (2020) 47:125–43. doi: 10.1111/jcpe.13244
91. Herlofson B, Barkvold P. The effect of two toothpaste detergents on the frequency of recurrent aphthous ulcers. *Acta Odontol. Scand.* (1996) 54:150–153. doi: 10.3109/00016359609003515
92. Pérez-López D, Varela-Centelles P, García-Pola MJ, Castelo-Baz P, García-Caballero L, Seoane-Romero JM. Oral mucosal peeling related to dentifrices and mouthwashes: a systematic review. *Med Oral Patol Oral Cir Bucal.* (2019) 24:e452–60. doi: 10.4317/medoral.22939
93. Alli BY, Erinoso OA, Olawuyi AB. Effect of sodium lauryl sulfate on recurrent aphthous stomatitis: A systematic review. *J Oral Pathol Med.* (2019) 48:358–64. doi: 10.1111/jop.12845
94. Tadin A, Gavic L, Govic T, Galic N, Vladislavic NZ, Zeljezic D. *In vivo* evaluation of fluoride and sodium lauryl sulphate in toothpaste on buccal epithelial cells toxicity. *Acta Odontol Scand.* (2019) 77:386–93. doi: 10.1080/00016357.2019.1577988
95. Pretty I, Edgar W, Higham S, A. study to assess the efficacy of a new detergent free, whitening dentifrice in vivo using QLF planimetric analysis. *Br Dent J.* (2004) 197:561–6. doi: 10.1038/sj.bdj.4811809
96. Nadelmann E, Czernik A. Wound care in immunobullous disease. *In: Autoimmune Bullous Diseases.* (2018).
97. Thongprasom K, Dhanuthai K. Steroids in the treatment of lichen planus: a review. *J Oral Sci.* (2008) 50:377–85. doi: 10.2334/josnusd.50.377
98. Heasman PA, Vernazza CR, Gaunt FL, Pennington MW. Cost-effectiveness of adjunctive antimicrobials in the treatment of periodontitis. *Periodontol.* (2011) 55:217–30. doi: 10.1111/j.1600-0757.2010.00341.x
99. Herrera D, Matesanz P, Martín C, Oud V, Feres M, Teughels W. Adjunctive effect of locally delivered antimicrobials in periodontitis therapy: A systematic review and meta-analysis. *J Clin Periodontol.* (2020) 47:239–56. doi: 10.1111/jcpe.13230



100. Teughels W, Feres M, Oud V, Martin C, Matesanz P, Herrera D. Adjunctive effect of systemic antimicrobials in periodontitis therapy: A systematic review and meta-analysis. *J Clin Periodontol.* (2020) 47:257–81. doi: 10.1111/jcpe.13264
101. Donos N, Calciolari E, Brusselsaers N, Goldoni M, Bostanci N, Belibasakis GN. The adjunctive use of host modulators in non-surgical periodontal therapy. A systematic review of randomized, placebo-controlled clinical studies. *J Clin Periodontol.* (2020) 47:199–238. doi: 10.1111/jcpe.13232
102. Sgolastra F, Petrucci A, Ciarrocchi I, Masci C, Spadaro A. Adjunctive systemic antimicrobials in the treatment of chronic periodontitis: A systematic review and network meta-analysis. *J Periodont Res.* (2021) 56:236–48. doi: 10.1111/jre.12821
103. Ide M. Periodontal Diseases. In: Palmer NO, editor. *Antimicrobial Prescribing in Dentistry - Good Practice, Guidelines, and 3rd ed.* London: Faculty of General Dental Practitioners. (2020). p. 49–63.
104. Fauci AS. Corticosteroids in autoimmune disease. *Hosp Pract.* (2016) 18:99–114. doi: 10.1080/21548331.1983.11702661
105. Bootun R. Effects of immunosuppressive therapy on wound healing. *Int Wound J.* (2013) 10:98–104. doi: 10.1111/j.1742-481X.2012.00950.x
106. Piette EW, Werth VP. Dapsone in the management of the autoimmune bullous diseases. *Dermatol Clin.* (2011) 29:561. doi: 10.1016/j.det.2011.06.018
107. Noe MH, Fairley JA. Bullous pemphigoid. In: *Autoimmune Bullous Diseases: Approach and Management.* Springer International Publishing. (2016). pp. 57–73.
108. Fagundes PPS, Santi CG, Maruta CW, Miyamoto D, Aoki V. Autoimmune bullous diseases in pregnancy: clinical and epidemiological characteristics and therapeutic approach. *An Bras Dermatol.* (2021) 96:581–90.
109. Williams HC, Wojnarowska F, Kirtschig G, Mason J, Godec TR, Schmidt E, et al. Doxycycline versus prednisolone as an initial treatment strategy for bullous pemphigoid: a pragmatic, non-inferiority, randomised controlled trial. *Lancet.* (2017) 389:1630–8. doi: 10.1016/S0140-6736(17)30560-3
110. Gabler WL, Creamer HR. Suppression of human neutrophil functions by tetracyclines. *J Periodontol Res.* (1991) 26:52–8. doi: 10.1111/j.1600-0765.1991.tb01626.x
111. Mensing H, Krause S. Therapy of bullous pemphigoid with erythromycin. *Med Klin.* (1990) 85:481–4.
112. Altomare G, Capella GL, Fracchiolla C, Frigerio E. Treatment of bullous pemphigoid with erythromycin: a reappraisal. *Eur J Dermatol.* (1999) 9:583–5.
113. Cooper SM, Powell J, Wojnarowska F. Linear IgA disease: successful treatment with erythromycin. *Clin Exp Dermatol.* (2002) 27:677–9. doi: 10.1046/j.1365-2230.2002.01133.x
114. Vangoitsenhoven R, Cresci GAM. Role of microbiome and antibiotics in autoimmune diseases. *Nutr Clin Pract.* (2020) 35:406–16. doi: 10.1002/ncp.10489
115. Feliciani C, Toto P, Pour SM, Coscione G, Amerio P, Amerio P, et al. In vitro and in vivo expression of interleukin-1 $\alpha$  and tumor necrosis factor- $\alpha$  mRNA in pemphigus vulgaris: interleukin-1 $\alpha$  and tumor necrosis factor- $\alpha$  are involved in acantholysis. *J Invest Dermatol.* (2000) 114:71–7. doi: 10.1046/j.1523-1747.2000.00835.x
116. Ruocco V, De Angelis E, Lombardi ML. Drug-induced pemphigus. II Pathomechanisms and experimental investigations. *Clin Dermatol.* (1993) 11:507–13. doi: 10.1016/0738-081X(93)90158-9
117. Markitziu A, Zafropoulos G, de Jacoby LF, Pisanty S. Periodontal alterations in patients with pemphigus vulgaris taking steroids. *J Clin Periodontol.* (1990) 17:228–32. doi: 10.1111/j.1600-051X.1990.tb00018.x
118. López-Jornet P, Camacho-Alonso F. Application of a motivation-behavioral skills protocol in gingival lichen planus: a short-term study. *J Periodontol.* (2010) 81:1449–54. doi: 10.1902/jop.2010.100245
119. Tomasi C, Leyland AH, Wennstrom JL. Factors influencing the outcome of non-surgical periodontal treatment: a multilevel approach. *J Clin Periodontol.* (2007) 34:682–90. doi: 10.1111/j.1600-051X.2007.01111.x
120. Sbordone L, Ramaglia L, Gulletta E, Iacono V. Recolonization of the subgingival microflora after scaling and root planing in human periodontitis. *J Periodontol.* (1990) 61:579–584. doi: 10.1902/jop.1990.61.9.579
121. Arduino PG, Lopetuso E, Carcieri P, Giacometti S, Carbone M, Tanteri C, et al. Professional oral hygiene treatment and detailed oral hygiene instructions in patients affected by mucous membrane pemphigoid with specific gingival localization: a pilot study in 12 patients. *Int J Dent Hygiene.* (2012) 10:138–41. doi: 10.1111/j.1601-5037.2011.00527.x
122. Gambino A, Carbone M, Arduino PG, Carcieri P, Carbone L, Broccoletti R. Conservative approach in patients with pemphigus gingival vulgaris: a pilot study of five cases international. *J Dent.* (2014) 2014:747506. doi: 10.1155/2014/747506
123. Stone SJ, Heasman PA, Staines KS, McCracken GI. The impact of structured plaque control for patients with gingival manifestations of oral lichen planus: a randomized controlled study. *J Clin Periodontol.* (2015) 42:356–62. doi: 10.1111/jcpe.12385
124. Garcia-Pola MJ, Rodriguez-López S, Fernández-Vigil A, Bagán L, Garcia-Martín JM. Oral hygiene instructions and professional control as part of the treatment of desquamative gingivitis. *System Rev Med Oral Patol Oral Cir Bucal.* (2019) 24:e136–44. doi: 10.4317/medoral.22782
125. Bianco L, Romano F, Maggiora M, Bongiovanni L, Guzzi N, Curmei E, et al. Effect of sonic versus manual supervised toothbrushing on both clinical and biochemical profiles of patients with desquamative gingivitis associated with oral lichen planus: A randomized controlled trial. *Int J Dent Hygiene.* (2019) 17:161–9. doi: 10.1111/idh.12377
126. Trombelli L, Simonelli A, Franceschetti G, Maietti E, Farina R. What periodontal recall interval is supported by evidence? *Periodontol.* (2020) 84:124–33. doi: 10.1111/prd.12340
127. Albander JM, Susin C, Hughes FJ. Manifestations of systemic diseases and conditions that affect the periodontal attachment apparatus: Case definitions and diagnostic considerations. *J Clin Periodontol.* (2018) 45:S171–89. doi: 10.1111/jcpe.12947
128. Papananou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, et al. Periodontitis: consensus report of workgroup 2 of the 2017 world workshop on the classification of periodontal and peri-implant diseases and conditions. *J Clin Periodontol.* (2018) 45:S162–70. doi: 10.1111/jcpe.12946

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Ide, Karimova and Setterfield. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.