



King's Research Portal

DOI:
[10.1093/ckj/sfac080](https://doi.org/10.1093/ckj/sfac080)

Document Version
Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):
Chilcot, J., Picariello, F., & Farrington, K. (2022). Depression and clinical outcomes in CKD: do anti-depressants play a role? (EQUAL STUDY). *Clinical Kidney Journal*, 15(8), 1628–1629. <https://doi.org/10.1093/ckj/sfac080>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

LETTER TO THE EDITOR

Depression and clinical outcomes in CKD: do anti-depressants play a role? (EQUAL study)

Joseph Chilcot ¹, Federica Picariello ¹ and Ken Farrington^{2,3}

¹Psychology Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK, ²Renal Unit, Lister Hospital, Stevenage, UK and ³Centre for Health Services and Clinical Research, University of Hertfordshire, Hatfield, UK

Correspondence to: Joseph Chilcot; E-mail: joseph.chilcot@kcl.ac.uk

We read with interest the impressive study 'Associations between depressive symptoms and disease progression in older patients with chronic kidney disease: results of the EQUAL study' by Eveleens Maarse [1]. This study examined the associations between depression symptoms and clinical outcomes in older people (≥ 65 years of age) with chronic kidney disease (CKD) (eGFR ≤ 20 mL/min/1.73 m²) and considered sex differences. The study found no association between baseline depressive symptoms and a decrease in eGFR. Furthermore, there was no association between depressive symptoms and time to dialysis initiation or a combined outcome (dialysis initiation and all-cause mortality). However, in men, there was a significant association between the presence of depressive symptoms [as defined by a score of ≤ 70 on the Mental Health Inventory (MHI-5)] and all-cause mortality compared with those with no symptoms [adjusted hazard ratio [aHR] 1.41 [95% confidence interval (CI) 1.03–1.93]]. While the study has several strengths, including the large sample size, comprehensive data defining the sample and rigorous approach to data analysis, we would like to comment on two aspects of the study: measurement of the depressive symptoms and the role of antidepressants.

First, it is important to note that the MHI-5 was used to measure depressive symptoms. While the authors recognize that the use of other validated depression tools may have improved the robustness of the findings, the MHI-5 is best conceptualized as a measure of psychological distress, particularly as two of the items relate to anxiety. This is particularly problematic when considered together with how item-level missingness was addressed ('the mental health score was calculated when minimal three of five questions were answered'). In addition, there are unique features of depression in older age that have measure-

ment implications. Therefore it is likely that the MHI-5 is not the optimal measure given the focus and sample here.

Second, in the discussion, we were interested to read the suggestion that in men, antidepressants may improve survival given the association between depressive symptoms and mortality ('might it be possible that at least in men, antidepressant treatment might not only improve HRQOL, but could potentially improve survival'). We are not convinced by this interpretation, particularly as evidence for the efficacy and safety of antidepressants in people with kidney disease is inconclusive [2–4]. An additional point in relation to the interpretation offered for the association between depression and mortality in men may be reduced help-seeking among men. It would be interesting to know whether the rates of past depression differed by sex (although we acknowledge numbers are relatively small).

We note that antidepressant data were available in 1032 patients (77.9%). While the analyses are suitably adjusted for confounders, we are intrigued by the lack of adjustment for the use of antidepressants. While we recognize the authors' comment regarding the challenges surrounding the number of confounders in survival models and the issues of over-adjustment, it would be interesting to know the findings of an exploratory set of subanalyses that adjust for antidepressant use. Furthermore, it would be important to know the crude and adjusted effect of antidepressant use on all-cause mortality.

Our reason for this interest stems from the retrospective analysis (albeit in haemodialysis patients and not CKD), which found that patients who initiated selective serotonin reuptake inhibitors (SSRIs) with higher QT-prolonging potential (citalopram, escitalopram) had a significantly increased risk of

Received: 22.2.2022; Editorial decision: 14.3.2022

© The Author(s) 2022. Published by Oxford University Press on behalf of the ERA. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

sudden cardiac death [aHR 1.18 (95% CI 1.05–1.31)] compared with patients who initiated SSRIs with lower QT-prolonging potential (fluoxetine, fluvoxamine, paroxetine and sertraline) [4]. We have previously raised the possibility that in some kidney patients, antidepressants may be doing more harm than good and suggest that trials evaluating the selective withdrawal of antidepressants in suitable patients may be appropriate [5].

CONFLICT OF INTEREST STATEMENT

None declared.

REFERENCES

1. Eveleens Maarse BC, Chesnaye NC, Schouten R *et al*. Associations between depressive symptoms and disease progression in older patients with chronic kidney disease: results of the EQUAL study. *Clin Kidney J* 2021; **15**: 786–797
2. Hedayati SS, Gregg LP, Carmody T *et al*. Effect of sertraline on depressive symptoms in patients with chronic kidney disease without dialysis dependence: the CAST randomized clinical trial. *JAMA* 2017; **318**: 1876–1890
3. Palmer SC, Natale P, Ruospo M *et al*. Antidepressants for treating depression in adults with end-stage kidney disease treated with dialysis. *Cochrane Database Syst Rev* 2016; **2016**: CD004541
4. Assimon MM, Brookhart, MA, Flythe JE. Comparative cardiac safety of selective serotonin reuptake inhibitors among individuals receiving maintenance hemodialysis. *J Am Soc Nephrol* 2019; **30**: 611–623
5. Chilcot J, Farrington K. The case for selective withdrawal of antidepressants in patients with advanced kidney disease. *J Am Soc Nephrol* 2019; **30**: 1339–1340