



King's Research Portal

DOI:

[10.1016/j.jss.2022.05.009](https://doi.org/10.1016/j.jss.2022.05.009)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Haiser, A., Aydin, A., Kunduzi, B., Ahmed, K., & Dasgupta, P. (2022). A Systematic Review of Simulation-Based Training in Vascular Surgery. *Journal of Surgical Research*, 279, 409-419.
<https://doi.org/10.1016/j.jss.2022.05.009>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.JournalofSurgicalResearch.com

Research review

A Systematic Review of Simulation-Based Training in Vascular Surgery



Alexander Haiser, BSc (Hons),^a Abdullatif Aydin, MBBS, PhD,^{b,*}
Basir Kunduzi, MBBS,^c Kamran Ahmed, PhD, FRCS,^b
and Prokar Dasgupta, MSc, MD, FRCS^b

^a Guy's, King's and St Thomas' School of Medical Education, King's College London, London, UK

^b MRC Centre for Transplantation, Guy's Hospital, King's College London, London, UK

^c Department of Transplant Surgery, Guy's and St. Thomas' NHS Foundation Trust, London, London, UK

ARTICLE INFO

Article history:

Received 10 January 2022

Received in revised form

20 April 2022

Accepted 22 May 2022

Available online xxx

Keywords:

Endovascular surgery

Open vascular surgery

Simulation

Training

Vascular skills

ABSTRACT

Introduction: Recent advancements in surgical technology, reduced working hours, and training opportunities exacerbated by the COVID-19 pandemic have led to an increase in simulation-based training. Furthermore, a rise in endovascular procedures has led to a requirement for high-fidelity simulators that offer comprehensive feedback. This review aims to identify vascular surgery simulation models and assess their validity and levels of effectiveness (LoE) for each model in order to successfully implement them into current training curricula.

Methods: PubMed and EMBASE were searched on January 1, 2021, for full-text English studies on vascular surgery simulators. Eligible articles were given validity ratings based on Messick's modern concept of validity alongside an LoE score according to McGaghie's translational outcomes.

Results: Overall 76 eligible articles validated 34 vascular surgery simulators and training courses for open and endovascular procedures. High validity ratings were achieved across studies for: content (35), response processes (12), the internal structure (5), relations to other variables (57), and consequences (2). Only seven studies achieved an LoE greater than 3/5. Overall, ANGIO Mentor was the most highly validated and effective simulator and was the only simulator to achieve an LoE of 5/5.

Conclusions: Simulation-based training in vascular surgery is a continuously developing field with exciting future prospects, demonstrated by the vast number of models and training courses. To effectively integrate simulation models into current vascular surgery curricula and assessments, there is a need for studies to look at trainee skill retention over a longer period of time. A more detailed discussion on cost-effectiveness is also needed.

© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author. MRC Centre for Transplantation, Guy's Hospital, King's College London, 5th Floor Southwark Wing, London SE1 9RT, UK. Tel.: +44 (0)20 7188 8580; fax: +44 (0)20 3312 6787.

E-mail address: abdullatif.aydin@kcl.ac.uk (A. Aydin).

0022-4804/\$ – see front matter © 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.1016/j.jss.2022.05.009>

Introduction

Influenced by the aviation and military sectors, simulation has become more apparent in medical education and has recently grown to be incorporated in the training curricula of various specialties.¹ Vascular surgical training has historically followed the Halstedian model of 'learning by doing'. This model is often criticized because current surgical trainees are limited by a lack of opportunity of 'what comes through the door'.² Working time restrictions have reduced surgical training in the United Kingdom by a third. Factors, including the shortening of residency time and reduced training opportunities, worsened by the COVID-19 pandemic, are having a large impact on vascular surgical training.²⁻⁴ Additionally, vascular surgery has evolved into a more independent and highly complex specialty. This is due to increasingly more endovascular procedures being conducted, with over half of aortic aneurysms in Europe now being treated by endovascular repair.⁵ These relatively new treatment methods require additional skills to be acquired. Furthermore, some studies have shown that vascular trainees experience fewer teaching opportunities for endovascular procedures than for open surgery, leading to a requirement for alternative teaching methods.^{6,7} It is still of vital importance to ensure that vascular trainees have proficiency in both open and minimally invasive procedures.^{8,9}

In recent years, there has been a greater emphasis on simulation as a surgical training tool to achieve the same level of competency and maintain patient safety.¹⁰ Due to the rise in endovascular procedures, which are known to offer improved patient outcomes and reduced hospital stay compared to open vascular procedures,^{5,11} there will be a subsequent rise in high-fidelity simulators utilized for training. Despite this, there is possibly an even greater demand for open simulators due to the requirement for junior trainees to safely practice common open procedures such as arterial anastomosis and additionally for senior trainees to undertake crisis management in a simulated operating theater.¹² These may actually be more expensive and face greater resistance than their endovascular counterparts, due to the challenge in replicating a three-dimensional field as opposed to a two-dimensional computer system featured in many endovascular simulators.¹² In order to ensure that vascular trainees can perform to a high standard, regulatory bodies and a European wide consensus have recognized the importance of simulation as part of the curriculum and as a method of assessment, which has been shown in other specialties to be an effective way of ensuring proficiency in surgical skills.^{2,10,13,14}

Based on what is currently known, the rationale for this review is to identify the current simulation modalities, assess their validity, and determine how effective they are for training vascular surgeons.

Methods

This review followed the PRISMA guidelines (Fig).¹⁵

Information sources and search

A preliminary search included the terms 'vascular surgery AND simulation' in order to identify MeSH terms. A systematic search of PubMed and Embase was carried out on January 1, 2021, with no limits using the search terms: 'Simulat* AND vascular surg* AND (EVAR OR aneurysm repair OR endovascular OR bypass OR endarterectomy OR angioplasty OR embolectomy) AND (education OR performance OR curriculum OR competence)'. Reference lists from eligible articles were manually searched for potential studies.

Study eligibility criteria

Original articles in full English text were included, with duplicates and conference abstracts excluded. Randomized control trials, cohort, case-control, and cross-sectional studies were included. Population inclusion criteria featured vascular surgeons or trainees at any stage, medical students, or doctors on a vascular placement. Models identified were separated into categories outlining the modality: virtual reality, bench, cadaver, animal, e-learning, computer, or application software.

Data extraction

Articles were screened and organized using the COVIDENCE software. Initially, this was done by title and abstract screening. Once duplicates and noneligible articles were excluded, the remaining articles were read in full. An Excel spreadsheet featured the name of the simulator, manufacturer, fidelity, availability, type of model, reference, number of participants, training level, procedure, validity, and effectiveness criteria.

Data analysis

Validity studies were assessed using Messick's modern concept of validity framework, which has been widely utilized in the literature.¹⁶⁻²⁶ This evaluated the strength of each source of validity evidence using five parameters (Table 1), quantified by Beckman *et al.*'s rating scale, which indicated the strength of validity of each parameter using a scale from 0 to 2 (Table 2).²⁷ Potential author bias and selective outcome reporting was assessed on a study level.

An adaption of McGaghie's translational outcomes was applied, and a level of effectiveness score (LoE) from 1 to 5 was given for each simulation model,²⁸ representing the translational outcome gained from using a simulator (Table 3).

Data analysis was discussed and practiced in detail by authors AH and AA prior to the formal analysis of validity studies. Data analysis was carried out by one author, AH, who applied ratings and scores to individual papers alongside a description of why individual scores and ratings were given. An additional author, AA, was brought in when there was a discrepancy regarding validity ratings or translational outcome scoring.




Fig. – Study selection process according to the PRISMA statement.¹⁵ Figure 1 shows that 525 articles were found after the initial search and exclusion of duplicates. After abstract and title screening, 178 articles were assessed for eligibility via full-text screening and 102 articles were excluded for various reasons. Overall, 76 eligible articles were found for 34 different simulators (See [Supplementary Table I](#)). Simulators were allocated to open vascular surgery, endovascular surgery, nontechnical skills, and simulation courses.

Results

Open vascular surgery

Abdominal aortic aneurysm (AAA) repair

Two simulators for AAA repair had validation criteria or showed an LoE in two articles. The Inanimate OAR bench simulator achieved excellent ratings across the board, rating

2/2 for content, response processes, internal structure, and consequences.²⁹ An LoE of 2/5 was awarded due to a decrease in the simulated procedure time and supervisor interference.

The Open Abdominal Aneurysm Simulation Model earned a rating of 2/2 for content, due to evaluation and refinement by experts and for relations to other variables. Similarly, an LoE of 2/5 was awarded due to an improvement in postsimulator performance.³⁰

Table 1 – Evaluation of validity definitions and examples based on Messick's framework.^{16,17}

Parameter	Definition	Example
Content validity	Test items are relevant and representative of the intended construct	Using expert opinions to ensure all domains are accurately represented
Response process	Thought processes and actions of the subjects and observers are made in accordance with the intended construct	Quality control of assessments, such as standardizing test administration, minimizing examiner bias, and providing a specific test for the task
Internal structure	Test scores across tasks can be reliably reproduced	Calculating interitem reliability and test-retest reliability
Relations to other variables	Test scores correlate with external independent measures that share a theoretic relationship	Comparing scores between groups with different levels of experience in the tested skill
Consequences	The impact of using the assessment	Determining the pass-fail score and considerations for the subject of obtaining a pass or fail, promotion, or privilege

Carotid endarterectomy (CEA)

Three validation studies were identified for three CEA simulators (See [Supplementary Table II](#)). The Carotid Bench model showed content validity with a rating of 2.³¹ An LoE of 2/5 was achieved as there was a significant improvement in surgical skills.

The low-cost Pulsatile Carotid Endarterectomy model by Fletcher et al.³² was tested upon 17 participants, achieving a content rating of 2/2. An increase in confidence and skills scored an LoE = 2/5.

Vascular anastomosis (VA)

VA was the most common type of simulation for open vascular surgery. Eight validation studies for eight simulator models were identified ([Supplementary Table II](#)), and all demonstrated high validity ratings alongside an LoE of 2/5 and above.

The Virtual Surgery simulator³³ achieved a rating of 2/2 for content and relations to other variables by comparing anastomosis performance between vascular surgeons and medical students. An improvement in anastomosis skills led to an LoE of 2/5.

A Frozen Porcine Aorta Anastomosis model discriminated between various levels of experience of residents within the group, thereby achieving a rating = 2/2 for relations to other variables.³⁴ Improvement in the post-test performance led to an LoE of 2/5.

The Vascular Anastomosis model scored highly for content and relations to other variables (rating = 2/2).³⁵ Performance in models was predictive of operative competency on real patients and was associated with less leakage and shorter operating time (LoE = 3/5).

Endovascular surgery

Angioplasty and stenting

These were the most commonly validated vascular procedures, with thirty-four validation studies found for five simulators.

ANGIO Mentor, a high-fidelity VR simulator, was validated by twelve studies as seen in [Supplementary Table III](#).³⁶⁻⁴⁷ Willaert et al. scored 2/2 for content by implementing an expert validated assessment scale on 20 endovascular experts and outlined a clear method of test design and standardization, scoring highly for response process (rating = 2/2).⁴⁰ Maertens et al. demonstrated in two studies that ANGIO Mentor could differentiate between endovascular performance in 32 surgical trainees and additionally 20 vascular surgeons and 29 medical students.^{36,37} All studies received favorable feedback and an improvement in technical metrics, such as decreased procedural and fluoroscopy time (LoE = 1-2/5). Wooster et al. scored an LoE = 3/5, as performance on patients post ANGIO Mentor use was associated with lower doses of radiation, contrast, and shorter procedure time.⁴⁶ Maertens et al. was awarded an LoE of 5/5 as proficiency levels were retained up to 3 mo, alongside an improvement in patient outcomes and better operative performance.³⁷

The Vascular Intervention System Trainer (VIST) was validated by 15 studies⁴⁸⁻⁶² (See [Supplementary Table III](#)). Bech et al. used a previously validated assessment method by experts, and Boyle created expert reviewed scoring sheets (content rating = 2/2).^{52,54,58} Response process ratings were mixed (N-2/2), with some studies featuring an explicit thought process of test design and error reduction methods such as randomization and rater blinding. Rolls et al. demonstrated

Table 2 – Beckman et al.'s rating scale.²⁷

Beckman's rating	Definition
N	No discussion of source of validity evidence
0	Discussion of the source of validity but no data presented
1	Data weakly supports the source of validity or is limited
2	Data strongly supports the source of validity

Table 3 – An adaption of McGaghie’s translational outcomes.²⁸

Parameter	Definition	Example	Level of effectiveness (LoE) score
Internal acceptability	The trainee’s satisfaction with using the simulator	Favorable responses from feedback forms or post-training survey questionnaires	1
Contained effects	Changes in performance in the simulation context	Development of knowledge and/or skills as measured by the simulator tool	2
Downstream effects	Behavioral changes in the clinical context	Adopting safer patient-care practices	3
Target effects	Direct changes to patient outcomes	Reduced rates of surgical complications	4
Collateral effects	Changes on a wider, systemic level	Cost saving; skill retention	5

excellent internal structure by calculating interitem and test-retest reliability (rating = 2/2).⁵¹ Video motion analysis was used in one study to successfully distinguish between experts and novices when performing a carotid artery stent simulation, scoring relations to other variables = 2/2, and this was rated highly across most studies.⁵⁰ Postsimulator surveys rated VIST’s fantastic training potential and realism, alongside a decreased procedure time, fluoroscopy time and contrast volume (LoE 1-2/5). Chaer *et al.* used VIST to trial a simulated iliofemoral angioplasty and found an improvement in all technical skill metrics when later performed on real patients (LoE = 3/5).⁵⁵ Hseino *et al.* found that practicing superior femoral angioplasty on VIST translated to real-world performance (LoE = 3/5).⁶¹

EVAR

As shown in [Supplementary Table III](#), nine studies validated four simulators. The 3D printed Aneurysm model simulated patient-specific EVAR and differentiated operative performance between simulation trained and nontrained residents (relations to other variables rating = 2/2).⁶³ Participants decreased procedure time, fluoroscopy time, and contrast used on real patients post simulator use (LoE = 3/5).

Rudarakanchana *et al.* tested VIST on a simulated ruptured AAA (rAAA) with a multidisciplinary team.⁶⁴ Differences between trainee and expert performance were clearly noted, scoring relations to other variables = 2/2. Another group used VIST to assess EVAR on 23 endovascular experts and established a pass/fail standard using the Angoff method (consequences = 2/2).⁶⁵

ANGIO Mentor has been validated by five studies for EVAR simulation.⁶⁶⁻⁷⁰ Saratzis *et al.* used expert reviewed assessment methods scoring content = 2/2.⁷⁰ Vento *et al.* used two expert vascular surgeons to perform simulated cases on ANGIO Mentor, obtaining mean values and comparing this with metrics obtained from residents to differentiate between varying levels of competence (relations to other variables = 2/2).⁶⁸ There was minimal discussion surrounding response processes; however, Descender *et al.* scored 2/2 for this criterion due to a clear thought process regarding assessment design.⁶⁹ The same group achieved an LoE of 4/5 after EVAR rehearsal, as there was a reduction in major and minor errors post simulator

use, having a direct effect on patient outcomes by reducing in-hospital mortality by 2%. The rest of the studies achieved an LoE of 2/5, due to a reduction in simulated fluoroscopy and procedural time, alongside fewer endoleaks and postsimulator use.

General endovascular procedures

Seven studies validated six simulators for general endovascular procedures (see [Supplementary Table III](#)). A Pulsatile Fresh Frozen Cadaver model scored an LoE = 1/5 as participants rated the simulation useful.⁷¹ Relations to other variables scored 2/2 as a difference in score was shown between endovascular experts and trainees. This was re-assessed in another study where an improvement in performance awarded an LoE of 2/5.⁷² Content was scored 2/2 as there was reference to a prior assessment method that demonstrated excellent face validity.

The 3D-Printed Endovascular Simulation model tested basic endovascular skills, using a low fidelity bench simulator on 96 endovascular physicians.⁷³ Favorable postfeedback responses scored an LoE = 1/5.

Nontechnical skills training (NTS)

For NTS, four studies were found for three simulators (see [Supplementary Table IV](#)).

The Resusci Anne Simulator simulated a rAAA.⁷⁴ A content score of 1/2 was awarded as simulation supervisors continuously evaluated methods of assessment. This process was poorly defined. There was a direct change to patient outcomes after the simulation scenario was introduced, including reduced door to occlusion time, door to needle time, and reduced 30-d mortality (LoE = 4/5).

ORCAMP was used in two studies assessing trainee sympathetic tone and teamwork performance.^{75,76} Content ratings were limited (0-1/2) due to reference of a prior validated assessment criterion but limited data. Relations to other variables scored highly (rating = 2/2) for both. Ramjeeawom *et al.*⁷⁵ scored 2/2 for response processes due to a clear thought process regarding quality control methods and test

standardization. An improvement in knowledge and team-work skills scored an LoE = 2/5.

Simulation courses

As in [Supplementary Table IV](#), seven studies validated seven training courses.

Robinson *et al.* designed a 1.5-d simulation course on open and endovascular procedures on a variety of simulation modalities.⁷⁷ The course was able to differentiate between residents and vascular fellows (relations to other variables = 2/2). Post-test procedural knowledge and self-rated procedural competence increased (LoE = 2/5).

Strøm *et al.* held a 1-d intensive course for EVAR sizing and stent-graft selection, using a CT Angiography computer program.⁷⁸ The test score comprised a subscore for anatomy measurements and modular stent-graft selection, which was devised by experts (content = 2/2). An improvement in knowledge and skill was recorded, scoring an LoE = 2/5.

The EduCas course was a 2-d simulation teaching for carotid stenting using ANGIO Mentor.⁷⁹ A previously validated rating scale by experts scored content = 2/2. An improvement in performance was shown by decreased procedural, fluoroscopy, and delivery-retrieval time of the embolic protection device (LoE = 2/5).

Discussion

This systematic review has identified a whole scope of simulators and courses that have been evaluated for their validity and translational outcomes. VR simulators, in particular ANGIO Mentor, were widely validated and translated to real-world performance for a variety of endovascular procedures. There is a need for novel, high fidelity simulators that offer greater sensory and haptic feedback for basic endovascular skills to address this paradigm shift from open to endovascular repair.⁸⁰ Despite this, it is still vital to ensure trainees are proficient in open vascular skills as these are procedures more likely to be performed. Therefore, it is important to remember that there is still a need for highly validated and effective open vascular simulators. Furthermore, the continuing development of surgical technology offers technical endovascular skill feedback such as fluoroscopy time, contrast volume, and even video-motion analysis to track hand-eye coordination.⁵¹ This has been demonstrated in this review to improve performance, translate skills to the operating theater, and improve patient outcomes, seen across multiple specialties where a transition to endovascular procedures has become more common. This overlap between specialties was highlighted in this review. For example, a considerable number of angioplasty and catheterization simulators were seen in the literature to test the skill of interventional radiology and cardiology trainees, often overlapping with vascular surgeons.^{81,82} Cadaver models are known to replicate procedures based on tissue feel and anatomy, permitting high fidelity training and hence were ranked most highly on a trainee review of simulator popularity, improving the confidence and competence for trainees with limited operative experience.⁸³ However, only four cadaver models were seen in this review,

not offering a well-balanced perspective on the validity and educational impact compared to other models. There is a need for greater implementation of cadaver models alongside or in combination with other simulation modalities by training institutions and designers.

VA models were the most popular simulators, mainly due to a low fidelity and simple design, alongside being a fundamental vascular procedure, as highlighted on a vascular surgery curriculum needs assessment.⁸ This assessment did identify other common open procedures, such as peripheral bypass, open embolectomy, and amputation, but these lacked available simulators in the literature. Open vascular simulators have met more resistance than endovascular counterparts, which assess technical skills via VR or computer software, which is often difficult to replicate for open procedures.¹² This explains the greater proportion of commercially available, high fidelity endovascular simulators in the literature compared to low-fidelity, open vascular models.

The rise in endovascular procedures has seen a subsequent rise in complexity and error rate, in particular when open and endovascular surgery are combined, often involving a diverse range of health professionals.⁸⁴ NTS or human factors such as teamwork, leadership, situational awareness, and communication, play an important role in patient safety, accounting for up to 85% of major errors in some studies.⁸⁵ However, only three simulators tested NTS; hence, it was neglected in this review.⁸⁶ Rudarakanchana *et al.* used a fully immersive Angiosuite combined with a high-fidelity VR simulator to simulate a rAAA scenario.⁶⁴ This was rated as highly realistic and tested the technical and nontechnical skills of a multidisciplinary team under realistic conditions. There needs to be a greater implementation of NTS testing within vascular surgery, given the variety of simulation modalities and courses available, and this combination of disciplines is highly recommended and rated by experts, despite the high costs of implementation and maintenance.⁸⁷

The use of modern definitions that identify validity as a unitary construct is vital in the in-depth assessment of simulators and their suitability for examining and training vascular surgeons.⁸⁸ This report featured Messick's validity criteria, which has been widely utilized as an acceptable and appropriate method for evaluating simulation tools within the literature, not only within vascular surgery but also within many other surgical specialties. Additionally, it was recommended as a standard by the American Educational Research Association.¹⁸⁻²⁶ Nevertheless, this was clearly limited as only one study (1.3%) followed Messick's modern concept of validity and featured all five validity criteria.⁶⁵ This corresponds to a surgical review that found only 6.6% of studies used the modern concept of validity.⁸⁹ None of the studies in this review showed complete validity for all domains, although Lawaetz *et al.*²⁹ achieved highly by scoring 2/2 for four of the validity criteria. A significant proportion of studies scored highly (rating = 2/2) for content (46%) and relations to other variables (76%). This was often due to an expert review of assessment methods, reference to a previously validated study, and comparing performance between novice and expert participants. Internal structure reporting was weak as many studies discussed measures of reliability, but only a small proportion (6.6%) scored 2/2 by using multiple measures

to calculate interitem and test-retest reliability. Only two studies (2.6%)^{29,65} scored high validity ratings for consequences criteria, the most neglected validity criteria, by setting a pass-fail score. This criterion was vital to set a certain trainee competence level before practice on real patients.²⁵ There was also a large population of novices featuring as study participants and rated simulators on their realism and effectiveness, meaning that the ratings of some simulators may not be entirely reflective of vascular experts.

Overall, 25% of studies used conventional definitions such as the face, content, and construct validity.⁹⁰ These definitions are outdated, possibly due to many studies being performed by clinicians without guidance from a medical education specialist or being published before the wide implementation of Messick's framework.^{25,90} Furthermore, commonly used definitions such as face and content validity are more subjective measures of validity, alongside there being no clear consensus on the exact definition of these measures, offering little educational relevance.^{90,91} This corresponds to similar reviews focused on urology, open vascular surgery, and additionally endovascular surgery.^{25,91,92} Another issue was that a significant proportion of studies (74%) lacked a formal validation process and were purely descriptive. This was evident in other specialties where reviews of ophthalmology and orthopedics showed a 45% and 38% prevalence of descriptive studies, respectively.^{93,94} This demonstrates that the validity evidence for simulation in vascular surgery is poor, and there needs to be greater use of the modern validity framework (Table 1).⁸⁹

Only seven simulators demonstrated improvement in the operating theater, patient outcomes, and skill retention (LoE 3+/5), with the majority of studies scoring an LoE = 1-2/5. The current literature demonstrates that simulation-based training has the potential to largely improve patient outcomes, but the current reporting is limited in this regard, and further research is needed to investigate the real-world transfer of skills.^{25,92}

Surgical simulation has significantly developed over recent years and offers exciting prospects for the future, such as augmented reality and 3D printed patient-specific models. These have been seen in this review to improve patient outcomes and are set to become more integrated into the vascular surgery curriculum and improve procedural training.⁹⁵ Simulation-based training has been shown to reduce the initial phase of the learning curve, but many trials in this review had a small sample size and did not include an adequate follow-up time. Future trials must validate simulators in large, well-designed randomized control trials with adequate follow-up time. This has been seen in the Simulation in Urological Training and Education (SIMULATE) trial,^{96,97} which assessed whether simulation reduced the number of procedures to reach technical skill proficiency and positively impacted patient outcomes. This offers high relevancy to the future of vascular surgery training.⁸⁵

Limitations

Despite a specific and methodical search criterion, the initial search result included irrelevant articles which focused solely on other endovascular specialties. Instead, the search terms

could feature negative Boolean operators to exclude such articles. The search also focused on more technical aspects of simulation-based training and could perhaps be restructured to allow a balance of nontechnical skills, which is known to complement the technical performance. Many articles featured outdated validation definitions, which were purely descriptive, lacked translational outcomes to real-world performance and lacked specialized expert opinions, which had a significant impact on reaching a conclusion. Finally, the heterogeneity of study methods and outcomes did not allow for a quantitative meta-analysis.

Conclusions

Simulation-based training in vascular surgery is a continuously developing field and of growing importance due to advances in technology and reduced training opportunities, exacerbated by the COVID-19 pandemic. This review applied the modern concept of validity criteria and found a vast number of studies validating simulation models and training courses. ANGIO Mentor was the most validated simulator, collectively demonstrating an improvement in real-world performance, knowledge, patient outcomes, and skill retention. In summary, the future of vascular surgery simulation has great potential. However, too few studies utilized the modern concept of validity framework, and prospective studies require greater use of this, alongside a longer follow-up time of trainee skill retention and discussion of cost-effectiveness. This is needed in order to effectively integrate simulation models into the current vascular surgery curriculum and methods of assessment.

Supplementary Materials

The following references [98–122] are cited in Supplementary Tables.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jss.2022.05.009>.

Author Contributions

AH and AA were involved in the initial planning and conceptualization of the review. AH conducted the literature search, alongside data collection and writing of the manuscript. AA and BK both provided guidance and revisions to the manuscript. KA and PD oversaw the project and provided guidance and access to necessary resources. All authors read and approved the final manuscript.

Disclosure

None.

Funding

None.

REFERENCES

- Fuller FW. The aviation paradigm and surgical education. *J Am Coll Surg*. 2005;201:110–117, 202. United States 2006. p. 200; author reply.
- Bismuth J, Donovan MA, O'Malley MK, et al. Incorporating simulation in vascular surgery education. *J Vasc Surg*. 2010;52:1072–1080.
- Lamont PM, Scott DJ. The impact of shortened training times on the discipline of vascular surgery in the United Kingdom. *Am J Surg*. 2005;190:269–272.
- Zingaretti N, Contessi Negrini F, Tel A, Tresoldi MM, Bresadola V, Parodi PC. The impact of COVID-19 on plastic surgery residency training. *Aesth Plast Surg*. 2020;44:1381–1385.
- Hoefer AC, Bouchagiar J, Goltz JP, et al. Development of an endovascular training model for simulation of eva procedures using 3D rapid prototyping for the production of exchangeable patient specific anatomic models. *Eur J Vasc Endovascular Surg*. 2019;58(Suppl 2):e290–e292.
- Suttie S, Guthrie G, Wilson A, Jamieson R, Marron C. Vascular training in Scotland and Northern Ireland: challenges for the introduction of the vascular curriculum. *Int J Surg*. 2013;11:735.
- Kashyap VS, Ahn SS, Petrik PV, Moore WS. Current training and practice of endovascular surgery: a survey. *Ann Vasc Surg*. 2001;15:294–305.
- Nayahangan LJ, Van Herzele I, Konge L, et al. Achieving consensus to define curricular content for simulation based education in vascular surgery: a Europe wide needs assessment initiative. *Eur J Vasc Endovasc Surg*. 2019;58:284–291.
- Joels CS, Langan EM, Cull DL, Kalbaugh CA, Taylor SM. Effects of increased vascular surgical specialization on general surgery trainees, practicing surgeons, and the provision of vascular surgical care. *J Am Coll Surg*. 2009;208:692–697. quiz 7.e1; discussion reply 7-9.
- Nicholas R, Humm G, MacLeod KE, et al. Simulation in surgical training: prospective cohort study of access, attitudes and experiences of surgical trainees in the UK and Ireland. *Int J Surg*. 2019;67:94–100.
- Lederle FA, Freischlag JA, Kyriakides TC, et al. Outcomes following endovascular vs open repair of abdominal aortic aneurysm: a randomized trial. *JAMA*. 2009;302:1535–1542.
- Pandey VA, Wolfe JH. Expanding the use of simulation in open vascular surgical training. *J Vasc Surg*. 2012;56:847–852.
- Arora A, Lau LY, Awad Z, Darzi A, Singh A, Tolley N. Virtual reality simulation training in Otolaryngology. *Int J Surg*. 2014;12:87–94.
- Gould DA. Training on simulators: limitations and relevance. *Eur J Vasc Endovascular Surg*. 2007;33:533–535.
- Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6:e1000097.
- Messick S. Validity of psychological assessment: validation of inferences from persons' responses and performances as scientific inquiry into score meaning. *Am Psychol*. 1995;50:741–749.
- Competency-based training and simulation: making a "valid" argument. *J Endourology*. 2018;32:84–93.
- Patel EA, Aydin A, Cearn M, Dasgupta P, Ahmed K. A systematic review of simulation-based training in neurosurgery, Part 1: cranial neurosurgery. *World Neurosurg*. 2020;133:e850–e873.
- Patel EA, Aydin A, Cearn M, Dasgupta P, Ahmed K. A systematic review of simulation-based training in neurosurgery, Part 2: spinal and pediatric surgery, neurointerventional radiology, and nontechnical skills. *World Neurosurg*. 2020;133:e874–e892.
- Lee R, Raison N, Lau WY, et al. A systematic review of simulation-based training tools for technical and non-technical skills in ophthalmology. *Eye (Lond)*. 2020;34:1737–1759.
- Vaidya A, Aydin A, Ridgley J, Raison N, Dasgupta P, Ahmed K. Current status of technical skills assessment tools in surgery: a systematic review. *J Surg Res*. 2020;246:342–378.
- Hovgaard LH, Al-Shahrestani F, Andersen SAW. Current evidence for simulation-based training and assessment of myringotomy and ventilation tube insertion: a systematic review. *Otol Neurotol*. 2021;42:e1188–e1196.
- Aydin A, Baig U, Al-Jabir A, Sarica K, Dasgupta P, Ahmed K. Simulation-based training models for urolithiasis. A systematic review. *J Endourology*. 2021;35:1098–1117.
- Davies MG. *Endovascular simulation: a systematic review of the validity evidence, methodology, quality and outcomes*. Chicago, IL: University of Illinois at Chicago; 2019.
- Lawaetz J, Skovbo Kristensen JS, Nayahangan LJ, Van Herzele I, Konge L, Eiberg JP. Simulation based training and assessment in open vascular surgery. *Eur J Vasc Endovascular Surg*. 2020;61:502–509.
- Nayahangan LJ, Lawaetz J, Strøm M, et al. Ensuring competency in open aortic aneurysm repair - development and validation of a new assessment tool. *Eur J Vasc Endovasc Surg*. 2020;59:767–774.
- Beckman TJ, Cook DA, Mandrekar JN. What is the validity evidence for assessments of clinical teaching? *J Gen Intern Med*. 2005;20:1159–1164.
- McGaghie WC, Issenberg SB, Barsuk JH, Wayne DB. A critical review of simulation-based mastery learning with translational outcomes. *Med Educ*. 2014;48:375–385.
- Lawaetz J, Nayahangan LJ, Strøm M, et al. Learning curves and competences of vascular trainees performing open aortic repair in a simulation-based environment. *Ann Vasc Surg*. 2020;72:430–439.
- Robinson WP, Baril DT, Taha O, et al. Simulation-based training to teach open abdominal aortic aneurysm repair to surgical residents requires dedicated faculty instruction. *J Vasc Surg*. 2013;58:247–253.e1-2.
- Duschek N, Assadian A, Lamont PM, et al. Simulator training on pulsatile vascular models significantly improves surgical skills and the quality of carotid patch plasty. *J Vasc Surg*. 2013;57:1148–1154.
- Markovic J, Peyser C, Cavoeres T, Fletcher E, Peterson D, Shortell C. Impact of endovascular simulator training on vascular surgery as a career choice in medical students. *J Vasc Surg*. 2012;55:1515–1521.
- Jensen AR, Milner R, Achildi O, Gaughan J, Wilhite DB, Grewal H. Effective instruction of vascular anastomosis in the surgical skills laboratory. *Am J Surg*. 2008;195:189–194.
- Bartline PB, O'Shea J, McGreevy JM, Mueller MT. A novel perfused porcine simulator for teaching aortic anastomosis increases resident interest in vascular surgery. *J Vasc Surg*. 2017;66:642–648.e4.
- Wilasrusmee C, Lertsithichai P, Kittur DS. Vascular anastomosis model: relation between competency in a laboratory-based model and surgical competency. *Eur J Vasc Endovasc Surg*. 2007;34:405–410.
- Maertens H, Aggarwal R, Desender L, Vermassen F, Van Herzele I. Development of a PROFiciency-based StePwise endovascular curricular training (PROSPECT) program. *J Surg Educ*. 2016;73:51–60.
- Maertens H, Aggarwal R, Moreels N, Vermassen F, Van Herzele I. A proficiency based stepwise endovascular curricular training (PROSPECT) program enhances operative

- performance in real life: a randomised controlled trial. *Eur J Vasc Endovasc Surg*. 2017;54:387–396.
38. Willaert W, Aggarwal R, Bicknell C, et al. Patient-specific simulation in carotid artery stenting. *J Vasc Surg*. 2010;52:1700–1705.
 39. Willaert WI, Aggarwal R, Van Herzeele I, et al. Patient-specific endovascular simulation influences interventionalists performing carotid artery stenting procedures. *Eur J Vasc Endovasc Surg*. 2011;41:492–500.
 40. Willaert W, Aggarwal R, Harvey K, et al. Efficient implementation of patient-specific simulated rehearsal for the carotid artery stenting procedure: part-task rehearsal. *Eur J Vasc Endovasc Surg*. 2011;42:158–166.
 41. Willaert WI, Cheshire NJ, Aggarwal R, et al. Improving results for carotid artery stenting by validation of the anatomic scoring system for carotid artery stenting with patient-specific simulated rehearsal. *J Vasc Surg*. 2012;56:1763–1770.
 42. Willaert WI, Aggarwal R, Daruwalla F, et al. Simulated procedure rehearsal is more effective than a preoperative generic warm-up for endovascular procedures. *Ann Surg*. 2012;255:1184–1189.
 43. Willaert WI, Aggarwal R, Van Herzeele I, et al. Role of patient-specific virtual reality rehearsal in carotid artery stenting. *Br J Surg*. 2012;99:1304–1313.
 44. Rolls AE, Riga CV, Rahim SU, et al. The use of video motion analysis to determine the impact of anatomic complexity on endovascular performance in carotid artery stenting. *J Vasc Surg*. 2019;69:1482–1489.
 45. Gosling AF, Kendrick DE, Kim AH, et al. Simulation of carotid artery stenting reduces training procedure and fluoroscopy times. *J Vasc Surg*. 2017;66:298–306.
 46. Wooster M, Doyle A, Hislop S, et al. REHEARSAL using patient-specific simulation to improve endovascular efficiency. *Vasc Endovascular Surg*. 2018;52:169–172.
 47. Lee JT, Qiu M, Teshome M, Raghavan SS, Tedesco MM, Dalman RL. The utility of endovascular simulation to improve technical performance and stimulate continued interest of preclinical medical students in vascular surgery. *J Surg Educ*. 2009;66:367–373.
 48. Nicholson WJ, Cates CU, Patel AD, et al. Face and content validation of virtual reality simulation for carotid angiography: results from the first 100 physicians attending the Emory NeuroAnatomy Carotid Training (ENACT) program. *Simul Healthc*. 2006;1:147–150.
 49. Dayal R, Faries PL, Lin SC, et al. Computer simulation as a component of catheter-based training. *J Vasc Surg*. 2004;40:1112–1117.
 50. Hsu JH, Younan D, Pandalai S, et al. Use of computer simulation for determining endovascular skill levels in a carotid stenting model. *J Vasc Surg*. 2004;40:1118–1125.
 51. Rolls AE, Riga CV, Bicknell CD, et al. A pilot study of video-motion analysis in endovascular surgery: development of real-time discriminatory skill metrics. *Eur J Vasc Endovasc Surg*. 2013;45:509–515.
 52. Van Herzeele I, Aggarwal R, Choong A, Brightwell R, Vermassen FE, Cheshire NJ. Virtual reality simulation objectively differentiates level of carotid stent experience in experienced interventionalists. *J Vasc Surg*. 2007;46:855–863.
 53. Bech B, Lönn L, Falkenberg M, et al. Construct validity and reliability of structured assessment of endovascular expertise in a simulated setting. *Eur J Vasc Endovasc Surg*. 2011;42:539–548.
 54. Bech B, Lönn L, Schroeder TV, Ringsted C. Fine-motor skills testing and prediction of endovascular performance. *Acta Radiol*. 2013;54:1165–1174.
 55. Chaer RA, Derubertis BG, Lin SC, et al. Simulation improves resident performance in catheter-based intervention: results of a randomized, controlled study. *Ann Surg*. 2006;244:343–352.
 56. Tedesco MM, Pak JJ, Harris EJ, Krummel TM, Dalman RL, Lee JT. Simulation-based endovascular skills assessment: the future of credentialing? *J Vasc Surg*. 2008;47:1008–1011. discussion 14.
 57. Aggarwal R, Black SA, Hance JR, Darzi A, Cheshire NJ. Virtual reality simulation training can improve inexperienced surgeons' endovascular skills. *Eur J Vasc Endovasc Surg*. 2006;31:588–593.
 58. Boyle E, O'Keeffe DA, Naughton PA, Hill AD, McDonnell CO, Moneley D. The importance of expert feedback during endovascular simulator training. *J Vasc Surg*. 2011;54:240–248.e1.
 59. Naughton PA, Aggarwal R, Wang TT, et al. Skills training after night shift work enables acquisition of endovascular technical skills on a virtual reality simulator. *J Vasc Surg*. 2011;53:858–866.
 60. Neequaye SK, Aggarwal R, Brightwell R, Van Herzeele I, Darzi A, Cheshire NJ. Identification of skills common to renal and iliac endovascular procedures performed on a virtual reality simulator. *Eur J Vasc Endovasc Surg*. 2007;33:525–532.
 61. Hseino H, Nugent E, Lee MJ, et al. Skills transfer after proficiency-based simulation training in superficial femoral artery angioplasty. *Simul Healthc*. 2012;7:274–281.
 62. Hseino H, Nugent E, Cantwell C, et al. Impact of an assistant on the technical skills of the primary operator in superficial femoral artery angioplasty. *Vasc Endovascular Surg*. 2012;46:635–639.
 63. Torres IO, De Luccia N. A simulator for training in endovascular aneurysm repair: the use of three dimensional printers. *Eur J Vasc Endovasc Surg*. 2017;54:247–253.
 64. Rudarakanchana N, Van Herzeele I, Bicknell CD, et al. Endovascular repair of ruptured abdominal aortic aneurysm: technical and team training in an immersive virtual reality environment. *Cardiovasc Intervent Radiol*. 2014;37:920–927.
 65. Strøm M, Lönn L, Konge L, et al. Assessment of EVAR competence: validity of a novel rating scale (EVARATE) in a simulated setting. *Eur J Vasc Endovasc Surg*. 2018;56:137–144.
 66. Kim AH, Kendrick DE, Moorehead PA, et al. Endovascular aneurysm repair simulation can lead to decreased fluoroscopy time and accurately delineate the proximal seal zone. *J Vasc Surg*. 2016;64:251–258.
 67. Kendrick DE, Gosling AF, Nagavalli A, Kashyap VS, Wang JC. Endovascular simulation leads to efficiency and competence in thoracic endovascular aortic repair procedures. *J Surg Educ*. 2015;72:1158–1164.
 68. Vento V, Cercenelli L, Mascoli C, et al. The role of simulation in boosting the learning curve in EVAR procedures. *J Surg Educ*. 2018;75:534–540.
 69. Desender LM, Van Herzeele I, Lachat ML, et al. Patient-specific rehearsal before EVAR: influence on technical and nontechnical operative performance. A randomized controlled trial. *Ann Surg*. 2016;264:703–709.
 70. Saratzis A, Calderbank T, Sidloff D, Bown MJ, Davies RS. Role of simulation in endovascular aneurysm repair (EVAR) training: a preliminary study. *Eur J Vasc Endovasc Surg*. 2017;53:193–198.
 71. Nesbitt C, Tingle SJ, Williams R, et al. A pulsatile Fresh frozen human cadaver circulation model for endovascular training: a trial of face validity. *Ann Vasc Surg*. 2018;46:345–350.
 72. Nesbitt CI, Tingle SJ, Williams R, et al. Educational impact of a pulsatile human cadaver circulation model for endovascular training. *Eur J Vasc Endovasc Surg*. 2019;58:602–608.
 73. Mafeld S, Nesbitt C, McCaslin J, et al. Three-dimensional (3D) printed endovascular simulation models: a feasibility study. *Ann Transl Med*. 2017;5:42.

74. Aho P, Vikatmaa L, Niemi-Murola L, Venermo M. Simulation training streamlines the real-life performance in endovascular repair of ruptured abdominal aortic aneurysms. *J Vasc Surg.* 2019;69:1758–1765.
75. Ramjeeawon A, Sharrock AE, Morbi A, Martin G, Riga C, Bicknell C. Using fully-immersive simulation training with structured debrief to improve nontechnical skills in emergency endovascular surgery. *J Surg Educ.* 2020;77:1300–1311.
76. Bakhsh A, Martin GFJ, Bicknell CD, Pettengell C, Riga C. An evaluation of the impact of high-fidelity endovascular simulation on surgeon stress and technical performance. *J Surg Educ.* 2019;76:864–871.
77. Robinson WP, Doucet DR, Simons JP, et al. An intensive vascular surgical skills and simulation course for vascular trainees improves procedural knowledge and self-rated procedural competence. *J Vasc Surg.* 2017;65:907–915.e3.
78. Strøm M, Rasmussen JL, Nayahangan LJ, et al. Learn EVAR sizing from scratch: the results of a one-day intensive course in EVAR sizing and stent graft selection for vascular trainees. *Vascular.* 2020;28:342–347.
79. Van Herzelee I, Aggarwal R, Neequaye S, et al. Experienced endovascular interventionalists objectively improve their skills by attending carotid artery stent training courses. *Eur J Vasc Endovasc Surg.* 2008;35:541–550.
80. Amin A, Salsamendi J, Sullivan T. High-fidelity endovascular simulation. *Tech Vasc Interv Radiol.* 2019;22:7–13.
81. Svensson LG, Cambria RP. Expanding surgical options using minimally invasive techniques for cardio-aortic and aortic procedures. *Arch Surg.* 1998;133:1160–1165.
82. Murphy TP, Soares GM. The evolution of interventional radiology. *Semin Intervent Radiol.* 2005;22:6–9.
83. Duran C, Bismuth J, Mitchell E. A nationwide survey of vascular surgery trainees reveals trends in operative experience, confidence, and attitudes about simulation. *J Vasc Surg.* 2013;58:524–528.
84. Rudarakanchana N, Van Herzelee I, Desender L, Cheshire NJW. Virtual reality simulation for the optimization of endovascular procedures: current perspectives. *Vasc Health Risk Manag.* 2015;11:195–202.
85. Iqbal MH, Khan O, Aydın A. Editorial commentary: simulation-based training in orthopaedic surgery: current evidence and limitations. *Arthroscopy. J Arthroscopic Relat Surg.* 2021;37:1008–1010.
86. Hull L, Arora S, Aggarwal R, Darzi A, Vincent C, Sevdalis N. The impact of nontechnical skills on technical performance in surgery. *A Syst Rev J Am Coll Surgeons.* 2012;214:214–230.
87. Kassab E, Tun JK, Arora S, et al. "Blowing up the barriers" in surgical training: exploring and validating the concept of distributed simulation. *Ann Surg.* 2011;254:1059–1065.
88. Noureldin YA, Lee JY, McDougall EM, Sweet RM. Competency-based training and simulation: making a "valid" argument. *J Endourology.* 2018;32:84–93.
89. Borgersen NJ, Naur TMH, Sørensen SMD, et al. Gathering validity evidence for surgical simulation: a systematic review. *Ann Surg.* 2018;267:1063–1068.
90. Downing SM. Face validity of assessments: faith-based interpretations or evidence-based science? *Med Educ.* 2006;40:7–8.
91. Aydın A, Shafi AMA, Shamim Khan M, Dasgupta P, Ahmed K. Current status of simulation and training models in urological surgery: a systematic review. *J Urol.* 2016;196:312–320.
92. See KW, Chui KH, Chan WH, Wong KC, Chan YC. Evidence for endovascular simulation training: a systematic review. *Eur J Vasc Endovasc Surg.* 2016;51:441–451.
93. Lee R, Raison N, Lau WY, et al. A systematic review of simulation-based training tools for technical and non-technical skills in ophthalmology. *Eye.* 2020;34:1737–1759.
94. Morgan M, Aydın A, Salih A, Robati S, Ahmed K. Current status of simulation-based training tools in orthopedic surgery: a systematic review. *J Surg Education.* 2017;74:698–716.
95. Veith FJ. A look at the future of vascular surgery. *J Vasc Surg.* 2016;64:885–890.
96. Aydın A, Ahmed K, Abe T, et al, The SIMULATE Trial Group. Effect of Simulation-based Training on Surgical Proficiency and Patient Outcomes: A Randomised Controlled Clinical and Educational Trial. *Eur Urol.* 2022;81:385–393.
97. Aydın A, Ahmed K, Van Hemelrijck M, Ahmed HU, Khan MS, Dasgupta P. Simulation in Urological Training and Education (SIMULATE): protocol and curriculum development of the first multicentre international randomized controlled trial assessing the transferability of simulation-based surgical training. *BJU Int.* 2020;126:202–211.
98. Aparajita R, Zayed MA, Casey K, Dayal R, Lee JT. Development and implementation of an introductory endovascular training course for medical students. *Ann Vasc Surg.* 2011;25:1104–1112.
99. Lee JT, Son JH, Chandra V, Lilo E, Dalman RL. Long-term impact of a preclinical endovascular skills course on medical student career choices. *J Vasc Surg.* 2011;54:1193–1200.
100. Hislop SJ, Hsu JH, Narins CR, et al. Simulator assessment of innate endovascular aptitude versus empirically correct performance. *J Vasc Surg.* 2006;43:47–55.
101. Dawson DL, Lee ES, Hedayati N, Pevac WC. Four-year experience with a regional program providing simulation-based endovascular training for vascular surgery fellows. *J Surg Educ.* 2009;66:330–335.
102. Dawson DL, Meyer J, Lee ES, Pevac WC. Training with simulation improves residents' endovascular procedure skills. *J Vasc Surg.* 2007;45:149–154.
103. Passman MA, Fleser PS, Dattilo JB, Guzman RJ, Naslund TC. Should simulator-based endovascular training be integrated into general surgery residency programs? *Am J Surg.* 2007;194:212–219.
104. Moorthy K, Munz Y, Adams S, Pandey V, Darzi A, Group IC-SMsHS. Self-assessment of performance among surgical trainees during simulated procedures in a simulated operating theater. *Am J Surg.* 2006;192:114–118.
105. Datta V, Bann S, Beard J, Mandalia M, Darzi A. Comparison of bench test evaluations of surgical skill with live operating performance assessments. *J Am Coll Surg.* 2004;199:603–606.
106. Kärkkäinen JM, Sandri G, Tenorio ER, et al. Simulation of endovascular aortic repair using 3D printed abdominal aortic aneurysm model and fluid pump. *Cardiovasc Intervent Radiol.* 2019;42:1627–1634.
107. Borger van der Burg BLS, Hörer TM, Eefting D, et al. Vascular access training for REBOA placement: a feasibility study in a live tissue-simulator hybrid porcine model. *J R Army Med Corps.* 2019;165:147–151.
108. Pandey VA, Black SA, Lazaris AM, et al. Do workshops improve the technical skill of vascular surgical trainees? *Eur J Vasc Endovascular Surg.* 2005;30:441–447.
109. Schwartz S, de Virgilio M, Chisum P, et al. A prospective randomized study assessing optimal method for teaching vascular anastomoses. *Ann Vasc Surg.* 2014;28:1087–1093.
110. Black SA, Nestel DF, Horrocks EJ, et al. Evaluation of a framework for case development and simulated patient training for complex procedures. *Simul Healthc.* 2006;1:66–71.
111. Malas T, Al-Atassi T, Brandys T, Naik V, Lapierre H, Lam BK. Impact of visualization on simulation training for vascular anastomosis. *J Thorac Cardiovasc Surg.* 2018;155:1686–1693.e5.
112. Berry M, Lystig T, Beard J, Klingestierna H, Reznick R, Lönn L. Porcine transfer study: virtual reality simulator training

- compared with porcine training in endovascular novices. *Cardiovasc Intervent Radiol*. 2007;30:455–461.
113. Robinson WP, Schanzer A, Cutler BS, et al. A randomized comparison of a 3-week and 6-week vascular surgery simulation course on junior surgical residents' performance of an end-to-side anastomosis. *J Vasc Surg*. 2012;56:1771–1781.
 114. Duran C, Estrada S, O'Malley M, et al. The model for Fundamentals of Endovascular Surgery (FEVS) successfully defines the competent endovascular surgeon. *J Vasc Surg*. 2015;62:1660–1666.e3.
 115. Taher F, Plimon M, Isaak A, et al. Ultrasound-guided percutaneous arterial puncture and closure device training in a pulsatile model. *J Surg Educ*. 2020;77:1271–1278.
 116. Sigounas VY, Callas PW, Nicholas C, et al. Evaluation of simulation-based training model on vascular anastomotic skills for surgical residents. *Simul Healthc*. 2012;7:334–338.
 117. Rosenthal R, Mujagic E, Jacob AL, Seelos R, Schäfer J, Gürke L. Impact of an intensive 2-day endovascular training course on technical performance of trainees. *Ann Vasc Surg*. 2013;27:1173–1181.
 118. Fletcher BP, Gusic ME, Robinson WP. Simulation training incorporating a pulsatile carotid endarterectomy model results in increased procedure-specific knowledge, confidence, and comfort in post-graduate trainees. *J Surg Educ*. 2020;77:1289–1299.
 119. Aeckersberg G, Gkremoutis A, Schmitz-Rixen T, Kaiser E. The relevance of low-fidelity virtual reality simulators compared with other learning methods in basic endovascular skills training. *J Vasc Surg*. 2019;69:227–235.
 120. Sidhu R, Weir-McCall J, Cochennec F, Riga C, DiMarco A, Bicknell CD. Evaluation of an electromagnetic 3D navigation system to facilitate endovascular tasks: a feasibility study. *Eur J Vasc Endovasc Surg*. 2012;43:22–29.
 121. Safi AF, Safi S, Tayeh M, et al. Vein graft interposition: a training model using gradually thawed cryopreserved vessels. *J Craniofac Surg*. 2019;30:e213–e216.
 122. O'Toole RV, Playter RR, Krummel TM, et al. Measuring and developing suturing technique with a virtual reality surgical simulator. *J Am Coll Surg*. 1999;189:114–127.