



Impact of inter-ethnic anatomical findings on current surgical outcome, established biomarkers, and future artificial intelligence algorithms

Carsten H. Meyer^{1,2} · Sandeep Saxena³

Received: 4 December 2022 / Revised: 4 December 2022 / Accepted: 20 December 2022
© The Author(s) 2023

Dear Editors,

Laviers et al. investigated in a multi-center study the impact of ethnicity on post-operative anatomical closure rate following macular hole surgery and determined that Black patients had an approximately twofold lower anatomic success rate than White patients [1]. Anatomic failure occurred in 38.5% of Black patients versus 12.6% of White patients, while this difference remained statistically significant despite adjustment for predefined demographic or socio-economic status as well as confounding anatomical variables.

Since the early beginning of optical coherence tomography (OCT), we learned that the neuroretinal anatomy of the foveola may vary within certain conditions, e.g., foveal hypoplasia in oculocutaneous albinism [2]. Previous studies also determined a variation in the foveal morphology among different ethnic cohorts: the eyes of Black patients presented on OCT a larger, deeper, and broader foveal pits associated with a thinner central foveal neuroretinal thickness compared to White and Asian counterparts [3]. Murphy et al. found that even the size in unilateral macular holes correlated with the foveal floor of the fellow eye [4].

In addition, more recent OCT angiography (OCTA) investigations [5] also demonstrated a greater variability in quantitative measures of the retinal microvasculature among racial and ethnic groups: Hispanic, Black, and Asian patients had larger FAZ areas, in comparison to White participants.

These differences persisted, even after adjusting for gender, age, and status in comparison to White patients.

These important neuroretinal and vascular OCT findings suggest that race and ethnicity may be important factors influencing established surrogated biomarkers [6], when evaluating the foveal morphology of healthy and diseased retina. The impact of these inter-ethnic findings is yet to be explored; however, it is conceivable that normative OCT parameters, biological determinants, and established threshold biomarkers may vary among different race and ethnicity groups and needed to elucidate in greater details to establish precise algorithms of future artificial intelligence systems.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

✉ Carsten H. Meyer
Meyer_Eye@yahoo.com

¹ Macula Center Grischun, Bahnhofstrasse 4, CH-7000 Chur, Switzerland

² Department of Ophthalmology, Philipps University, Marburg, Germany

³ Department of Ophthalmology, King George's Medical University, Lucknow, India

References

1. Laviers H, Papavasileiou E, Bruce C, et al. (2023) The effect of ethnicity on anatomic success following macular hole surgery: a multicentre cohort study” for Graefe’s Archive for Clinical and Experimental Ophthalmology. <https://doi.org/10.1007/s00417-022-05950-w>
2. Meyer CH, Lapolice DJ, Freedman SF (2002) Foveal hypoplasia in oculocutaneous albinism demonstrated by optical coherence tomography. *Am J Ophthalmol* 133(3):409–410
3. Patel PJ, Foster PJ, Grossi CM et al (2016) Spectral-domain optical coherence tomography imaging in 67 321 adults: associations with macular thickness in the UK Biobank Study. *Ophthalmology* 123:829–840. <https://doi.org/10.1016/j.ophtha.2015.11.009>
4. Murphy DC, Melville HJR, George G et al (2021) The association between foveal floor measurements and macular hole size. *Ophthalmol Retina* 5:680–686
5. Laotaweerungsawat S, Psaras C, Haq Z, Liu X, Stewart JM (2021) Racial and ethnic differences in foveal avascular zone in diabetic and nondiabetic eyes revealed by optical coherence tomography angiography. *PLoS One* 16(10):e0258848. <https://doi.org/10.1371/journal.pone.0258848> eCollection 2021
6. Phadikar P, Saxena S, Ruia S et al (2017) The potential of spectral domain optical coherence tomography imaging based retinal biomarkers. *Int J Retina Vitreous* 3:1. <https://doi.org/10.1186/s40942-016-0054-7>

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.