



King's Research Portal

DOI:

[10.1007/s00011-023-01722-2](https://doi.org/10.1007/s00011-023-01722-2)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Freidin, M. B., Cheetham, N., Duncan, E. L., Steves, C. J., Doores, K. J., Malim, M. H., Rossi, N., Lord, J. M., Franks, P. W., Borsini, A., Granville Smith, I., Falchi, M., Pariente, C., & Williams, F. M. K. (2023). Long-COVID fatigue is not predicted by pre-pandemic plasma IL-6 levels in mild COVID-19. *Inflammation Research*, 72(5), 947-953. <https://doi.org/10.1007/s00011-023-01722-2>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Long-COVID fatigue is not predicted by pre-pandemic plasma IL-6 levels in mild COVID-19 infection

Maxim B Freidin^{1,2}, Nathan Cheetham¹, Emma Duncan¹, Claire J Steves¹, Katherine J Doores³, Michael H Malim³, Niccolo Rossi¹, Janet M Lord^{4,5}, Paul W Franks^{6,7}, Alessandra Borsini⁸, Isabelle Granville Smith¹, Mario Falchi¹, Carmine Pariante⁸, Frances MK Williams¹

1 – Department of Twin Research and Genetic Epidemiology, School of Life Course Sciences, King's College London, London, UK

2 – Department of Biology, School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK

3 – Department of Infectious Diseases, School of Immunology & Microbial Sciences, King's College London, London, UK

4 – MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, UK

5 – NIHR Birmingham Biomedical Research Centre, University Hospital Birmingham and University of Birmingham, Birmingham, UK

6 – Lund University Diabetes Center, Lund University, Malmö, Sweden

7 – Department of Clinical Sciences, Lund University, Malmö, Sweden

8 – Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

Correspondence to: Dr Maxim Freidin, Department of Twin Research and Genetic Epidemiology, 4th Floor, Block D, South Wing St Thomas' Hospital, King's College London, London, UK

tel. +44(0)20 7882 6453; email: maxim.freydin@kcl.ac.uk

Abstract

Objective and design: Fatigue is a prominent symptom in the general population and may follow viral infection, including SARS-CoV2 infection which causes COVID-19. Chronic fatigue lasting more than three months is a major symptom of the long-COVID syndrome. The mechanisms underlying long-COVID fatigue are unknown. We hypothesized that the development of long-COVID chronic fatigue is driven by the pro-inflammatory immune status of an individual prior to COVID-19 infection.

Subjects and methods: We tested this hypothesis by analysing pre-pandemic plasma levels of IL-6, which plays a key role in persistent fatigue, in N = 1274 community dwelling adults from TwinsUK. Subsequent COVID-19 positive and COVID-19 negative participants were categorized based on antigen and antibody testing. Chronic fatigue was assessed using the Chalder Fatigue Scale.

Results: COVID-19 positive participants exhibited mild symptoms of infection. Chronic fatigue was a prevalent symptom among this population and was significantly higher in the COVID-19 positive participants than COVID-19 negative participants (17% vs 11%, respectively; $p = 0.001$). The qualitative nature of chronic fatigue as determined by individual questionnaire responses was similar in COVID-19 positive and COVID-19 negative participants. Pre-pandemic plasma IL-6 levels were positively associated with chronic fatigue in COVID-19 negative, but not COVID-19 positive individuals. Raised BMI was associated with chronic fatigue in COVID-19 positive participants.

Conclusions: We found evidence that pre-existing increased levels of IL-6 provide the ground for chronic fatigue symptoms but there was no specific link seen with mild COVID-19 status. Elevated BMI increased the risk of chronic fatigue in mild COVID-19 infection consistent with previous reports.

KEY WORDS

long-COVID, chronic fatigue, IL-6, inflammation, BMI

Introduction

Fatigue is a common symptom of COVID-19 infection and is one of its most pronounced acute and post-acute clinical manifestations (1). Long lasting manifestations – so called long-COVID – is a subject of intense interest. Affected individuals report fatigue along with shortness of breath, headache and loss of sense of taste and smell (2). Mechanisms underlying the persistence of long-COVID symptoms are yet to be determined. To date, exploration of blood-borne biomarkers following COVID-19 infection has not revealed an association with inflammatory markers or white cell count (3).

The cytokine interleukin (IL)-6 has a recognized role in fatigue development in many clinical settings including autoimmune inflammatory arthritis (reviewed in (4)) and cancer (5). It is an established driver of acute COVID-19 infection and treatment with anti-IL-6 monoclonal antibodies reduces mortality in severe COVID-19 illness (6, 7). IL-6 is considered a key mediator of neuropsychiatric symptoms of long-COVID, including fatigue (8). Finally, a Mendelian randomization study of depressive symptoms has suggested that IL-6 manifests a causal influence on fatigue, as well as sleep problems and suicidality (9).

We hypothesized that long-COVID fatigue is driven at least in part by the pre-existing immune status of an individual. We have shown previously that chronic fatigue induced by treatment with interferon-alpha (IFN- α) for chronic hepatitis B viral infection is predicted by higher baseline IL-6 and IL-10 levels, as well as an exaggerated elevation of IL-6 and IL-10, in response to treatment (10). In addition, increasing age is a major risk factor for both infection with SARS-CoV2 and COVID-19 illness (2, 11). Aging has been associated with elevated levels of pro-inflammatory cytokines such as TNF- α and IL-6 and reduced levels of anti-inflammatory mediator IL-10 (12). (13). An age-related, chronic, pro-inflammatory milieu may mediate the risk for susceptibility to COVID-19 and long-COVID fatigue. To test this hypothesis, we analysed levels of pro-inflammatory cytokine IL-6 in plasma collected prior to the SARS-CoV2 pandemic in a longitudinal cohort sample of UK adults, who were assessed during the pandemic for chronic fatigue symptoms and COVID-19 infection.

Material and Methods

Sample and phenotyping

Participants were selected from the UK Twin Registry (TwinsUK), an adult cohort which has been shown to be representative of the general population for lifestyle and health-related traits (14). Ethics permission was obtained and participants have provided fully informed consent; the Declaration of Helsinki was adhered to. The registry comprises 14,500 same sex mono- and dizygotic

twin volunteers from the general UK population recruited from previous twin registers and national media campaigns. The cohort is predominantly female (83%), and mainly of Northern European descent (15). Samples and data for this project were collected as part of ongoing research initiatives into inflammaging as well as more recent research into COVID-19. Participant selection and grouping is depicted on Figure 1. Participants (n = 5755) were invited to complete a COVID-19 Personal Experience (CoPE) Questionnaire which included questions about COVID-19 infection, related symptoms and fatigue over the previous three months (16). CoPE was completed in multiple waves during the pandemic: April 2020, August 2020, November 2020, and April 2021. Participants also provided serum for COVID-19 antigen and antibody testing at multiple points during the pandemic, with primary collections of n = 506 in April-June 2020, n = 5165 in August 2020, n = 137 in November-December 2020, and n = 4291 in April-May 2021.

We followed guidance on interpretation of antibody test results from the Centres for Disease Control and Prevention (<https://www.cdc.gov/coronavirus/2019-ncov/lab/resources/antibody-tests-guidelines.html>) to assign natural COVID-19 infection status from the results of swab antigen tests; and enzyme-linked immunosorbent assays (ELISA) anti-Nucleocapsid (anti-N) and anti-Spike (anti-S) Roche antibody tests (17) performed at both King's College London and 3rd party laboratories as described previously (18). Individuals with a positive antigen test at any point over the CoPE questionnaire administration period, who were positive for anti-S antibodies prior to self-reported COVID-19 vaccination date, or positive for anti-N antibodies at any point, were classified as COVID-19 positive cases. Individuals with negative antigen or anti-N antibody results, or negative anti-S antibody results before COVID-19 vaccination, were classified as COVID-19 negative, or controls. Those with negative anti-N antibody and negative antigen results but positive anti-S after COVID-19 vaccination were excluded from the analysis as anti-S antibodies are generated in response to vaccination. Individuals with no laboratory antigen or antibody test results were also excluded from the analysis.

Within the CoPE questionnaire the Chalder Fatigue Scale (CFQ) (19) was used to classify participants into those experiencing chronic fatigue and those who did not. CFQ comprises 11 questions concerning physical and mental aspects of fatigue. Reliability of CFQ has been shown in clinical and non-clinical settings (19, 20). CFQ responses were extracted from the CoPE questionnaires administered in August and November 2020. Responses were coded as 0 ("Less than usual", "No more than usual") or 1 ("More than usual", "Much more than usual") followed by summing the scores for different questions and assigning a diagnosis of fatigue to those with summary score of 4 or more (20). Volunteers reporting fatigue at both timepoints were diagnosed as having chronic

fatigue because it lasted 3 months or longer. Those who reported fatigue at a single timepoint were removed from the study.

Pre-pandemic IL-6 levels were ascertained using plasma specimens obtained in 1997-2018 (median 2009). Samples were assayed using Olink Target 96 Inflammation assay (<https://www.olink.com/products-services/target/inflammation/>). Where multiple plasma specimens were available, we selected the most recent, pre-pandemic specimen. IL-6 levels were expressed as normalized protein expression on the Olink arbitrary unit in log₂ scale.

Statistical analysis

A generalized mixed-effects model with chronic fatigue as a dichotomous categorical response variable and IL-6 levels as a predictor was examined, adjusting for age, body mass index (BMI) and sex as fixed effects. Family structure and zygosity were considered as random factors. IL-6 levels were adjusted for age and BMI followed by transformation of the residuals to a normal distribution using *qqnorm* function in R statistical environment. The model was fitted for COVID-19 positive- and COVID-19 negative participants separately. A set of sensitivity analyses was performed. The first investigated sample integrity over time and comprised only plasma samples collected two years before the pandemic; the second was analysis without adjustment of IL-6 levels for age and BMI levels. Finally, we repeated the analysis after excluding individuals with major inflammatory disease (rheumatoid arthritis, systemic lupus erythematosus, ulcerative colitis, and Crohn's disease; n = 11).

Results

Prevalence of fatigue

After selecting participants reporting the same fatigue status at both timepoints, the sample for analysis comprised total n = 1274 participants, of which n=282 were classified COVID-19 positive and n=162 were classified as having chronic fatigue (Table 1). None of the COVID-19 positive participants had been hospitalised so their COVID-19 infection was considered relatively mild. The prevalence of long-term fatigue was 17.4% among COVID-19 positive participants and 11.4% in COVID-19-negative participants (Fisher's exact p = 0.011). COVID-19 positive participants were on average two years younger than COVID-19 negative participants, there was no difference in BMI observed (Table 1). Two (1.1%) and nine (0.7%) cases of major inflammatory disease were reported by those with and without chronic fatigue, respectively (Fisher's exact test p = 0.6408).

Structure of fatigue

The qualitative nature of chronic fatigue was similar in COVID-19 positive and COVID-19 negative participants, with no differences in answer prevalence in the CFQ (Table 2).

IL-6 levels

In the total sample, chronic fatigue was found to be associated with elevated pre-pandemic plasma IL-6 levels, which persisted after adjusting the cytokine levels for age, sex, BMI, and COVID-19 status; as well as with higher levels of BMI (Figure 2).

Stratified analysis established, that pre-pandemic plasma IL-6 levels were elevated in participants with chronic fatigue in the COVID-19 negative group. The same relationship was not seen in the COVID-19 positive group (Table 3). We also calculated prevalence of fatigue in COVID-19 positive and COVID-19 negative participants stratified by high and low levels of pre-pandemic IL-6. We defined high and low IL6 levels as values equal to or above 75% percentile of IL-6 distribution and equal to or below 25% percentile, respectively. The prevalence of fatigue was found to be significantly higher in high IL-6 level group compared to low IL-6 level group in COVID-negative participants (17.5% vs 7.6%, $p = 0.001$); however, no such differences were found in COVID-positive group (19.2% vs 19.0%, $p = 0.999$). Sensitivity analysis of participants having plasma collected no earlier than 2017, showed similarity with the main analysis: $\beta = 0.653 \pm 0.375$, $p = 0.0814$; and $\beta = -0.760 \pm 0.625$, $p = 0.224$ for COVID-19 negative and COVID-19 positive participants, respectively). Sensitivity analysis without adjusting IL-6 levels for age and BMI levels at the time of plasma collection, produced almost identical results to the main analysis: $\beta = 0.321 \pm 0.115$, $p = 0.005$; and $\beta = -0.061 \pm 0.180$, $p = 0.774$ for COVID-19 negative and COVID-19 positive participants, respectively. Sensitivity analysis with excluding cases of major inflammatory disease, produced almost identical results, too: $\beta = 0.307 \pm 0.108$, $p = 0.004$; and $\beta = -0.078 \pm 0.166$, $p = 0.639$, for COVID-19 negative and COVID-19 positive participants, respectively.

BMI

Higher BMI was associated with chronic fatigue in COVID-19 positive participants and in the whole sample (Table 3). To explore the relationship between chronic fatigue and BMI in COVID-19 positive participants, we examined groups by BMI (BMI < 18.5; $n = 6$), healthy weight (BMI > 18.5 and < 25; $n = 115$), overweight (BMI > 25 and < 30; $n = 102$), and obese (BMI ≥ 30 ; $n = 59$). There was a significant difference in the prevalence of these groups between COVID-19 positive cases with and without chronic fatigue ($\chi^2 = 8.3$, $df = 3$, p -value = 0.040; Figure 3). This was largely driven by a lower prevalence of healthy weight and higher prevalence of obesity in COVID-19 positive participants with chronic fatigue.

Discussion

This is among the first studies to examine pre-infective inflammatory cytokine levels and subsequent long-COVID chronic fatigue in participants who experienced mild COVID-19 illness. We found pre-pandemic IL-6 levels to increase the risk of developing fatigue in adult volunteers who had never had COVID-19 illness, however, pre-pandemic IL-6 levels did not predict fatigue in our COVID-19 positive group. This finding is in keeping with other post-viral and post-infective fatigue research and studies in a general population (21, 22). We previously demonstrated that greater pro-inflammatory cytokine elevations in response to IFN- α treatment was associated with developing persistent fatigue (10). In that study we also observed a role for pre-treatment higher cytokine levels of IL-6 and lower IL-10 and the development chronic fatigue – which led to the current hypothesis.

While our COVID-19 positive group were not hospitalized and therefore classified with ‘mild’ illness it is likely IL-6 and other pro-inflammatory cytokines reach high levels during COVID-19 infection (21), well in excess of levels associated with well-known risk factors such as elevated BMI (22). Taken together, these and our findings suggest *any* elevation of circulating pro-inflammatory cytokines, be it a protracted, low-grade inflammatory dysregulation or an acute sickness response to infection will increase chronic fatigue risk, with higher cytokine levels associated with higher risk. Baseline immune status or cytokine levels of an apparently healthy individual may only predict chronic fatigue in the absence of pronounced cytokine elevation, such as an acute sickness response to viral or bacterial infection.

Our hypothesis was that chronic fatigue in community dwelling adults develops after COVID-19 infection because of pre-existing pro-inflammatory immune state, which was not supported by our results.

We did however find a statistically significant association between higher BMI and long-COVID fatigue in COVID-19 positive participants (Table 3). This is consistent with a study demonstrating association between obesity and fatigue while controlling for other potential contributors including IL-6 levels (23). Indeed, almost a third of circulating IL-6 is thought to be produced by adipocytes (24), in keeping with BMI being an important risk factor for chronic fatigue. Of interest, the relationship with BMI and chronic fatigue was not evident in COVID-19 negative participants, an association usually apparent in the population (25).

Allied to this, increasing BMI and advancing age are well-established risk factors for severe COVID-19 disease (23) and both are associated with systemic inflammation (24). Our findings show that following mild COVID-19 infection chronic fatigue does not appear to be qualitatively different from

other forms of chronic fatigue, at least according to the CFQ, which is a well validated tool known to discriminate between clinical and non-clinical conditions (20). Whether this is true for more severe disease requiring hospitalization is unclear.

The most pronounced limitation to this study is the focus on a single cytokine and we do recognize other inflammatory mediators such as TNF- α are likely to contribute to chronic fatigue also.

In summary, our work sheds light on the role of IL-6 in general chronic fatigue but it does not support a specific role for IL-6 levels in the development of chronic fatigue following mild COVID-19 infection.

Acknowledgements

The study was supported by Kennedy Trust grant #KENN 19-20-10. JML is supported by the NIHR Birmingham Biomedical Research Centre, and CMP by the South London and Maudsley NHS Foundation Trust & King's College London Biomedical Research Centre; the views expressed here are those of the authors and not necessarily those of the NHS, NIHR or Department for Health and Social Care. Mario Falchi is supported by MRC grant MR/T004142/1. TwinsUK is funded by the Wellcome Trust, Medical Research Council, Versus Arthritis, European Union Horizon 2020, Chronic Disease Research Foundation (CDRF), Zoe Ltd and the National Institute for Health Research (NIHR) Clinical Research Network (CRN) and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London.

References

1. Carfi A, Bernabei R, Landi F, Gemelli Against C-P-ACSG. Persistent Symptoms in Patients After Acute COVID-19. *JAMA*. 2020;324(6):603-5.
2. Sudre CH, Murray B, Varsavsky T, Graham MS, Penfold RS, Bowyer RC, et al. Attributes and predictors of long COVID. *Nat Med*. 2021;27(4):626-31.
3. Townsend L, Dyer AH, Jones K, Dunne J, Mooney A, Gaffney F, et al. Persistent fatigue following SARS-CoV-2 infection is common and independent of severity of initial infection. *PLoS One*. 2020;15(11):e0240784.
4. Grygiel-Gorniak B, Puszczewicz M. Fatigue and interleukin-6 - a multi-faceted relationship. *Reumatologia*. 2015;53(4):207-12.
5. Kolak A, Kamińska M, Wysokińska E, Surdyka D, Kieszko D, Pakieła M, et al. The problem of fatigue in patients suffering from neoplastic disease. *Contemp Oncol (Pozn)*. 2017;21(2):131-5.
6. Castelnovo L, Tamburello A, Lurati A, Zaccara E, Marrazza MG, Olivetti M, et al. Anti-IL6 treatment of serious COVID-19 disease: A monocentric retrospective experience. *Medicine (Baltimore)*. 2021;100(1):e23582.
7. Abidi E, El Nekidy WS, Alefishat E, Rahman N, Petroianu GA, El-Lababidi R, et al. Tocilizumab and COVID-19: Timing of Administration and Efficacy. *Front Pharmacol*. 2022;13:825749.
8. Kappelmann N, Dantzer R, Khandaker GM. Interleukin-6 as potential mediator of long-term neuropsychiatric symptoms of COVID-19. *Psychoneuroendocrinology*. 2021;131:105295.
9. Milaneschi Y, Kappelmann N, Ye Z, Lamers F, Moser S, Jones PB, et al. Association of inflammation with depression and anxiety: evidence for symptom-specificity and potential causality from UK Biobank and NESDA cohorts. *Mol Psychiatry*. 2021.
10. Russell A, Hepgul N, Nikkheslat N, Borsini A, Zajkowska Z, Moll N, et al. Persistent fatigue induced by interferon-alpha: a novel, inflammation-based, proxy model of chronic fatigue syndrome. *Psychoneuroendocrinology*. 2019;100:276-85.
11. The COVID-19 Host Genetics Initiative, a global initiative to elucidate the role of host genetic factors in susceptibility and severity of the SARS-CoV-2 virus pandemic. *Eur J Hum Genet*. 2020;28(6):715-8.
12. Franceschi C, Campisi J. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J Gerontol A Biol Sci Med Sci*. 2014;69 Suppl 1:S4-9.
13. Bartlett DB, Firth CM, Phillips AC, Moss P, Baylis D, Syddall H, et al. The age-related increase in low-grade systemic inflammation (Inflammaging) is not driven by cytomegalovirus infection. *Aging Cell*. 2012;11(5):912-5.
14. Andrew T, Hart DJ, Snieder H, de Lange M, Spector TD, MacGregor AJ. Are twins and singletons comparable? A study of disease-related and lifestyle characteristics in adult women. *Twin Res*. 2001;4(6):464-77.
15. Verdi S, Abbasian G, Bowyer RCE, Lachance G, Yarand D, Christofidou P, et al. TwinsUK: The UK Adult Twin Registry Update. *Twin Res Hum Genet*. 2019;22(6):523-9.
16. Suthahar A, Sharma P, Hart D, García MP, Horsfall R, Bowyer RCE, et al. TwinsUK COVID-19 personal experience questionnaire (CoPE): wave 1 data capture April-May 2020. *Wellcome Open Research*. 2020;6(123).
17. Muench P, Jochum S, Wenderoth V, Ofenloch-Haehnle B, Hombach M, Strobl M, et al. Development and Validation of the Elecsys Anti-SARS-CoV-2 Immunoassay as a Highly Specific Tool for Determining Past Exposure to SARS-CoV-2. *Journal of Clinical Microbiology*. 2020;58(10):e01694-20.
18. Seow J, Graham C, Merrick B, Acors S, Pickering S, Steel KJA, et al. Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. *Nature Microbiology*. 2020;5(12):1598-607.
19. Jackson C. The Chalder Fatigue Scale (CFQ 11). *Occup Med (Lond)*. 2015;65(1):86.
20. Cella M, Chalder T. Measuring fatigue in clinical and community settings. *J Psychosom Res*. 2010;69(1):17-22.

21. Cho HJ, Kivimaki M, Bower JE, Irwin MR. Association of C-reactive protein and interleukin-6 with new-onset fatigue in the Whitehall II prospective cohort study. *Psychol Med*. 2013;43(8):1773-83.
22. Hickie I, Davenport T, Wakefield D, Vollmer-Conna U, Cameron B, Vernon SD, et al. Post-infective and chronic fatigue syndromes precipitated by viral and non-viral pathogens: prospective cohort study. *BMJ*. 2006;333(7568):575.
23. Lim W, Hong S, Nelesen R, Dimsdale JE. The association of obesity, cytokine levels, and depressive symptoms with diverse measures of fatigue in healthy subjects. *Arch Intern Med*. 2005;165(8):910-5.
24. Fried SK, Bunkin DA, Greenberg AS. Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *J Clin Endocrinol Metab*. 1998;83(3):847-50.
25. Collin SM, Nikolaus S, Heron J, Knoop H, White PD, Crawley E. Chronic fatigue syndrome (CFS) symptom-based phenotypes in two clinical cohorts of adult patients in the UK and The Netherlands. *J Psychosom Res*. 2016;81:14-23.

Figure legends

Figure 1 – Participant flow diagram. CoPE COVID-19 Personal Experience Questionnaire CFQ Chadler Fatigue Scale.

Figure 2 – Association between fatigue and pre-pandemic levels of IL-6 and current levels of BMI. First panel shows unadjusted IL6 values, second panel shows values undusted for age, sex, BMI, and COVID-19 status (via residuals). IL-6 values or the residuals have been transformed to achieve normal distribution.

Figure 3 – Prevalence of different BMI classes among COVID-19 positive participants with and without chronic fatigue. BMI groups have been defined as the following: underweight (BMI<18.5; n = 6), normal weight (BMI >18.5 and < 25; n = 115), overweight (BMI >25 and <30; n = 102), and obesity (BMI ≥30; n = 59). The difference in the prevalence of the BMI groups is statistically significant ($\chi^2 = 8.3$, df = 3, p-value = 0.040).

Table 1 – Characteristics of the TwinsUK sample

Group	Sample size	Number with chronic fatigue (%)	Age (SD), years	BMI (SD), kg/m²
COVID-19 cases	282	49 (17.4)	65.4 (10.2)	26.5 (5.3)
COVID-19 controls	992	113 (11.4)	67.4 (9.6)	26.2 (4.8)
p-value		0.011	0.0004	0.356

Table 1 legend: P-value provided for comparisons between COVID-19 cases and controls using Fisher's exact test or Student's t-test. SD, standard deviation.

Table 2 – Comparison of Chalder Fatigue Scale responses in COVID-19 cases and controls

Chalder Fatigue Scale Question	COVID-19 cases (n=49)	COVID-19 controls (n=113)	p-value
Do you have problems with tiredness?	0.68±0.05	0.74±0.03	0.533
Do you need to rest more?	0.70±0.05	0.69±0.03	0.999
Do you feel sleepy or drowsy?	0.60±0.05	0.59±0.03	0.272
Do you have problems starting things?	0.64±0.05	0.70±0.03	0.199
Do you lack energy?	0.82±0.04	0.77±0.03	0.904
Do you have less strength in your muscles?	0.60±0.05	0.56±0.03	0.542
Do you feel weak?	0.48±0.05	0.49±0.03	0.381
Do you have difficulties concentrating?	0.72±0.05	0.65±0.03	0.299
Do you make slips of the tongue when speaking?	0.63±0.05	0.57±0.03	0.902
Do you find it more difficult to find the right word?	0.70±0.05	0.71±0.03	0.896
How is your memory?	0.65±0.05	0.61±0.03	0.345

Table 2 legend: Prevalence (\pm standard error) of positive answers among those who was classified as having long-term fatigue is provided; P-values estimated using Fisher's exact test.

Table 3 – Risk factors for chronic fatigue in TwinsUK participants by COVID-19 antibody status

Group	Variable	Estimate	Std. Error	z value	p value
COVID-19 positive	(Intercept)	-1.679	1.356	-1.238	0.216
	IL-6	-0.084	0.166	-0.509	0.611
	Age	-0.023	0.017	-1.389	0.165
	Sex	-1.175	1.051	-1.118	0.263
	BMI	0.063	0.030	2.146	0.032
COVID-19 negative	(Intercept)	-1.402	0.920	-1.524	0.127
	IL-6	0.304	0.107	2.849	0.004
	Age	-0.020	0.011	-1.841	0.066
	Sex	-0.144	0.510	-0.282	0.778
	BMI	0.025	0.021	1.170	0.242

Table 3 legend: Modelling pre-pandemic plasma IL-6 and other factors on the risk of chronic fatigue. IL-6 levels were adjusted for age and BMI followed by transformation to a normal distribution using *qqnorm* function in R. Model was fitted using age and BMI at the time CoPE questionnaire have been administered. Further adjustment was made for family structure and zygosity as random factors.