



## King's Research Portal

DOI:

[10.1007/s00127-016-1245-0](https://doi.org/10.1007/s00127-016-1245-0)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Valmaggia, L. R., Day, F., & Rus-Calafell, M. (2016). Using virtual reality to investigate psychological processes and mechanisms associated with the onset and maintenance of psychosis: a systematic review. *Social Psychiatry and Psychiatric Epidemiology*, 51(7), 921–936. <https://doi.org/10.1007/s00127-016-1245-0>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Using virtual reality to investigate psychological processes and mechanisms associated with the onset and maintenance of psychosis: a systematic review.**

Lucia R. Valmaggia<sup>1,2\*</sup> Fern Day<sup>1,3</sup>, Mar Rus-Calafell<sup>1,2</sup>

1. King's College London, Institute of Psychiatry, Psychology and Neuroscience, United Kingdom
2. South London and Maudsley NHS Trust, United Kingdom
3. City University London, United Kingdom

\*Corresponding author:

Dr Lucia R. Valmaggia, King's College London, Institute of Psychiatry, Psychology and Neuroscience, Department of Psychology (PO 77), De Crespigny Park, SE5 8AF London, United Kingdom

E-mail: [Lucia.Valmaggia@kcl.ac.uk](mailto:Lucia.Valmaggia@kcl.ac.uk)

Total words abstract: 185

Total words (excluding abstract, table, and references): 3174

## **Abstract**

**Purpose** In the last decade researchers have embraced virtual reality to explore the psychological processes and mechanisms that are involved in the onset and maintenance of psychosis. A systematic review was conducted to synthesise the evidence of using virtual reality to investigate these mechanisms.

**Methods** Web of Science, PsycINFO, Embase, and Medline. Reference lists of collected papers were also visually inspected to locate any relevant cited journal articles. A total of 6,001 articles potentially eligible for inclusion; of these, 16 studies were included in the review.

**Results** The review identified studies investigating the effect of interpersonal sensitivity, childhood bullying victimisation, physical assault, perceived ethnic discrimination, social defeat, population density and ethnic density on the real-time appraisal of VR social situations. Further studies demonstrated the potential of VR to investigate paranoid ideation, anomalous experiences, self-confidence, self-comparison, physiological activation and behavioural response.

**Conclusions** The reviewed studies suggest that VR can be used to investigate psychological processes and mechanisms associated with psychosis. Implications for further experimental research, as well as for assessment and clinical practice are discussed. The present review has been registered in the PROSPERO register: CRD42016038085.

**Keywords:** Virtual reality; adverse life events; daily stressors; stress sensitivity, psychosis.

## **Introduction**

Cognitive models of psychosis have investigated the psychological processes and mechanisms that play a role in psychosis and suggest that stressful life events can influence the way we perceive ourselves, others, and the world around us and can contribute to the onset and maintenance of psychosis [1-5]. These models have been supported by research which suggest that stress is a key feature in the onset and maintenance of mental health symptoms and disorders [6,7]. Stress can be defined as a physiological reaction to a threat that involves the activation of the hypothalamic-pituitary-adrenal (HPA) axis [8-10]. This reaction is similar across all mammalian species. In humans the psychological conceptualisation of stress proposes that the emotional response and physiological activation that occur in a situation are dependent on the interpretation of a threat and on whether we think we will be able to cope with the event or not [11]. Stress is a dominant feature of many aetiological models of psychosis [12,9,13], however stress is an elusive concept that is defined differently in different contexts. Reported stressful events can range from mild feelings of everyday stress and minor hassles (for example, living in a densely populated city; travelling in crowded public transport), through significant adverse life events (for example, bully victimization; ethnic discrimination), to traumatic events (for example, physical assault; sexual abuse).

A number of social factors that can be viewed as stressors have themselves been linked to an increased risk of psychosis. Urbanicity has been associated with higher levels of stress [14] and higher rates of psychosis [15,16]. Higher psychosis rates have also been found among migrants [17,18]. Higher incidence rates of psychosis have been linked to the perception of ethnic density [19,20], and associations between perceived ethnic discrimination and psychosis in ethnic minority and immigrant groups have been reported in previous research

[21-23]. Childhood bully-victimisation has also been associated with an increase in psychosis and other mental health problems in adulthood [24,25].

Also worth considering in this context is the evidence showing that individuals with schizophrenia have an increased sensitivity to daily hassles and report greater subjective stress [26-28]. These findings have stimulated interest in stress reactivity as an element of vulnerability [29] and an interest in how individuals with different levels of vulnerability for psychosis respond to daily stressors [30-32]. Interpersonal sensitivity is described as “*an undue and excessive awareness of, and sensitivity to, the behaviour and feelings of others... particularly to perceived or actual situations of criticism or rejection...*” (p.342 [33]). Higher interpersonal sensitivity is linked to problems in interpersonal relationships, low self-esteem, as well as negative self-concepts, and it has been found to be higher in individuals at ultra high risk for psychosis [32,34]. Previous studies have also found that a negative self-concept [35,36]) and low or unstable self-esteem [35] are linked to higher levels of paranoia in clinical and non-clinical samples.

When thinking about possible underlying mechanisms in the onset of psychosis is important to consider that prolonged exposure to social adversity and social exclusion have been linked to high prevalence rates of anomalous experiences (such as a propensity to hallucinations, high endorsement of psychic items or higher trait paranoia) in the general population and have been found to be associated with the same risk factors present in people with psychosis [37-41]. Stressful life events have also been linked to paranoid ideation [42]. Paranoid ideation has been defined as “*the unfounded fear that others intend to cause you harm*” [43,44], it is the most common type of delusional belief recorded by individuals with psychosis [45] and it is a common delusional belief in the general population [46].

While there is a long tradition of research in these processes and mechanisms they are usually assessed with retrospective self-report questionnaires or interviews. Limited evidence is

available researching underlying psychological processes and mechanisms in real-life or in ecologically valid environments [47,48]. In the last decade, researchers in the field of psychosis have embraced Immersive Virtual Reality (hereafter shortened to VR) as a method to explore the psychological mechanisms associated with the onset and maintenance of psychosis. VR environments are displayed in colour and in 3D using a head-mounted-display. The user is fully immersed in the environment and can interact with the environment either by head movements, full body turning or with a joystick. The sounds of the environment are heard through headphones [49] (please see <https://www.youtube.com/watch?v=svG6kXC2PQA> for an example). VR environments elicit sense of presence [50] and trigger real time cognitive, emotional, behavioural, and physiological responses to real time situations [51]. This sense of ‘being there’ as if it were a real life situation has led researcher to describe VR environments as ecologically valid [51,52]. Aside from allowing to simulate “real life” social situation and bring them into a laboratory setting, the added value of VR is that it allows the researcher to measure real time responses, and at the same time it enables to control and manipulate each aspect of an environment [53]. VR enables thus researchers to expose participants to exactly the same social environment and to measure the individual cognitive, emotional, behavioural and physiological response to the environment. It also enables to investigate the association between an individual prior exposure to social and environmental factors (e.g. history of migration, or bully victimisation) and their real-time response to a social environment. The VR environment itself can also be manipulated, for example, the number of virtual characters in an environment can be changed, as can be their ethnicity, their facial expression, their behaviour or what they say. Sounds, like for example indistinct background conversations, crowd murmuring or laughter or other noises like background music or sound of traffic can also be manipulated.

The aim of this manuscript was to review experimental studies that used VR to investigate the underlying psychological processes and mechanisms associated with the onset and maintenance of psychosis.

## **Method**

### **Design**

We conducted a systematic review of the literature and described the results in a qualitative synthesis. The review was registered on the PROSPERO register CRD42016038085.

### **Selection Procedure**

The databases used to conduct the search were Web of Science, PsycINFO, Embase, and Medline. Reference lists of collected papers were also visually inspected to locate any relevant cited journal articles. Studies were included in the review if they were: written in English; contained original empirical findings, and published in a peer-reviewed journal. Studies were excluded from the review if they were treatment studies; case studies; reviews; not available in English; theses; or book chapters. Only studies of immersive VR were included.

### **Search Criteria**

Studies for review were identified following a keyword search for the terms ‘virtual reality’ OR ‘VR’ in conjunction with ‘social stress’ OR ‘adverse life events’ OR ‘urban’, OR ‘migration’, OR ‘bullying’, OR ‘discrimination’, OR ‘social defeat’, OR ‘social entrapment’,

OR 'stress', AND 'psychological processes', OR 'mechanisms', OR 'mental health', OR 'paranoia', OR 'anxiety', OR 'depression', OR 'trauma', OR 'psychosis' OR 'schizophrenia'. Appropriate truncations and wild cards were used to identify permutation of the terms searched, e.g. stress\* to search for: stress, stressors, stressful; paranoi\* to search for: paranoia, paranoid.

### **Quality Assessment**

Quality of the reviewed studies was assessed independently by LV and MRC using the Evaluation of Public Health Practice Project (EPHPP) Quality Assessment Tool for Quantitative Studies [54]. A global rating for the paper is described as follows: Strong=no weak ratings; Moderate=one weak rating; Weak= two or more weak ratings on the subscales (selection bias; study design; confounders; blinding; data collection methods; withdrawals and dropouts) The EPHPP is available online (<http://ca/tools.html>). Rating disagreements were resolved by consensus.

### **Results**

#### **Information Extraction**

The search resulted in 5,999 articles (final search conducted 4<sup>th</sup> March 2016); two more manuscripts were identified by publication alerts email from journals. From these 6,001 studies we identified 16 experimental studies which used full immersive VR to research adverse life events, daily stressors and psychological processes associated with psychosis (see Figure 1).



-- Figure 1 --

## **Qualitative Synthesis**

The reviewed studies were organised in two categories. The first category describes studies which used VR to investigate the association between baseline characteristics and the real-time response to a neutral social VR environment. The second category includes studies that manipulated one or more aspects of the environment to measure how the manipulation would influence the real-time response to VR environment. Finally, we review overarching findings from both categories regarding whether VR can be used to investigate real-time response to social situations.

### **Studies exploring the association between baseline characteristics and the real-time response to a neutral social VR environment.**

The ten studies summarised in Table 1 used VR to explore the association between existing characteristics of participants and their response to a VR social situation. In two studies the environment was a library, the remaining studies used an underground train environment. Both the library environment and the underground were designed to be perceived as neutral social environments. Virtual characters or avatars showed neutral behaviours typical of such social situations, but that could be interpreted as ambiguous like for example, looking at the participant, smiling, or avatars talking to each other. This was done on purpose as the main aim of the studies was to explore ‘unfounded’ real-time paranoid ideation in VR rather than a response to a genuinely hostile environment [55].

-- Table 1 ---

*Childhood Bullying Victimization, Physical Assault, Perceived Ethnic Discrimination, and Social Defeat*

Five studies investigated whether having experienced adverse life events was associated with higher levels of paranoid ideations in the neutral social VR environments. All five studies used the VR underground train environment. Childhood bullying victimisation [56]; being the victim of a physical assault [57,58]; perceived ethnic discrimination [59] and social defeat [60] were found to be associated with real-time paranoid ideation in the VR social environment.

*Interpersonal sensitivity*

The correlation between interpersonal sensitivity and real-time paranoid ideation during VR was investigated in five studies [61-65]. All studies found that heightened interpersonal sensitivity was associated with higher levels of paranoia in the VR library or in the underground train.

**Studies investigating the effect of a manipulation on the response to a VR environment.**

The six studies listed in Table 2 have used a manipulation to investigate the real-time response to the VR environment. The studies have used different environments: a bar, a student flat, and the underground train described earlier.

--Table 2 –

### *Population Density and Ethnic Density*

Two studies investigated how manipulating population density and ethnic density effected paranoid ideation in a VR bar environment [66,67]. The findings suggested that increased population density, hostility of the avatars, and a different ethnicity were associated with higher levels of paranoia in the VR bar.

### *Self-Confidence and Social Comparison*

Two experiments indicated that VR can be useful to measure self-confidence and social comparison. In the first study participants underwent a manipulation aimed to induce low or high self-confidence. In the high self-confidence induction they were instructed to think about a time in their life when they felt self-confident and they were provided with a list of confidence statements. In the low-confidence induction, participants underwent a similar procedure but regarding their least self-confident experience. Low self-confidence was associated with higher paranoid ideation in VR underground train [68].

The second experiment manipulated the height of participants in the VR environment and found that a lowered height was associated with negative social-comparison [69].

### *Physiological Activation and Behavioural response*

Manipulation of the VR environment triggered not only the cognitive and emotional response to a social situation, but also the physiological changes and behavioural response. Galvanic

skin response, heart rate and behavioural response (distance kept from the avatars) was found to be triggered by changes in population density and ethnic density in a VR bar environment [70]. The same environment was used to investigate galvanic skin response in a sample of individuals with psychosis [66]. Distance kept from the avatar was also measured in a study which manipulated the contingency behaviour of avatars while participants were exploring a VR student flat [61].

## **VR to investigate real time responses to social situations**

### *Paranoid Ideations*

As summarised in Tables 1 and 2, the large majority of studies investigated paranoid ideation and the associated correlates. VR was found to be a valid research method, not only to elicit and measure paranoid ideation in participants, but also to generate participant's attributions about positive, neutral and negative mental states to virtual characters as if they were real people [63,62,59,60,56,64,66,67]. Furthermore, as illustrated in Tables 1 and 2, VR was found to be effective in generating paranoid ideation in people from general population samples [61,57,69,63,71,62,58] as well as from individuals with an ultra high risk for psychosis [72,59,60,56,64,67], and those already experiencing a psychotic disorder [66,67].

### *Anomalous experiences*

Individuals experiencing anomalous experiences (such as a propensity to hallucinations, high endorsement of psychic items or higher trait paranoia) reported higher levels of paranoid ideation when exposed to a VR environment [61,63,71,65,64,67]. These results confirm the validity of VR to assess real-time paranoia in response to a social situation, as it would be

expected that an individual who reports trait paranoia would also experience paranoid ideation in an ecologically valid social VR environment.

### **Quality Assessment**

Several studies reached a strong rating on the EPHPP global rating (see Table 1 and 2, last column). No studies were excluded based on their quality. However, it is worth mentioning that a strong global rating on the EPHPP allows a moderate score on one of more subscales. Table 3 illustrates the subscale scores for each study: the majority of studies reach moderate scores on selection bias, indicating that the participants recruited for the study were not very likely to be representative of the target population. Indeed quite a few studies were conducted with healthy controls and student samples. The majority of studies had a within subject design, did not always controlled for confounders and the outcome assessor was not always blind to the experimental condition. Most studies had a sound data collection method and few reported withdrawals or drop-outs.

-- Table 3 --

### **Discussion**

We sought to review experimental studies that used VR to investigate the underlying psychological processes and mechanisms associated with the onset and maintenance of psychosis. The term stress was consider to encompass a broad range of events and experiences encountered in the psychosocial domain that provoke feelings of distress, mental

strain, or physiological symptoms of stress. These experiences included traumatic events, social adversity, stressful life events, and subjective feelings of stress and inability to cope. Childhood bullying victimisation [56]; being the victim of a physical assault [57,58]; perceived ethnic discrimination [59] and social defeat [60] were associated with paranoid ideation in a VR social environment. The ability to use VR to study the effect of adverse life events on real-time response to social situations in a controlled environment offers an exciting new way to research what contributes to stress sensitivity. Measuring the real-time physiological activation in social situations offers a number of possibilities not only for research purposes but also for assessment and treatment [10,48]. Interestingly, VR has been shown to be an effective way to overcome excessive self-criticism in people with depression [73,74], suggesting that VR can be not just a method to assess components of interpersonal sensitivity, but that it also has the potential to be used in the treatment of decrease interpersonal sensitivity.

It has generally been assumed that stress exposure leads to the development of psychosis and psychotic-like-experiences, however reverse causality is also possible. Symptoms, personality traits and unusual behaviour may predispose individuals to stressful experiences [75]: for example, early symptoms may predispose an individual to bullying victimisation [76,77,25]. VR could be embedded in longitudinal research to allow the repeated measurements of current stress response alongside repeated measure of life stressors and symptoms. This type of research could also help in defining resilience factors in people who experience adverse life events, but do not go on to develop psychosis [78,79]. Previous studies have recognised the similarity between anomalous experiences and intrusive symptoms of post-traumatic stress disorder (PTSD) [80] as well as co-morbid PTSD in people with psychosis (e.g. [81]). While not the topic of the current review, is also worth mentioning that the search revealed a rich body of literature demonstrating the use of VR-

assisted therapy for mental health problems (for recent reviews see [72,82,83]), and the use of VR in the research, assessment and treatment of psychosis [84,85] and PTSD [86-88,49] has previously been the topic of extensive reviews. Lessons can therefore be learned from those studies to inform future experimental studies into the psychological mechanisms of psychosis.

Further evidence for the fact that VR is an ecologically valid method to assess real-time responses to social environments was found by studies looking at the effect of manipulating population density and the ethnic density of avatars on the levels of paranoid ideation and physiological activation [70,66,67]. Of interest in the context of urbanicity is the finding that changes in the architectural characteristics of a room in VR can be used to change the physiological reaction to closed spaces: compared to participants in a VR room without windows, participants in the room with windows and a nice calming view had lower cortisol reactivity to stress and a quicker recovery after a stress induction test [89].

Using VR to investigate anomalous experiences opens up engaging avenues for further research in the association between paranoid ideation and the mechanisms which underlie the propensity to experience hallucinations, such as self-monitoring, attributional biases, and mental imagery (e.g.[90,91]. VR studies can also contribute to research the differences between clinical and non-clinical samples of people who report anomalous experiences [92,93].

## **Limitations**

There were limitations regarding the research conducted to date. The design of all studies was cross-sectional and life events were assessed retrospectively. A number of studies had small sample sizes and healthy control participants in some of the studies were exclusively from a student population, thus differences in social class, employment status, and level of

education between people with psychosis and student samples may contribute to some of the differences found between groups. A final possible limitation is also the fact that the reviewed studies were conducted by a relatively small group of researchers who collaborate with each other and who have a strong research interest in paranoid ideation. The focus of research on paranoid ideation reflects on one hand, the long tradition and large body of research of the role of psychological mechanisms in the aetiology of paranoid symptoms [94-97]. On the other hand, this strong focus on paranoid ideation also reflects the interest of the clinical academics who conducted the studies, with the large majority of reviewed studies conducted by Freeman and colleagues, or by researchers who collaborated with him. Future VR research should ensure that other areas of stress research are not neglected.

## **Conclusion**

The reviewed studies suggest that VR can be used to investigate psychological processes and mechanisms associated with psychosis. The ability to assess real-time, cognitive, emotional, behavioral and physiological responses in a controlled but ecologically valid environment has enormous potential for future research in the mechanisms involved in the onset and maintenance of psychosis.

## **Acknowledgments**

We acknowledge the NIHR Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and Institute of Psychiatry King's College London for their support.



### **Author Disclosure Statement**

The authors work in a VR lab and have published some of the studies reviewed in this review.



**Table 1.** Studies exploring the association between baseline characteristics of participants and the cognitive, emotional, behavioural, and physiological response to an immersive VR social environment to (n=10)

Study (Country)	Area studied	Design	Participants	Mean age (SD)	VR equipment	VR task	Measures	Main findings	EPHPP Global Rating
Freeman, Slater et al, 2003 [62]  (United Kingdom)	Persecutory ideation	Within subjects design	24 healthy participants (12 male and 12 females)	26 (6)	CAVE Immersive projection system and Crystal Eyes shutter-glasses.	Library with 5 avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	BSI; SSPS; STAI; VR paranoia questionnaire; Semi structured interview and observer rating of persecutory ideation; SUS	Individuals attributed mental states to avatars, including paranoid intentions.  Persecutory ideation was associated with interpersonal sensitivity	Moderate
Freeman, Garety, et al., 2005 [63]  (United Kingdom)	Persecutory ideation	Within subjects design	30 healthy participants (15 males and 15 females; 70% White ethnicity)	22 (5)	CAVE Immersive projection system and Crystal Eyes shutter-glasses	Library with 5 avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	SSPS; LSHS; Structured Interview for assessing perceptual abnormalities; Need for closure: DASS-21; IPSM; PSCS; Probabilistic reasoning task ; SADS; VR paranoia questionnaire; VR-Social avoidance and distress scale; SUS	Individuals attributed mental states to avatars, including paranoid intentions.  Persecutory ideation was predicted by baseline anxiety, timidity and hallucination predisposition  No association was found with probabilistic reasoning or need for closure	Moderate
Freeman, Gittins et al, 2008 [71]  (United Kingdom)	Paranoid ideation and social anxiety	Within subjects design	200 participants from general population (100 males and 100 females; 63% White ethnicity; 47% graduate degree or higher ;17% unemployed)	37.5 (13.3)	HMD: VR1280	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	WAIS; DASS-21; PSWQ; Worry domains questionnaire; Cathastrophizing interview; BCSS; IPMS; Cognitive flexibility; Probabilistic reasoning; CAPS; MAP; Life stressors checklist; SSQ; SELSA; SSPS; SADS	Higher baseline perceptual abnormalities was a predictor of paranoid ideation but not of social anxiety.	Strong
Freeman, Pugh et al, 2010 [65]  (United Kingdom)	Persecutory ideation and associated factors	Within subjects design	Group 1 (n=30): low non-clinical paranoia (83% White ethnicity)  Group 2 (n=30): high non-clinical paranoia (77% White ethnicity)  Gropu 3(n=30) : Clinical	Group 1: 44.2 (11.2)  Group 2: 36 (11.7)	HMD: VR1280	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	GPTS; SSPS; VAS hostility; DASS-21; PSWQ; IPSM; Beads task; CAPS; Life stressors checklist; WAIS; SSQ Simulation sickness questionnaire	There was an increase in levels of anxiety, worry, interpersonal sensitivity, depression, anomalous experiences and trauma history across the three groups of	Strong

			Persecutory delusions (53% White ethnicity)  Each group had 18 males and 12 females	Group 3: 44.2 (11.7)				paranoia.  Jumping to conclusion was only present in the persecutory delusions group.	
Freeman, Thompson et al, 2013 [58]  (United Kingdom)	Paranoia and PTSD following an assault	Within subject comparison	106 participants recruited in an A&E department after having experienced a distressing physical assault (79 males and 27 females ; 52% White ethnicity; 31% unemployed)	34.4 (11.6)	HMD: NVIS SX111	Participants experienced a underground train ride with avatar avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	Posttraumatic diagnostic scale; Structural Clinical Interview for DSM-IV; GPTS; PANSS; SSPS; Thoughts and feelings during the assault scale; CPQ; MDS; TMQ; IQQ; RIQ; SBQ; PCI; Perceived negative response of others; CSS; PSWQ; ISM; Catastrophizing Interview; BCSS; Cognitive flexibility scale; CAPS; MAP; VAS: PTSD; Paranoia	PTSD and Paranoia were correlated but distinct experiences	Strong
Freeman, Antley et al, 2014 [57]  (United Kingdom)	Paranoia in VR as predictor of paranoia and PTSD 6 months after experiencing an assault	Within subject comparison	As above	As above	As above	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	PANSS; GPTS; VAS: PTSD; Paranoia; PSRS; Posttraumatic diagnostic scale; SSPS	Response to the VR environment predicted the severity of PTSD and paranoia symptoms at 6 months.	Strong
Shaikh et al, 2016 [59]  (United Kingdom)	Perceived ethnic discrimination and paranoid ideation	Cross sectional between participants independent design	64 Ultra High Risk individuals (UHR) (38 males 26 females; 36% White British; 56% Unemployed; 13% Higher education)  43 Healthy Controls (20 males, 23 females; 37% White British; 7% Unemployed; 33% Higher education)	UHR 22.5 (4.0)  HC 24 (4.0)	HMD: VR1280	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	Prodromal questionnaire; SSPS; PEDQ-Community version	UHR reported higher levels of perceived ethnic discrimination  Higher levels of perceived ethnic discrimination were associated with greater paranoia in VR in the entire sample.	Strong
Valmaggia, Freeman et al,	Paranoid ideation	Within subject	21 Ultra High Risk individuals (UHR) (13	25 (4.7)	CAVE Immersive	Participants experienced a	GPTS; PS; ISM; DASS; LSHS; Paranoia Scale; Beads	Paranoid ideation in VR were predicted by	Strong

2007 [64]  (United Kingdom)		comparison	males and 8 females; 57% White ethnicity; 33% unemployed)		projection system and Crystal Eyes shutter-glasses.	underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	Task; Wisconsin Card Sorting Task; National Reading Test; Pre and Post VR VAS: anxiety; VR Questionnaire; Semi structured interview and observer rating of persecutory ideation  Follow-up assessment after one week to monitor intrusions and adverse effects in the week following the experiment	baseline paranoia; anxiety; stress, perseveration and interpersonal sensitivity  VR did not increase anxiety or cause negative experiences in the week following the experiment.	
Valmaggia, Day, Kroll et al, 2015 [56]  (United Kingdom)	Childhood bullying victimisation and paranoid ideation	Cross sectional between participants independent design	64 Ultra High Risk individuals (UHR) (38 males 26 females; 36% White British; 56% Unemployed; 13% Higher education  43 Healthy Controls (20 males, 23 females; 37% White British; 7% Unemployed; 33% Higher education)	UHR 22.5 (4.0)  HC 24 (4.0)	HMD: VR1280	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	Retrospective bullying questionnaire; SSPS	UHR were more likely to have experienced childhood bullying  Childhood bullying victimisation was associated with paranoid ideation in the VR independent of clinical status.	Strong
Valmaggia, Day, Garety et al, 2015 [60]  (United Kingdom)	Social defeat and paranoid ideation	Cross sectional between participants independent design	As above	As above	HMD: VR1280	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	Comprehensive Assessment of the at risk mental state; Prodromal questionnaire; SCS; Defeat Scale; Entrapment Scale; DASS; SSPS	UHR subjects reported significantly higher levels of social defeat than controls and their paranoid appraisals were predicted by level of social defeat at baseline	Strong
<p><i>NOTE:</i> HMD: Head Mounted Display; GPTS-B: Green Paranoia Thoughts Scale-B; VAS: Visual Analogical Scale; SCS: Social comparison scale; SSPS: Social State Paranoia Scale; GSR: Galvanic skin response; HR: Heart rate; STAI: Spielberg Trait Anxiety Inventory; SUS: Slater, Usoh, Steed Sense of Presence Scale; BSI: Brief Symptom Inventory; LSHS: Launay Slade Hallucination Scale; DASS: Depression Anxiety Stress Scale; IPSM: Interpersonal sensitivity Measure; PSCS-Private Self-Consciousness Scale; SADS: Social Avoidance and distress scale; WAIS: Wechsler Abbreviated scale for intelligence; PSWQ: Penn State Worry questionnaire; BCSS: Brief core schema scale; CAPS: Cardiff Anomalous perceptions scale; MAP: Maudsley Addiction Profile; SSQ: Social support questionnaire; SELSA: Social and emotional loneliness scale for adults; SCS: Social Comparison Scale; PANSS: Positive and Negative Symptom Scale; CPQ: Cognitive processing questionnaire; MDS: Mental defeat Scale; TMQ: Trauma Memory questionnaire; IQQ: Intrusions Qualities questionnaire; RIQ: Response to intrusion questionnaire; SBQ: Safety behaviour questionnaire; PCI: Posttraumatic cognition inventory; CSS: Crisis support scale; PSRS: Psychotic Symptoms rating Scale; PEDQ: Perceived Ethnic Discrimination Questionnaire- Community version; SIAS: Social Interaction Anxiety Scale; DACOBS: Davos Assessment of Cognitive Biases Scale; RSES: Rosenberg Self-esteem rating scale.</p>									



**Table 2.** Studies exploring the cognitive, emotional, behavioural, and physiological response to a manipulation in the VR social environment to (n=6)

Study (Country)	Area studied	Design	Participants	Mean age (SD)	VR equipment	VR task	Measures	Main findings	EPHPP Global Rating
Atherton et al, 2014 [68]  (United Kingdom)	Self-confidence and paranoid ideation	Within subjects design	26 males (from a general population sample who scored >17 on GPTS-Paranoia)	43.4 (16.3)	HMDs: NVIS SX111 or VR1280	Before VR participants received a low or a high self-confidence manipulation.  Participants experienced an underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)	GPTS-B; VAS social confidence; SCS; SSPS	Induced low self confidence led to higher levels of paranoia and more negative views of the self in the VR environment.	Strong
Brinkman et al, 2011 [70]  (The Netherlands)	Physiological response to changes in population density and ethnic density	Two by two within-subject design	2 Males (1 with schizoaffective disorder)  24 Healthy Controls (20 male and 4 female; all White Dutch)	29 (9.2)	HMD: Emagin Z800 3DVisor	Participants moved freely in a bar with different population density and avatar ethnic density.	GSR, HR, Subjective discomfort; Distance from avatar	Increased population density and ethnic density are associated with more fluctuation in physiological arousal in the VR environment	Weak
Freeman, Evans et al, 2013 [69]  (United Kingdom)	Persecutory ideation and social comparison after height was altered	Two condition, two period crossover trial	60 females from the general population (with paranoid thinking in the last month; 78% White ethnicity; 55% graduate degree or higher)	31.5 (13)	HMD: NVIS SX111	Participants experienced a underground train ride with avatars who occasionally showed potentially ambiguous behaviour (e.g. looking, smiling, talking)  The height of the participant was lowered during the experimental condition.	GPTS-B; SSPS; SCS	A lowered height was associated with higher levels of paranoid ideation and in more negative views about the self compared to others.	Strong
Fornells-Ambrojo et al, 2016 [61]  (United Kingdom)	Perception of trustworthiness and interpersonal distance	Between group comparison	61 healthy male (77%: White ethnicity; 90% students).	25.3 (7.3)	CAVE Immersive projection system and Crystal Eyes shutter-glasses..	Participants moved in a student flat and interacted with a virtual flatmate. The contingency behaviour of the flatmate was manipulated to be high in one condition	SSPS; STAI; Relationship Questionnaire; Distance kept from avatar; Trustworthiness of the avatar; SUS	High subjective trust was not associated with less distance from the avatar.  Higher trait paranoia was associated with heightened sensitivity to contingency	Moderate

						and low in the other.		behaviour.	
Veling, Brinkman et al, 2014 [66]  (The Netherlands)	Paranoid thoughts and associated social anxiety, cognitive bias, self esteem	Cross sectional between participants independent design	17 patients with a first episode of psychosis (FEP) (14 males, 3 females; 35% White Dutch ethnicity)  24 healthy controls (20 males, 4 females; 100% White Dutch ethnicity)	FEP 27.3 (5.5)  HC 29 (9.2)	Emagin Z800 3D Visor	Participants moved freely in a bar with different population density and avatar ethnic density.	GPTS; SIAS; DACOBS; RSES; Simulation sickness questionnaire; SSPS; Galvanic skin response	Patients reported more paranoid thoughts, showed more proximity to the avatars and higher galvanic skin response to avatars of a different ethnicity from their own.	Strong
Veling, Pot-Kolder et al, 2016 [67]  (The Netherlands)	Paranoid thoughts and stress sensitivity	Cross sectional between participants independent design	Group 1: 20 UHR (7 males, 13 females 75% White Dutch ethnicity)  Group 2: 55 Individuals with psychosis (42 males, 13 females; 53% White Dutch ethnicity)  Group 3: 42 siblings of people with psychosis (23 males, 19 females; 75% White Dutch ethnicity)  Group 4: 53 controls with negative first degree family history of psychosis (25 males, 28 females; 70% White Dutch ethnicity)	UHR 24 (4.5)  Psychosis 26 (4.7)  Siblings 26.4 (4.8)  Controls 24.6 (4.4)	Sony HMZ-T1	Participants moved freely in a bar with different population density, avatar ethnic density and hostility of facial expression of the avatars.	GPTS; SIAS; Community Assessment of psychic experiences; VAS: subjective distress; SSPS	Increase social stress was associated with increase in paranoia and subjective distress in VR.  Negative affect and psychosis liability were associated with higher levels of paranoia in VR	Strong
<p><i>NOTE:</i> HMD: Head Mounted Display; GPTS-B: Green Paranoia Thoughts Scale-B; VAS: Visual Analogical Scale; SCS: Social comparison scale; SSPS: Social State Paranoia Scale; GSR: Galvanic skin response; HR: Heart rate; STAI: Spielberg Trait Anxiety Inventory; SUS: Slater, Usoh, Steed Sense of Presence Scale; BSI: Brief Symptom Inventory; LSHS: Launay Slade Hallucination Scale; DASS: Depression Anxiety Stress Scale; IPSM: Interpersonal sensitivity Measure; PSCS-Private Self-Consciousness Scale; SADS: Social Avoidance and distress scale; WAIS: Wechsler Abbreviated scale for intelligence; PSWQ: Penn State Worry questionnaire; BCSS: Brief core schema scale; CAPS: Cardiff Anomalous perceptions scale; MAP: Maudsley Addiction Profile; SSQ: Social support questionnaire; SELSA: Social and emotional loneliness scale for adults; SCS; Social Comparison Scale; PANSS; Positive and Negative Symptom Scale; CPQ: Cognitive processing questionnaire; MDS: Mental defeat Scale; TMQ: Trauma Memory questionnaire; IQQ: Intrusions Qualities questionnaire; RIQ: Response to intrusion questionnaire; SBQ: Safety behaviour questionnaire; PCI: Posttraumatic cognition inventory; CSS: Crisis support scale; PSRS: Psychotic Symptoms rating Scale; PEDQ: Perceived Ethnic Discrimination Questionnaire- Community version; SIAS: Social Interaction Anxiety Scale; DACOBS: Davos Assessment of Cognitive Biases Scale; RSES: Rosenberg Self-esteem rating scale.</p>									



**Table 3. Immersive VR studies EPHPP quality assessment rating**

<b>Study</b>	<b>A. Selection bias</b>	<b>B. Study design</b>	<b>C. Confounders</b>	<b>D. Blinding</b>	<b>E. Data collection method</b>	<b>F. Withdrawals and drop out</b>	<b>Global rating</b>
Atherton et al, 2014 [68]	Moderate	Moderate	Strong	Moderate	Strong	Strong	Strong
Brinkman et al, 2011 [70]	Weak	Moderate	Weak	Weak	Strong	Strong	Weak
Fornells-Ambrojo et al, 2016 [61]	Weak	Strong	Moderate	Moderate	Strong	Strong	Moderate
Freeman, Slater et al, 2003 [62]	Moderate	Moderate	Weak	Moderate	Moderate	Strong	Moderate
Freeman, Garety, et al., 2005 [63]	Strong	Moderate	Weak	Moderate	Strong	Strong	Moderate
Freeman, Gittins et al, 2008 [71]	Moderate	Moderate	Strong	Moderate	Strong	Strong	Strong
Freeman, Pugh et al, 2010 [65]	Strong	Moderate	Strong	Moderate	Strong	Strong	Strong
Freeman, Evans et al, 2013 [69]	Strong	Moderate	Strong	Moderate	Strong	Strong	Strong
Freeman, Antley et al, 2014 [57]	Strong	Moderate	Strong	Moderate	Strong	Strong	Strong

Freeman, Thompson et al, 2013 [58]	Moderate	Moderate	Strong	Moderate	Strong	Strong	Strong
Shaikh et al, 2016 [59]	Moderate	Moderate	Strong	Moderate	Strong	Moderate	Strong
Valmaggia , Freeman et al, 2007 [64]	Moderate	Moderate	Moderate	Moderate	Strong	Moderate	Strong
Valmaggia, Day, Kroll et al, 2015 [56]	Moderate	Moderate	Moderate	Moderate	Strong	Moderate	Strong
Valmaggia, Day, Garety et al, 2015 [60]	Moderate	Moderate	Strong	Moderate	Strong	Moderate	Strong
Veling, Brinkman et al, 2014 [66]	Moderate	Moderate	Moderate	Moderate	Moderate	Moderate	Strong
Veling, Pot-Kolder et al, 2016 [67]	Moderate	Moderate	Moderate	Moderate	Moderate	Moderate	Strong

## References

1. Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington PE (2001) A cognitive model of the positive symptoms of psychosis. *Psychological medicine* 31 (2):189-195
2. Garety PA, Bebbington P, Fowler D, Freeman D, Kuipers E (2007) Implications for neurobiological research of cognitive models of psychosis: a theoretical paper. *Psychological medicine* 37 (10):1377-1391. doi:10.1017/S003329170700013X
3. Freeman D, Garety PA, Kuipers E, Fowler D, Bebbington PE (2002) A cognitive model of persecutory delusions. *The British journal of clinical psychology / the British Psychological Society* 41 (Pt 4):331-347
4. Bentall RP, Fernyhough C, Morrison AP, Lewis S, Corcoran R (2007) Prospects for a cognitive-developmental account of psychotic experiences. *The British journal of clinical psychology / the British Psychological Society* 46 (Pt 2):155-173. doi:10.1348/014466506X123011
5. Morrison AP, Wells A (2003) A comparison of metacognitions in patients with hallucinations, delusions, panic disorder, and non-patient controls. *Behaviour research and therapy* 41 (2):251-256
6. Van Dam DS, Van Nierop M, Viechtbauer W, Velthorst E, Van Winkel R, Bruggeman R, Cahn W, De Haan L, Kahn RS, Meijer CJ, Myin-Germeys I, Van Os J, Wiersma D (2015) Childhood abuse and neglect in relation to the presence and persistence of psychotic and depressive symptomatology. *Psychological Medicine* 45 (7):1363-1377. doi:10.1017/S0033291714001561
7. Pariante CM (2015) Psychoneuroimmunology or immunopsychiatry? *The Lancet Psychiatry* 2 (3):197-199. doi:10.1016/S2215-0366(15)00042-5
8. Aiello G, Horowitz M, Hepgul N, Pariante CM, Mondelli V (2012) Stress abnormalities in individuals at risk for psychosis: A review of studies in subjects with familial risk or with "at risk" mental state. *Psychoneuroendocrinology* 37 (10):1600-1613. doi:<http://dx.doi.org/10.1016/j.psyneuen.2012.05.003>
9. Walker E, Mittal V, Tessner K (2008) Stress and the hypothalamic pituitary adrenal axis in the developmental course of schizophrenia. *Annu Rev Clin Psychol* 4:189-216. doi:10.1146/annurev.clinpsy.4.022007.141248
10. Day FL, Valmaggia LR, Mondelli V, Papadopoulos A, Papadopoulos I, Pariante CM, McGuire P (2014) Blunted cortisol awakening response in people at ultra high risk of developing psychosis. *Schizophr Res* 158 (1-3):25-31. doi:10.1016/j.schres.2014.06.041
11. Lazarus RS (1991) Progress on a cognitive-motivational-relational theory of emotion. *Am Psychol* 46 (8):819-834
12. Corcoran C, Walker E, Huot R, Mittal V, Tessner K, Kestler L, Malaspina D (2003) The stress cascade and schizophrenia: etiology and onset. *Schizophr Bull* 29 (4):671-692
13. Zubin J, Spring B (1977) Vulnerability--a new view of schizophrenia. *J Abnorm Psychol* 86 (2):103-126
14. Lederbogen F, Kirsch P, Haddad L, Streit F, Tost H, Schuch P, Wust S, Pruessner JC, Rietschel M, Deuschle M, Meyer-Lindenberg A (2011) City living and urban upbringing affect neural social stress processing in humans. *Nature* 474 (7352):498-501. doi:<http://www.nature.com/nature/journal/v474/n7352/abs/nature10190-f1.2.html - supplementary-information>
15. Allardyce J, Gilmour H, Atkinson J, Rapson T, Bishop J, RG. M (2005) Social fragmentation, deprivation and urbanicity: relation to first-admission rates for psychoses. *British Journal of Psychiatry* 187:401-4016
16. Vassos E, Pedersen CB, Murray RM, Collier DA, Lewis CM (2012) Meta-analysis of the association of urbanicity with schizophrenia. *Schizophr Bull* 38 (6):1118-1123. doi:10.1093/schbul/sbs096
17. Morgan C, Hutchinson G (2009) The social determinants of psychosis in migrant and ethnic minority populations: a public health tragedy. *Psychol Med*:1-5. doi:10.1017/S0033291709005546
18. Seeman MV (2011) Canada: psychosis in the immigrant Caribbean population. *Int J Soc Psychiatry* 57 (5):462-470. doi:10.1177/0020764010365979

19. Boydell J, van Os J, McKenzie K, Allardyce J, Goel R, McCreadie RG, Murray RM (2001) Incidence of schizophrenia in ethnic minorities in London: ecological study into interactions with environment. *BMJ* 323 (7325):1336-1338
20. Das-Munshi J, Becares L, Boydell JE, Dewey ME, Morgan C, Stansfeld SA, Prince MJ (2012) Ethnic density as a buffer for psychotic experiences: findings from a national survey (EMPIRIC). *Br J Psychiatry* 201 (4):282-290. doi:10.1192/bjp.bp.111.102376
21. Karlsen S, Nazroo JY, McKenzie K, Bhui K, Weich S (2005) Racism, psychosis and common mental disorder among ethnic minority groups in England. *Psychol Med* 35 (12):1795-1803. doi:10.1017/S0033291705005830
22. Veling W, Selten JP, Susser E, Laan W, Mackenbach JP, Hoek HW (2007) Discrimination and the incidence of psychotic disorders among ethnic minorities in The Netherlands. *Int J Epidemiol* 36 (4):761-768. doi:10.1093/ije/dym085
23. Stowkowy J, Liu L, Cadenhead KS, Cannon TD, Cornblatt BA, McGlashan TH, Perkins DO, Seidman LJ, Tsuang MT, Walker EF, Woods SW, Bearden CE, Mathalon DH, Addington J (2016) Early traumatic experiences, perceived discrimination and conversion to psychosis in those at clinical high risk for psychosis. *Social Psychiatry and Psychiatric Epidemiology*:1-7. doi:10.1007/s00127-016-1182-y
24. Lereya ST, Copeland WE, Costello EJ, Wolke D (2015) Adult mental health consequences of peer bullying and maltreatment in childhood: two cohorts in two countries. *Lancet Psychiatry* 2 (6):524-531. doi:10.1016/S2215-0366(15)00165-0
25. Arseneault L, Cannon M, Fisher HL, Polanczyk G, Moffitt TE, Caspi A (2011) Childhood trauma and children's emerging psychotic symptoms: A genetically sensitive longitudinal cohort study. *Am J Psychiatry* 168 (1):65-72. doi:10.1176/appi.ajp.2010.10040567
26. Lukoff D, Snyder K, Ventura J, Nuechterlein KH (1984) Life events, familial stress, and coping in the developmental course of schizophrenia. *Schizophrenia bulletin* 10 (2):258-292
27. Norman RMG, Malla AK (1993) Stressful life events and schizophrenia. II: Conceptual and methodological issues. *British Journal of Psychiatry* 162 (FEB.):166-174
28. Corcoran C, Walker E, Huot R, Mittal V, Tessner K, Kestler L, Malaspina D (2003) The Stress Cascade and Schizophrenia: Etiology and Onset. *Schizophrenia Bulletin* 29 (4):671-692
29. Myin-Germeys I, Krabbendam L, Delespaul PAEG, Van Os J (2003) Do life events have their effect on psychosis by influencing the emotional reactivity to daily life stress? *Psychological Medicine* 33 (2):327-333
30. Lee SY, Kim KR, Park JY, Park JS, Kim B, Kang JI, Lee E, An SK, Kwon JS (2011) Coping strategies and their relationship to psychopathologies in people at ultra high-risk for psychosis and with schizophrenia. *The Journal of nervous and mental disease* 199 (2):106-110. doi:10.1097/NMD.0b013e3182083b96
31. Jalbrzikowski M, Sugar CA, Zinberg J, Bachman P, Cannon TD, Bearden CE (2014) Coping styles of individuals at clinical high risk for developing psychosis. *Early intervention in psychiatry* 8 (1):68-76. doi:10.1111/eip.12005
32. Masillo A, Day F, Laing J, Howes O, Fusar-Poli P, Byrne M, Bhattacharyya S, Fiori Nastro P, Girardi P, McGuire PK, Valmaggia LR (2012) Interpersonal sensitivity in the at-risk mental state for psychosis. *Psychological medicine* 42 (9):1835-1845. doi:10.1017/S0033291711002996
33. Boyce P, Parker G (1989) Development of a scale to measure interpersonal sensitivity. . *Australasian Psychiatry* 23 (3):341-351
34. Masillo A, Valmaggia LR, Saba R, Brandizzi M, Lindau JF, Solfanelli A, Curto M, Narilli F, Telesforo L, Kotzalidis GD, Di Pietro D, D'Alema M, Girardi P, Fiori Nastro P (2016) Interpersonal sensitivity and functioning impairment in youth at ultra-high risk for psychosis. *Eur Child Adolesc Psychiatry* 25 (1):7-16. doi:10.1007/s00787-015-0692-6
35. Tiernan B, Tracey R, Shannon C (2014) Paranoia and self-concepts in psychosis: a systematic review of the literature. *Psychiatry research* 216 (3):303-313. doi:10.1016/j.psychres.2014.02.003

36. Lincoln TM, Mehl S, Ziegler M, Kesting ML, Exner C, Rief W (2010) Is fear of others linked to an uncertain sense of self? The relevance of self-worth, interpersonal self-concepts, and dysfunctional beliefs to paranoia. *Behav Ther* 41 (2):187-197. doi:10.1016/j.beth.2009.02.004
37. van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L (2009) A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol Med* 39 (2):179-195. doi:10.1017/S0033291708003814
38. Kelleher I, Devlin N, Wigman JT, Kehoe A, Murtagh A, Fitzpatrick C, Cannon M (2014) Psychotic experiences in a mental health clinic sample: implications for suicidality, multimorbidity and functioning. *Psychol Med* 44 (8):1615-1624. doi:10.1017/S0033291713002122
39. Kelleher I, Wigman JT, Harley M, O'Hanlon E, Coughlan H, Rawdon C, Murphy J, Power E, Higgins NM, Cannon M (2015) Psychotic experiences in the population: Association with functioning and mental distress. *Schizophr Res* 165 (1):9-14. doi:10.1016/j.schres.2015.03.020
40. Wigman JT, Devlin N, Kelleher I, Murtagh A, Harley M, Kehoe A, Fitzpatrick C, Cannon M (2014) Psychotic symptoms, functioning and coping in adolescents with mental illness. *BMC Psychiatry* 14:97. doi:10.1186/1471-244X-14-97
41. Fusar-Poli P, Borgwardt S, Bechdolf A, Addington J, Riecher-Rossler A, Schultze-Lutter F, Keshavan M, Wood S, Ruhrmann S, Seidman LJ, Valmaggia L, Cannon T, Velthorst E, De Haan L, Cornblatt B, Bonoldi I, Birchwood M, McGlashan T, Carpenter W, McGorry P, Klosterkötter J, McGuire P, Yung A (2013) The psychosis high-risk state: a comprehensive state-of-the-art review. *JAMA Psychiatry* 70 (1):107-120. doi:10.1001/jamapsychiatry.2013.269
42. Bentall RP, de Sousa P, Varese F, Wickham S, Sitko K, Haarmans M, Read J (2014) From adversity to psychosis: pathways and mechanisms from specific adversities to specific symptoms. *Soc Psychiatry Psychiatr Epidemiol* 49 (7):1011-1022. doi:10.1007/s00127-014-0914-0
43. Freeman D, Garety PA (2000) Comments on the content of persecutory delusions: does the definition need clarification? *The British journal of clinical psychology / the British Psychological Society* 39 ( Pt 4):407-414
44. Freeman D, Pugh K, Antley A, Slater M, Bebbington P, Gittins M, Dunn G, Kuipers E, Fowler D, Garety P (2008) Virtual reality study of paranoid thinking in the general population. *The British journal of psychiatry : the journal of mental science* 192 (4):258-263. doi:10.1192/bjp.bp.107.044677
45. Garety PA, Everitt BS, Hemsley DR (1988) The characteristics of delusions: a cluster analysis of deluded subjects. *Eur Arch Psychiatry Neurol Sci* 237 (2):112-114
46. Freeman D (2007) Suspicious minds: the psychology of persecutory delusions. *Clin Psychol Rev* 27 (4):425-457. doi:10.1016/j.cpr.2006.10.004
47. Myin-Germeys I, Oorschot M, Collip D, Lataster J, Delespaul P, van Os J (2009) Experience sampling research in psychopathology: opening the black box of daily life. *Psychol Med* 39 (9):1533-1547. doi:10.1017/S0033291708004947
48. Reininghaus U, Depp CA, Myin-Germeys I (2016) Ecological Interventionist Causal Models in Psychosis: Targeting Psychological Mechanisms in Daily Life. *Schizophr Bull* 42 (2):264-269. doi:10.1093/schbul/sbv193
49. Rizzo A, Buckwalter G, Forbell E, Reist C, Difede J, Rothbaum BO, Ijzerman H, Koenig S, Talbot S (2013) Virtual reality Applications to Address the Wounds of War. *Psychiatric Annals* 43 (3). doi:10.3928/00485713-20130306-08
50. Slater M (2004) Presence and emotions. *Cyberpsychol Behav* 7 (1):121; author reply 123. doi:10.1089/109493104322820200
51. Parsons TD, Courtney CG, Arizmendi B, Dawson M (2011) Virtual Reality Stroop Task for neurocognitive assessment. *Stud Health Technol Inform* 163:433-439
52. Seidel RJ, Chatelier PR (2013) *Virtual Reality, Training's Future?: Perspectives on Virtual Reality and Related Emerging Technologies*, vol 6. Springer Science & Business Media,
53. Rizzo A, Kim GJ (2005) A SWOT analysis of the field of virtual reality rehabilitation and therapy. *Presence, Teleoperators and Virtual Environments* 14:119-146

54. Thomas BH, Ciliska D, Dobbins M, Micucci SA (2004) A Process for Systematically Reviewing the Literature: Providing the Research Evidence for Public Health Nursing Interventions. *Worldviews on Evid Based Nursing* 1 (3):176-184
55. Freeman D (2008) Studying and treating schizophrenia using virtual reality: a new paradigm. *Schizophrenia bulletin* 34 (4):605-610
56. Valmaggia LR, Day FL, Kroll J, Laing J, Byrne M, Fusar-Poli P, McGuire P (2015) Bullying victimisation and paranoid ideation in people at ultra high risk for psychosis. *Schizophrenia Research* 168 (1-2):68-73
57. Freeman D, Antley A, Ehlers A, Dunn G, Thompson C, Vorontsova N, Garety P, Kuipers E, Glucksman E, Slater M (2014) The use of immersive Virtual Reality (VR) to predict the occurrence 6 months later of paranoid thinking and posttraumatic stress symptoms assessed by self-report and interviewer methods: A study of individuals who have been physically assaulted. *Psychological Assessment* 26 (3):841-847
58. Freeman D, Thompson C, Vorontsova N, Dunn G, Carter LA, Garety P, Kuipers E, Slater M, Antley A, Glucksman E, Ehlers A (2013) Paranoia and post-traumatic stress disorder in the months after a physical assault: a longitudinal study examining shared and differential predictors. *Psychological medicine* 43 (12):2673-2684
59. Shaikh M, Ellett L, Dutt A, Day F, Laing K, Kroll J, Pterella S, McGuire P, Valmaggia L (2016) Perceived ethnic discrimination and persecutory paranoia in individuals at ultra-high risk for psychosis. *Psychiatry Research*
60. Valmaggia LR, Day F, Garety P, Freeman D, Antley A, Slater M, Swapp D, Myin-Germeys I, McGuire P (2015) Social defeat predicts paranoid appraisals in people at high risk for psychosis. *Schizophrenia Research* 168 (1-2):16-22
61. Fornells-Ambrojo M, Elenbaas M, Barker C, Swapp D, Navarro X, Rovira A, Sanahuja JMT, Slater M (2016) Hypersensitivity to contingent behavior in paranoia: A new virtual reality paradigm. *Journal of Nervous and Mental Disease* 204 (2):148-152
62. Freeman D, Slater M, Bebbington PE, Garety PA, Kuipers E, Fowler D, Met A, Read CM, Jordan J, Vinayagamoorthy V (2003) Can virtual reality be used to investigate persecutory ideation? *The Journal of nervous and mental disease* 191 (8):509-514
63. Freeman D, Garety PA, Bebbington P, Slater M, Kuipers E, Fowler D, Green C, Jordan J, Ray K, Dunn G (2005) The psychology of persecutory ideation II: A virtual reality experimental study. *Journal of Nervous and Mental Disease* 193 (5):309-315
64. Valmaggia LR, Freeman D, Green C, Garety P, Swapp D, Antley A, Prescott C, Fowler D, Kuipers E, Bebbington P, Slater M, Broome M, McGuire PK (2007) Virtual reality and paranoid ideations in people with an 'at-risk mental state' for psychosis. *British Journal of Psychiatry* 191 (SUPPL. 51):s63-s68
65. Freeman D, Pugh K, Vorontsova N, Antley A, Slater M (2010) Testing the Continuum of Delusional Beliefs: An Experimental Study Using Virtual Reality. *Journal of Abnormal Psychology* 119 (1):83-92
66. Veling W, Brinkman WP, Dorrestijn E, van der Gaag M (2014) Virtual reality experiments linking social environment and psychosis: a pilot study. *Cyberpsychology, behavior and social networking* 17 (3):191-195
67. Veling W, Pot-Kolder R, Counotte J, van Os J, van der Gaag M (2016) Environmental Social Stress, Paranoia and Psychosis Liability: A Virtual Reality Study. *Schizophr Bull.* doi:10.1093/schbul/sbw031
68. Atherton S, Antley A, Evans N, Cernis E, Lister R, Dunn G, Slater M, Freeman D (2016) Self-confidence and paranoia: An experimental study using an immersive virtual reality social situation. *Behavioural and Cognitive Psychotherapy* 44 (1):56-64
69. Freeman D, Evans N, Lister R, Antley A, Dunn G, Slater M (2014) Height, social comparison, and paranoia: An immersive virtual reality experimental study. *Psychiatry Research* 218 (3):348-352
70. Brinkman WP, Veling W, Dorrestijn E, Sandino G, Vakili V, van der Gaag M (2011) Virtual reality to study responses to social environmental stressors in individuals with and without psychosis. *Studies in health technology and informatics* 167:86-91

71. Freeman D, Gittins M, Pugh K, Antley A, Slater M, Dunn G (2008) What makes one person paranoid and another person anxious? The differential prediction of social anxiety and persecutory ideation in an experimental situation. *Psychological Medicine* 38 (8):1121-1132
72. Goncalves R, Pedrozo AL, Coutinho ESF, Figueira I, Ventura P (2012) Efficacy of Virtual Reality Exposure Therapy in the Treatment of PTSD: A Systematic Review. *PLoS ONE* 7 (12):no pagination
73. Falconer CJ, Slater M, Rovira A, King JA, Gilbert P, Antley A, Brewin CR (2014) Embodying compassion: A virtual reality paradigm for overcoming excessive self-criticism. *PLoS ONE* 9 (11):no pagination
74. Falconer CJ, Rovira A, King JA, Gilbert P, Antley A, Fearon P, Ralph N, Slater M, Brewin CR (2016) Embodying self-compassion within virtual reality and its effects on patients with depression. *British Journal of Psychiatry Open* 2 (1):74-80
75. Bebbington PE, Bhugra D, Brugha T, Singleton N, Farrell M, Jenkins R, Lewis G, Meltzer H (2004) Psychosis, victimisation and childhood disadvantage: evidence from the second British National Survey of Psychiatric Morbidity. *Br J Psychiatry* 185:220-226. doi:10.1192/bjp.185.3.220
76. Shakoor S, McGuire P, Cardno AG, Freeman D, Plomin R, Ronald A (2015) A shared genetic propensity underlies experiences of bullying victimization in late childhood and self-rated paranoid thinking in adolescence. *Schizophr Bull* 41 (3):754-763. doi:10.1093/schbul/sbu142
77. Goldman-Mellor S, Caspi A, Arseneault L, Ajala N, Ambler A, Danese A, Fisher H, Hucker A, Odgers C, Williams T, Wong C, Moffitt TE (2016) Committed to work but vulnerable: self-perceptions and mental health in NEET 18-year olds from a contemporary British cohort. *J Child Psychol Psychiatry* 57 (2):196-203. doi:10.1111/jcpp.12459
78. Brett CMC, Peters ER, McGuire PK (2015) Which psychotic experiences are associated with a need for clinical care? *European Psychiatry* 30 (5):648-654. doi:10.1016/j.eurpsy.2014.12.005
79. Underwood R, Kumari V, Peters E (2016) Appraisals of psychotic experiences: An experimental investigation of symptomatic, remitted and non-need-for-care individuals. *Psychological Medicine* 46 (6):1249-1263. doi:10.1017/S0033291715002780
80. Hardy A, Fowler D, Freeman D, Smith B, Steel C, Evans J, Garety P, Kuipers E, Bebbington P, Dunn G (2005) Trauma and hallucinatory experience in psychosis. *J Nerv Ment Dis* 193 (8):501-507
81. Varese F, Smeets F, Drukker M, Lieveise R, Lataster T, Viechtbauer W, Read J, van Os J, Bentall RP (2012) Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull* 38 (4):661-671. doi:10.1093/schbul/sbs050
82. Eichenberg C, Wolters C (2012) Virtual Realities in the Treatment of Mental Disorders: A Review of the Current State of Research. In: *InTech*. doi:<http://dx.doi.org/10.5772/50094>
83. Valmaggia LR, Latif L, Kempton MJ, Rus-Calafell M (2016) Virtual reality in the psychological treatment for mental health problems: A systematic review of recent evidence. *Psychiatry Research* 236:189-195
84. Veling W, Moritz S, van der Gaag M (2014) Brave New Worlds-Review and Update on Virtual Reality Assessment and Treatment in Psychosis. *Schizophrenia Bulletin* 40 (6):1194-1197. doi:10.1093/schbul/sbu125
85. Macedo M, Marques A, Queiros C (2015) Virtual reality in assessment and treatment of schizophrenia: A systematic review. *Jornal Brasileiro de Psiquiatria* 64 (1):70-81
86. Jiandani N, Nair SR, Shukla H (2014) Efficacy of virtual reality exposure therapy in the management of symptoms associated with post traumatic stress disorder. *Value in Health* 17 (7):A572
87. Wiederhold BK, Wiederhold MD (2014) Virtual Reality for Posttraumatic Stress Disorder. *Advances in virtual reality and anxiety disorders*:211-233
88. Rizzo A, Parsons TD, Lange B, Kenny P, Buckwalter JG, Rothbaum B, Difede J, Frazier J, Newman B, Williams J, Reger G (2011) Virtual reality goes to war: A brief review of the future of military behavioral healthcare. *Journal of Clinical Psychology in Medical Settings* 18 (2):176-187

89. Fich LB, Jonsson P, Kirkegaard PH, Wallergard M, Garde AH, Hansen T (2014) Can architectural design alter the physiological reaction to psychosocial stress? A virtual TSST experiment. *Physiology and Behavior* 135:91-97
90. Johns LC, Kompus K, Connell M, Humpston C, Lincoln TM, Longden E, Preti A, Alderson-Day B, Badcock JC, Cella M, Fernyhough C, McCarthy-Jones S, Peters E, Raballo A, Scott J, Siddi S, Sommer IE, Laroi F (2014) Auditory verbal hallucinations in persons with and without a need for care. *Schizophrenia Bulletin* 40 (SUPPL. 4):S255-S264. doi:10.1093/schbul/sbu005
91. Stinson K, Valmaggia LR, Antley A, Slater M, Freeman D (2010) Cognitive triggers of auditory hallucinations: an experimental investigation. *J Behav Ther Exp Psychiatry* 41 (3):179-184. doi:10.1016/j.jbtep.2009.12.003
92. Brett C, Heriot-Maitland C, McGuire P, Peters E (2014) Predictors of distress associated with psychotic-like anomalous experiences in clinical and non-clinical populations. *British Journal of Clinical Psychology* 53 (2):213-227. doi:10.1111/bjc.12036
93. Ward TA, Gaynor KJ, Hunter MD, Woodruff PWR, Garety PA, Peters ER (2014) Appraisals and responses to experimental symptom analogues in clinical and nonclinical individuals with psychotic experiences. *Schizophrenia Bulletin* 40 (4):845-855. doi:10.1093/schbul/sbt094
94. Freeman D, Garety P (2014) Advances in understanding and treating persecutory delusions: a review. *Soc Psychiatry Psychiatr Epidemiol* 49 (8):1179-1189. doi:10.1007/s00127-014-0928-7
95. Howes OD, Murray RM (2014) Schizophrenia: an integrated sociodevelopmental-cognitive model. *Lancet* 383 (9929):1677-1687. doi:10.1016/S0140-6736(13)62036-X
96. Myin-Germeys I, Delespaul P, van Os J (2005) Behavioural sensitization to daily life stress in psychosis. *Psychol Med* 35 (5):733-741
97. Reininghaus U, Kempton MJ, Valmaggia L, Craig TK, Garety P, Onyejiaka A, Gayer-Anderson C, So SH, Hubbard K, Beards S, Dazzan P, Pariante C, Mondelli V, Fisher HL, Mills JG, Viechtbauer W, McGuire P, van Os J, Murray RM, Wykes T, Myin-Germeys I, Morgan C (2016) Stress Sensitivity, Aberrant Salience, and Threat Anticipation in Early Psychosis: An Experience Sampling Study. *Schizophr Bull* 42 (3):712-722. doi:10.1093/schbul/sbv190