



King's Research Portal

DOI:

[10.1002/ejhf.716](https://doi.org/10.1002/ejhf.716)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Behar, J. M., Rajani, R., & Rinaldi, C. A. (2017). Guided left ventricular lead placement for cardiac resynchronization therapy: An opportunity for image integration. *EUROPEAN JOURNAL OF HEART FAILURE*. Advance online publication. <https://doi.org/10.1002/ejhf.716>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Guided left ventricular lead placement for cardiac resynchronisation therapy - an opportunity for image integration.

Jonathan M Behar MBBS BSc MRCP^{1,3}, Ronak Rajani MD MRCP FESC FSCCT²,
Christopher Aldo Rinaldi MD FHRS^{1,2}

1. Department of Imaging Sciences and Biomedical Engineering, King's College London, UK.
2. Department of Cardiology, St Thomas' Hospital, London, UK.
3. Department of Cardiology, Barts Heart Centre, London, UK.

Corresponding author:

Dr Jonathan M Behar

+44 7956183661

Email: jonathanbehar@gmail.com

Key words: Image guidance, CRT, Image integration, LV lead, Multi modality imaging

Word count: 295

Conflicts of interest: None declared.

Funding.

The authors would like to acknowledge the support of the National Institute for Health Research – Biomedical Research Centre at Guys' and St Thomas' Hospital NHS Foundation Trust.

Acknowledgements.

None.

Dear Editors

We read with interest two manuscripts in this month's edition of the journal both exploring the effect of a multi modality imaging approach to guide LV lead implantation for delivery of cardiac resynchronisation therapy (CRT). Suboptimal LV lead positioning is a well established contributor to poor CRT response and as such, techniques utilising cardiac imaging to avoid areas of myocardial fibrosis(1) and / or target dyssynchrony (2) have been of great interest. The studies by Sommer et al (3) and Bertini et al (4) take an interesting approach by combining the strengths of different imaging modalities to guide LV lead placement and in doing so, reporting a reduction in the proportion of patients with a suboptimal response to CRT.

Whilst these results are certainly encouraging, one of the major limitations is that the data output from the imaging modality in each study is reviewed separate to, and alongside X-Ray rather than integrated together. Given the radiolucency of the cardiac silhouette and high variability in the rotation of the left and right sided chambers relative to one another, it is not altogether surprising that using fluoroscopy to determine regional anatomy can be highly inaccurate, particularly with regard to CRT and lead position(5). The next logical step is the co-registration of cardiac imaging data with X-Ray; the resulting visual integration likely to facilitate more accurate image guidance for cardiac procedures. Our group has indeed, designed a purpose built platform which permits real-time cardiac MRI defined scar and dyssynchrony in order to guide LV lead implantation(6). Rapidly evolving technology is enabling real-time image fusion to become a reality for an increasing number of cardiac centres using cardiac MRI, echocardiography and electro-anatomical mapping; future image guided CRT studies should incorporate this highly valuable and translational clinical application.

REFERENCES

1. Leyva F, Foley PWX, Chalil S, Ratib K, Smith RE a, Prinzen F, et al. Cardiac resynchronization therapy guided by late gadolinium-enhancement cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* [Internet]. BioMed Central Ltd; 2011 Jan [cited 2014 Nov 16];13(1):29. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3141552&tool=pmc-entrez&rendertype=abstract>
2. Khan FZ, Virdee MS, Palmer CR, Pugh PJ, O'Halloran D, Elvik M, et al. Targeted left ventricular lead placement to guide cardiac resynchronization therapy: the TARGET study: a randomized, controlled trial. *J Am Coll Cardiol* [Internet]. Elsevier Inc.; 2012 Apr 24 [cited 2014 Nov 12];59(17):1509–18. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22405632>
3. Sommer A, Kronborg MB, Nørgaard BL, Poulsen SH, Bouchelouche K, Böttcher M, et al. Multimodality imaging-guided left ventricular lead placement in cardiac resynchronization therapy: a randomized controlled trial. *Eur J Heart Fail* [Internet]. 2016 Apr 18 [cited 2016 Jul 29]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27087019>
4. Bertini M, Mele D, Malagù M, Fiorencis A, Toselli T, Casadei F, et al. Cardiac resynchronization therapy guided by multimodality cardiac imaging. *Eur J Heart Fail* [Internet]. 2016 Nov [cited 2016 Nov 6];18(11):1375–82. Available from: <http://doi.wiley.com/10.1002/ejhf.605>
5. Sommer A, Kronborg MB, Nørgaard BL, Gerdes C, Mortensen PT, Nielsen JC. Left and right ventricular lead positions are imprecisely determined by fluoroscopy in cardiac resynchronization therapy: a comparison with cardiac computed tomography. *Europace* [Internet]. 2014 Sep [cited 2014 Nov 16];16(9):1334–41. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24687965>
6. Behar JM, Claridge S, Jackson T, Sieniewicz B, Porter B, Webb J, et al. The role of multi modality imaging in selecting patients and guiding lead placement for the delivery of cardiac resynchronization therapy. *Expert Rev Cardiovasc Ther* [Internet]. 2016 Oct 26 [cited 2016 Nov 6];14779072.2016.1252674. Available from: <https://www.tandfonline.com/doi/full/10.1080/14779072.2016.1252674>