



#### **King's Research Portal**

DOI: 10.5281/zenodo.495887

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

*Citation for published version (APA):* Charlton, P. H., Camporota, L., Smith, J., Chaffey, G., Nandi, M., Christie, M. I., Aston, P., Beale, R., & Alastruey-Arimon, J. (2017). Clinical Applications of Attractor Reconstruction Analysis. https://doi.org/10.5281/zenodo.495887

#### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

#### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

#### Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

# **Clinical Applications of Attractor Reconstruction Analysis**

P.H. Charlton <sup>12</sup>, L. Camporota <sup>2</sup>, J. Smith <sup>2</sup>, G. Chaffey <sup>3</sup>,

M. Nandi<sup>1</sup>, M. Christie<sup>1</sup>, P.J. Aston<sup>3</sup>, R. Beale<sup>2</sup>, J. Alastruey<sup>1</sup>

<sup>1</sup> King's College London

<sup>2.</sup> Guy's and St Thomas' NHS Foundation Trust

<sup>3.</sup> University of Surrey

## 1. Analysing pulse waveforms using Attractor Reconstruction (AR)

Pulse waveforms contain a wealth of information

Physiological information in a pulse waveform



**Attractor Reconstruction (AR)** AR is a novel mathematical

	Arterial Blood Pressure Signals (mean)															)					
140																					
	Å	Å	Å	Å	Å	A	Å	Å	A	Å						A	Å	Å	A		







**NHS Foundation Trust** 

Blood pressure and pulse oximetry signals are routinely acquired during physiological assessments. These pulse waveforms contain much physiological information (see right). Currently only simple measures such as heart rate and blood pressure values are routinely extracted from them.

technique for analysing periodic signals. It transforms a **signal** into an **attractor** (see right). The resulting attractor can be used to extract information on the variability and shape of the signal. Changes in the shape or variability of an attractor may indicate changes in cardiovascular health, or disease progression.



# 2. Physiological Insights

## **Vascular Properties**

The properties of the smaller arteries affect the morphology of attractors obtained from a pulse oximetry waveform. This is demonstrated for vascular stiffness and diameter below.

### **Cardiac Properties**

The properties of the heart affect attractors, even when measured distally using a pulse oximeter waveform. Two examples are shown below: ejection time and

## **Physiological State**

The changes shown to the left were obtained from simulations of blood flow. Changes in attractors are also observed during physiological changes, as shown below.

## **Measurement Site**

The morphology of attractors varies according to the site at which they are measured, as shown by the four blood pressure attractors below.



**Detection of Sepsis** 3.

Changes in attractor shape which may be associated with the progression of sepsis

## 4. Respiratory Monitoring 5. Measurement Accuracy





AR may provide early detection of sepsis

AR can improve respiratory rate (RR)

AR can be used to identify and eliminate

Sepsis is a heightened response to infection, which increases morbidity and mortality. Early detection is essential to prevent its progression. AR may allow markers of sepsis to be derived from physiological signals. This is due to the impact of sepsis on the cardiac, vascular and autonomic nervous systems, all of which influence the pulse waveform.

#### estimation from the pulse oximeter

RR is usually measured by manually counting breaths. Techniques have been developed to estimate RR electronically from the pulse oximetry signal, although these are susceptible to inaccuracies. We have used AR to improve the accuracy of RRs derived from a pulse oximeter.

#### inaccurate clinical measurements

Several clinical measurements are derived from physiological signals. The signals must be of high quality to ensure accurate measurements. Attractors change dramatically during periods of low signal quality. Therefore, AR could be used to automatically discard inaccurate measurements.

## **Contact:** Peter.Charlton [at] kcl.ac.uk

# III KING'S HEALTH PARTNERS

This research was supported by: an EPSRC Impact Acceleration Award to PHC and JA; a National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) Early Career Award to PHC; the NIHR BRC based at Guy's and St Thomas' NHS Foundation Trust (GSTT) and King's College London (KCL). The views expressed are those of the authors and not necessarily those of the EPSRC, NHS, NIHR, Department of Health, GSTT or KCL.

## Pioneering better health for all