



## King's Research Portal

DOI:

[10.1017/S0033291717003506](https://doi.org/10.1017/S0033291717003506)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

McCutcheon, R., Bloomfield, M. A. P., Dahoun, T., Quinlan, M., Terbeck, S., Mehta, M., & Howes, O. (2018). Amygdala Reactivity in Ethnic Minorities and its Relationship to the Social Environment: An fMRI Study. *Psychological Medicine*, 48(12), 1985-1992. <https://doi.org/10.1017/S0033291717003506>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## **Amygdala Reactivity in Ethnic Minorities and its Relationship to the Social Environment: An fMRI Study**

Robert McCutcheon<sup>1,2,3</sup>, MRCPsych

Michael A.P. Bloomfield<sup>1,2,3,4,5</sup>, PhD

Tarik Dahoun<sup>2,3</sup>, MD

Marina Quinlan<sup>2</sup>, BSc

Sylvia Terbeck<sup>6</sup>, PhD

Mitul Mehta<sup>7</sup>, PhD

Oliver Howes<sup>1,2,3,\*</sup>, PhD

1. Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, Kings College London, De Crespigny Park, London SE5 8AF, UK
2. Psychiatric Imaging Group, Robert Steiner MR Unit, MRC London Institute of Medical Sciences, Hammersmith Hospital, London W12 0NN, UK
3. Psychiatric Imaging Group, Institute of Clinical Sciences, Faculty of Medicine, Imperial College London, Du Cane Road, London W12 0NN, UK
4. Division of Psychiatry, University College London, 6th Floor, Maple House, 149 Tottenham Court Road, London WC1T 7NF, UK
5. Clinical Psychopharmacology Unit, Research Department of Clinical, Educational and Health Psychology, University College London, 1–19 Torrington Place, London WC1E 6BT, UK
6. School of Psychology, Plymouth University, Drake Circus, Plymouth, PL48AA
7. Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, Kings College London, De Crespigny Park, London SE5 8AF, UK

\*Corresponding author: oliver.howes@kcl.ac.uk tel: ++44 (0)207 848 0355, Box 67, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, Kings College London, De Crespigny Park, London SE5 8AF, UK

**Key Words:** Schizophrenia, Psychosis, Amygdala, fMRI, Ethnic density, Ethnicity, Ethnic minority, risk factors, Paranoia

## **Abstract**

**Background:** Ethnic minority individuals have an increased risk of developing a psychotic disorder, particularly if they live in areas of ethnic segregation, or low own group ethnic density. The neurobiological mechanisms underlying this ethnic minority associated risk are unknown. We used functional MRI to investigate neural responses to faces of different ethnicity, in individuals of black ethnicity, and a control group of white British ethnicity individuals

**Methods:** 20 individuals of black ethnicity, and 22 individuals of white British ethnicity underwent a 3T MRI scan while viewing faces of black and white ethnicity. Own group ethnic density was calculated from the 2011 census. Neighbourhood segregation was quantified using the Index of Dissimilarity method.

**Results:** At the within group level, both groups showed greater right amygdala activation to outgroup faces. Between groups, the black ethnicity group showed greater right amygdala activation to white faces, compared to the white ethnicity group. Within the black ethnicity group, individuals living in areas of lower own group ethnic density showed greater right amygdala reactivity to white faces ( $r=-0.61$ ,  $p=0.01$ ).

**Conclusions:** This is the first time an increased amygdala response to white faces has been demonstrated in individuals of black ethnicity. In the black ethnicity group, correlations were observed between amygdala response and neighbourhood variables associated with increased psychosis risk. These results may have relevance for our understanding of the increased rates of paranoia and psychotic disorders in ethnic minority individuals.

**Funding:** Medical Research Council and Wellcome trust

## Background

Being a member of an ethnic minority group is one of the most well established environmental risk factors for schizophrenia (Cantor-graae *et al.* 2005; Bourque *et al.* 2011). Early research focused on the risk associated with migration (Ødegaard 1932). It has subsequently become clear that this increased risk is not secondary to selective migration (Selten *et al.* 2002; van der Ven *et al.* 2015), and the effect persists among second-generation migrants, indicating that the risk is associated with being a member of a minority group, rather than the act of migration itself (Hutchinson *et al.* 1996; Bresnahan *et al.* 2007; Bourque *et al.* 2011). In various settings the increased risk appears greatest for individuals of black ethnicity (Cantor-graae *et al.* 2005; Bresnahan *et al.* 2007). In the United Kingdom, individuals of black ethnicity were found to have incidence rate ratios compared to the white British population of between six and nine (Fearon *et al.* 2006). This risk also appears to be somewhat specific for psychosis. Although there is some evidence that affective disorders may have slightly increased incidence rates in first generation immigrants, this appears to be driven by manic presentations, and there is no evidence of increased rates of depression or anxiety in the UK black ethnicity population (Shaw *et al.* 1999; Brugha *et al.* 2004; Weich *et al.* 2004; Swinnen & Selten 2007; Mindlis & Boffetta 2017). Similarly, in the United States a number of studies have reported reduced rates of depression and anxiety in African Americans (Kessler *et al.* 2003; Riolo *et al.* 2005; Breslau *et al.* 2006; Himle *et al.* 2009; Asnaani *et al.* 2010; Gibbs *et al.* 2013).

Interestingly, despite the fact that communities with high proportions of ethnic minority individuals are frequently the most deprived, living in an area where one's own ethnic group forms a larger proportion of the population is relatively protective

McCutcheon et al.  
against schizophrenia (Shaw *et al.* 2012; Kirkbride *et al.* 2014). One hypothesis is that higher own group ethnic density buffers the individual from the adverse psychiatric consequences of social isolation and racist experiences (Bécares *et al.* 2009). Epidemiological work has also suggested that greater levels of segregation between members of ethnic minority groups and the rest of the community may be linked to a greater incidence of psychotic disorders (Kirkbride *et al.* 2014).

Cognitive models propose that minority status is associated with greater levels of social threat (Combs *et al.* 2002; Morgan *et al.* 2010). However, the neurobiological mechanisms underlying both the increased risk of psychosis in ethnic minority groups, and the protective effects of increased own group ethnic density, are not known. One previous fMRI study investigated the neural correlates of the stress response in Turkish migrants living in Germany, and identified differences in anterior cingulate activation (Akdeniz *et al.* 2014). The amygdala, is also an area of potential interest given it is a key component of the neural circuit processing threatening experiences (Chekroud *et al.* 2014; Fox *et al.* 2015; Underwood *et al.* 2016).

A greater right amygdala neural response to pictures of faces of people with an outgroup ethnicity, compared to those with in-group ethnicity, has been interpreted as representing threat appraisal (Chekroud *et al.* 2014). This outgroup effect has been consistently demonstrated in multiple studies of individuals of white ethnicity, but has not been thoroughly investigated in individuals of black ethnicity (Cunningham *et al.* 2004; Lieberman *et al.* 2005; Wheeler & Fiske 2005; Ronquillo *et al.* 2007; Chekroud *et al.* 2014; Fox *et al.* 2015). In psychosis, meanwhile, amygdala hyperactivity has been linked to paranoid symptoms (Goghari *et al.* 2010; Pinkham *et al.* 2015; Underwood *et al.* 2016). Individuals from ethnic minority populations typically have much greater exposure to outgroup individuals than the native

population. Amygdala hyperactivity to frequent outgroup exposure could therefore partially underlie the greater levels of paranoia seen in both ethnic minority patient and population samples (Bhugra *et al.* 2000; Combs *et al.* 2002; Cohen *et al.* 2004; Veling *et al.* 2007; Wickham *et al.* 2014), and potentially have relevance as regards the increased rates of psychotic illnesses in these groups.

In view of this evidence, we hypothesised that individuals who identified as being of black ethnicity would display a greater right amygdala response to white faces, than individuals who identified as being of white ethnicity; who would in turn show a greater response to black faces. Within the black ethnic minority group we further hypothesised that increased amygdala reactivity to white faces would correlate with increased ethnic segregation, and decreased own group ethnic density, as these ethnicity related neighbourhood variables are associated with increased psychosis risk.

## **Methods**

### *Participants*

Healthy volunteers were recruited throughout the UK via newspaper, online and leaflet advertising, and in person recruitment, as part of a study examining environmental risk factors for psychosis. Participants were aged 18-45, with no history of mental illness, in good physical health, and with capacity to give informed consent.

The black ethnic minority group needed to be a first or second generation migrant, to self-identify as being of black ethnicity, and were classified as of either black Caribbean or black African ethnicity based on either their own (first generation

migrants), or their parents' (second generation migrants) country of origin. The white British ethnicity group needed to self-identify as being of white British ethnicity and have at least one parent of UK nationality. The white British ethnicity group was age matched within 5 years to the black ethnic minority group. All subjects provided written informed consent after the study had been fully explained.

### *Sociodemographic assessment*

Population density, and own group ethnic density as a fraction of the total population, were obtained from the 2011 census; Indices of Multiple Deprivation (where a lower ranking equates to a more deprived area) were obtained from 2015 data (*English Indices of Deprivation 2015*). Values were obtained for each participant's current statistical lower layer super output area (LSOA, geographical areas with populations of roughly 1500 individuals).

In order to quantify ethnic segregation, the Index of Dissimilarity method was used to calculate an index of segregation at the ward-level, using census output areas as the smaller geographic unit. A score of 0 is equivalent to no segregation, while a score of 1 is equivalent to complete segregation (see supplementary methods for further information) (Yalonetzky 2015). Own group ethnic density and ethnic segregation were both calculated with participants assigned to either black African, black Caribbean or white British ethnicity.

### *fMRI task*

An event related, implicit face perception task was employed based on a previously reported experimental design (see figure 1) (Cunningham *et al.* 2004). During the task, participants indicated whether a visual stimulus appeared to the left or right of a fixation cross. The stimuli consisted of either black or white male faces presented for

30 ms (short trials) or 525 ms (long trials); or a white square also presented for 30 or 525 ms. Faces were taken from the Stanford University faces database ([https://stanforduniversity.qualtrics.com/SE/?SID=SV\\_aX0ovSkASZR9Py4](https://stanforduniversity.qualtrics.com/SE/?SID=SV_aX0ovSkASZR9Py4)), 18 faces of each ethnicity were chosen and matched on ratings of attractiveness, stereotypicality and age. In order to mask the short duration stimuli, all stimuli were preceded by an abstract picture for 30 ms and followed by the same picture for either 525 ms (short trials) or 30 ms (long trials). As a result all trials lasted a total of 585ms. During the short trials, the participants experience was of seeing an abstract picture and they were not aware of the implicitly presented face. We checked this after the experiment and no participant reported seeing a face during presentation of these stimuli. During the long trials they saw either a white square or a face. A fixation cross appeared for 2055 ms between trials. Each face trial was followed by 4 white square trials giving an interstimulus interval between faces ranging from 10.62 to 11.12s. Participants completed 3 runs. During each run participants would be presented with 6 short trials of black faces, 6 long trials of black faces, 6 short trials of white faces and 6 long trials of white faces, shown in a random order without replacement.

### *Data acquisition*

Imaging data was acquired using a Philips 3T Inera magnetic resonance imaging system. Functional imaging consisted of T2\* weighted transverse echo planar image (EPI) slices. Each run consisted of 161 volumes, collected in an ascending direction, with 2.19mm x 2.19mm voxel dimensions in plane, repetition time (TR) - 2000 ms, and echo time (TE) – 30 ms.

The EPI scans were followed by a gradient-echo scan (TR= 9.7ms, TE=4.6ms, flip angle – 9°, slice thickness 1.20 mm, 0.94 x 0.94 mm in plane resolution, 150 slices).



*fMRI data analysis*

Analysis of fMRI data was undertaken using SPM8 (<http://www.fil.ion.ucl.ac.uk/spm/>) in Matlab 7.9. The first two volumes for each participant were discarded so as to avoid T1 equilibration effects. Slice timing correction was applied to each volume. Spatial realignment to the third volume was followed by coregistration of each participant's functional and anatomical data. Images were then normalized into standard MNI space using the normalization parameters estimated by T1 structural image unified segmentation. Smoothing of the resampled images (3mmx3mmx3mm) was with a Gaussian kernel of 8mm full-width-half-maximum.

First level analysis using the general linear model included six conditions (short and long presentations of white square, white faces and black faces), and individual events were convolved with the canonical haemodynamic response function. The model included the six movement parameters from realignment, and six volume to volume movement parameters, as regressors of no interest, additionally any volumes where between volume movement was greater than 0.5mm were removed. We excluded any runs with extended shifts of over 2mm. Participants were included in the second level analysis if they had at least two usable runs. First level contrasts were constructed between faces and white squares.

Second level within group analyses were performed using paired t-tests. Second level between groups analyses were performed using independent samples t-tests, and a 2 by 2 flexible factorial ANOVA implemented in SPM8. Following from a priori hypotheses, region of interest (ROI) analyses were conducted for the amygdala using Marsbar (<http://marsbar.sourceforge.net/>) and the included Automated

Anatomical Labelling ROI library. As per SPM conventions t-test p values are reported as one-tailed.

### *Statistical Analysis*

Analyses were carried out using SPSS for Macintosh version 23.0. After exclusion of outliers of more than 1.5 interquartile ranges below or above the 1<sup>st</sup> or 3<sup>rd</sup> quartiles, continuous variables were assessed for normality using the Shapiro-Wilks test. Differences between group means were assessed using an independent samples *t*-test for normally distributed variables, after using Levene's test to check for equality of variances. Mann-Whitney *U* tests were used for non-normally distributed variables. Between group differences in terms of the magnitude of the amygdala outgroup effect, was tested with a two-tailed independent samples *t*-test. Correlations between normally distributed variables were assessed using Pearson's product moment coefficient, and were only reported if they remained significant after removal of outliers defined as a Cook's *d* of  $>n/4$ . Correlations involving non-normally distributed variables were assessed using Spearman's rank correlation coefficient. All correlations were two-tailed with  $p<0.05$  defined as significant. Neighbourhood variables found on bivariate testing to show a significant correlation with right amygdala response were subsequently entered into a stepwise regression with right amygdala response as the dependent variable.

### *Data Availability*

Imaging and demographic data is available from the corresponding author on request.

## **Results**

### *Sociodemographic and neighbourhood variables*

42 individuals were included in the study (black ethnic minority group  $n=20$ , white British ethnicity group  $n=22$ ). There were no significant differences between groups in terms of sex or age. In terms of neighbourhood variables the black ethnic minority group had significantly lower own group ethnic density percentages and lived in significantly more deprived, less segregated, and more densely populated areas (demographics of the entire sample are reported in supplementary table 1, while details of those included in the imaging analysis are described in table 1). In the black ethnicity group, participant characteristics in terms of age, gender, and deprivation were similar to UK averages. In the white ethnicity group the study population was younger compared to the UK median age but otherwise similar (see supplementary material).

Within the black ethnic minority group, own group ethnic density negatively correlated with degree of segregation ( $r_s=-.809$ ,  $p<0.001$ ) (correlations between other neighbourhood variables are described in the supplementary data).

### *Functional imaging results*

3 participants from each group were excluded due to excessive movement or failed scans. The 30 ms presentations did not reliably induce right amygdala activation. Neither black nor white faces elicited significant right amygdala activation in either the white British or black ethnic minority groups (all contrasts against the white square baseline  $p>0.05$ ). In addition, there were no significant differences between groups, or within groups between stimuli, for the 30 ms presentations. All results reported below are for 525 ms presentations.

At the within group level, both groups individually showed greater right amygdala activation to outgroup faces (white British ethnicity group  $t=1.90$ ,  $p=0.036$ ; black

ethnic minority group  $t=2.38$ ,  $p=0.015$ ) (see figure 2). This was reflected in a significant group x task interaction ( $F=7.85$ ,  $p=0.008$ ) (see figure 2). The magnitude of the outgroup effect was not significantly different between groups ( $t=1.1$ ,  $p=0.3$ ).

Between groups, compared to the white British ethnicity group, the black ethnic minority group showed significantly greater right amygdala activation to white faces ( $t=1.84$ ,  $p=0.038$ ), and significantly lower right amygdala activation to black faces ( $t=1.80$ ,  $p=0.040$ ) (see figure 2).

Whole brain exploratory analyses were conducted to investigate the potential involvement of other regions in the effects and to enable comparisons with studies investigating other regions (peak threshold  $p<0.001$ , cluster threshold  $p<0.05$  FWE corrected). No significant results were observed for the between groups, or group by task interaction analyses. For the black ethnicity group, white faces>black faces contrast one small cluster was observed in the right cerebellum ( $p=0.007$ , cluster size - 76 voxels, coordinates  $x=33$ ,  $y=-58$ ,  $z=-26$ ).

#### *Relationships between imaging results and neighbourhood variables*

Within the black ethnic minority group right amygdala response to white faces significantly correlated with living in an area of greater ethnic segregation ( $r_p=0.831$ ,  $p<0.001$ ) and lower own group ethnic density ( $r_s=-0.611$ ,  $p=0.009$ ) (see figure 3 and supplementary material for results using alternative measures of ethnic density). The right amygdala response also positively correlated with greater population density ( $r_p=0.627$ ,  $p=0.002$ ) and lower levels of deprivation ( $r_s=0.601$ ,  $p=0.011$ ). These findings survive Bonferroni correction accounting for the four separate tests ( $p<0.0125$ ). All four neighbourhood variables were entered into a stepwise regression with right amygdala response as the dependent variable; from this only

own group ethnic density remained a significant predictor ( $r=0.689$ ,  $p=0.002$ , see Table 2). In the white ethnicity group no significant correlations were observed between neighbourhood variables and right amygdala response to black faces.

## Discussion

We demonstrated increased amygdala activation to white faces in a group of individuals of black ethnicity – both within group compared to black faces, and between groups when compared white British individuals. The white British ethnicity group showed a greater amygdala response to black faces both within, and between group. In addition, within the black ethnic minority group, a stepwise regression including neighbourhood variables that correlated with amygdala reactivity, found that lower own group ethnic density was significantly associated with greater amygdala reactivity to white faces.

Our finding of increased right amygdala reactivity to black faces in the white British ethnicity group, has been repeatedly demonstrated in white ethnicity participants (Cunningham *et al.* 2004; Lieberman *et al.* 2005; Wheeler & Fiske 2005; Ronquillo *et al.* 2007; Chekroud *et al.* 2014). This is, however, to our knowledge the first time that increased reactivity to white faces has been unambiguously demonstrated in individuals of black ethnicity. One previous study demonstrating greater amygdala activity to outgroup faces included four participants of black ethnicity, but did not comment on this subgroup specifically (Hart *et al.* 2000). A study by Lieberman *et al.* found that African-American ethnic minority individuals showed an increased amygdala response while viewing black faces (Lieberman *et al.* 2005). This study, however, included only nine African-American subjects and used a different task that focused on differences in perceptual and verbal encoding, which may explain the inconsistency with our findings. While our findings suggest a general out group

effect, this finding may potentially have greater psychological impact for ethnic minority groups given their generally greater exposure to outgroups.

This is also the first time that amygdala reactivity has been linked to neighbourhood variables related to ethnic minority associated risk for psychosis. Living in areas of low own group ethnic density has been shown to increase the risk of psychosis in ethnic minorities (Bosqui *et al.* 2014). Our results extend these findings by indicating that lower own group ethnic density is also linked to greater amygdala response to outgroup faces in black individuals. Living in an environment where one is more isolated from one's own ethnic group (i.e. an area of low own group ethnic density), could conceivably contribute to outgroup individuals being perceived as a greater threat. Taken with evidence that experiences of racism are correlated with amygdala activation to white faces in black individuals (Greer *et al.* 2012), this suggests that environmental exposures are associated with functional alterations in the brain circuits involved in threat processing. While a number of our results were significant when examining the bilateral amygdala (see supplementary material), our findings were driven by the right amygdala. The right amygdala appears to be particularly implicated in the processing of race, although the underlying reasons for this remain unclear (Chekroud *et al.* 2014).

### *Methodological Considerations*

We investigated a group of black ethnic minority individuals who were first or second generation migrants relative to the white British ethnic majority. This group appears to display the greatest increase in incidence of psychosis, compared to white British individuals (Fearon *et al.* 2006; Coid *et al.* 2008). However, it should be recognised that other ethnic groups, including non-British white individuals, and Asian individuals, have increased rates of psychosis as well, and migrants may be exposed

to other adverse experiences in addition to those associated with being a member of a minority group (Fearon *et al.* 2006; Coid *et al.* 2008). Ethnic minority status itself is likely a crude proxy for multiple complex, interacting factors that cumulatively act to increase the risk of psychosis. Our study used a convenience sampling procedure and future work should determine how representative the sample is of the population in general.

Our finding of an association between ethnicity related neighbourhood variables and amygdala reactivity suggests that this mechanism may have some relevance for our understanding of the neurobiological basis of ethnic minority associated psychosis risk. This, however, is a speculative interpretation, and the current study does not establish causality. Capturing the relationship between amygdala functioning and real world experiences of being in a minority group could potentially be accomplished using experience sampling methodologies (Gevonden *et al.* 2016; Reininghaus *et al.* 2016).

Study participants were intentionally unaffected by mental illness, which precluded investigating direct associations with psychopathology. As our participants did not have a mental illness it is important to recognise the possibility that our findings may be interpreted as physiologically appropriate or neurobiological correlates of psychological resilience. However, the fact that in patients with schizophrenia greater amygdala responses are associated with paranoid symptoms suggests this may not be the case (Goghari *et al.* 2010; Pinkham *et al.* 2015; Underwood *et al.* 2016). Amygdala dysfunction is implicated in a wide range of disorders, so even if aberrations identified in the current study have pathoetiological relevance, it may not be specific to psychotic disorders. Furthermore, drawing parallels with patient groups is limited by the fact that our task specifically measured differences in

amygdala reactivity to faces of differing ethnicities, whereas most probes used in patient populations tend to involve emotion processing. Investigating amygdala response to faces of differing ethnicity in patient populations would assist with determining if this mechanism is linked to psychopathology.

When relating neighbourhood variables to amygdala response we were primarily interested in the influence of own group ethnic density, and ethnic segregation. We additionally measured population density and deprivation ranking. The stepwise regression resulted in a model only including own group ethnic density. Nearly all these variables, however, are strongly correlated with one another and thus determining which play mediating as opposed to confounding roles is not possible in the current study. Larger sample sizes would allow for the investigation of interactions between variables, while replication in other settings would establish whether the degree to which the effect is generalizable. Additionally while the link between psychosis risk and own group ethnic density is well established, the evidence for the influence of ethnic segregation is less clear (Kirkbride *et al.* 2007, 2014).

We did not observe significant between or within group differences, for the 30 ms stimuli presentations. This is surprising given that this subliminal presentation has previously been found to evoke greater amygdala responses to outgroup faces than the 525ms presentation (Cunningham *et al.* 2004). Potential reasons underlying this discrepancy may include differences in masking stimuli between the current and previous study, or differences as regards image characteristics such as levels of brightness and contrast which subliminal stimuli may be particularly sensitive to.



*Implications for understanding the mechanisms underlying ethnic minority associated risk for psychosis*


Cognitive models of psychosis propose an increased level of social threat anticipation as part of a pathway to the development of persecutory beliefs (Bentall & Fernyhough 2008). Of note, a higher degree of paranoid symptomatology has been described in ethnic minority groups, both in patient (Bhugra *et al.* 2000; Veling *et al.* 2007), and healthy populations (Combs *et al.* 2002; Cohen *et al.* 2004; Wickham *et al.* 2014). In the current study both the black ethnic minority group and the white British ethnicity group displayed increased amygdala reactivity to outgroup faces. The consequence for each group, however, may be different given that ethnic minorities typically have much greater outgroup contact in day-to-day life. Following from this, if the increased amygdala response to outgroup faces is interpreted as a marker of threat, this suggests that ethnic minority individuals have more frequent threat experience.

Although amygdala hyperactivity has been linked to paranoid symptoms in psychosis (Goghari *et al.* 2010; Pinkham *et al.* 2015; Underwood *et al.* 2016), reactivity during facial processing is not consistently raised in schizophrenia (Taylor *et al.* 2012). The black ethnic minority group in the current study did not show overall increased amygdala reactivity, only to specific stimuli. A differential sensitivity to various stimuli may partially explain the variability of findings observed in studies of amygdala function in schizophrenia.


## **Conclusions**

Amygdala reactivity to white faces is increased in black ethnic minority individuals, and correlated with measures of own group ethnic density and segregation. These


findings indicate that black ethnic minority individuals show the same response to outgroup faces seen in white ethnic majority groups, and also suggest that in this population ethnic segregation and lower own group ethnic density is associated with greater amygdala reactivity to outgroup faces.



**Figure 1:** fMRI task. Participants judge whether an image is to the left or right of a fixation cross. Each face trial (top row) is followed by four white square trials (bottom row).



**Figure 2:** (A) Increased amygdala activation in black ethnicity group compared to white British ethnicity group for white>black faces contrast during 525ms presentations. Functional maps are unmasked and thresholded at  $P < 0.001$  whole brain, uncorrected for display purposes. (B) Mean parameter estimates for right amygdala ROI for both groups, for faces (525 ms) vs white square contrasts, error bars =  $\pm$  SEM. \* $p < 0.05$



**Figure 3:** Scatter plots for right amygdala response to whitefaces within the black ethnicities group, against: (A) % own group ethnicity of current LSOA ( $r_s=-.611$ ,  $p=0.009$ ) (B) Index of segregation ( $r_p=0.831$ ,  $p<0.001$ )

## **Acknowledgements**

### **Author contributions**

R.M. designed and performed experiments, analysed data, and wrote the paper. M.B. designed and performed experiments, and reviewed the manuscript. T.D. and M.Q. performed experiments and reviewed the paper. S.T. designed experiments and reviewed the manuscript. M.M. analysed data and reviewed the paper. O.H. obtained funding, designed the study and wrote the paper.

### **Acknowledgements**

This study was funded by Medical Research Council-UK (no. MC-A656-5QD30), Maudsley Charity (no. 666), Brain and Behavior Research Foundation, and Wellcome Trust (no. 094849/Z/10/Z) grants to O.H. and the *National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London*. R.M.'s work is supported by the Wellcome Trust (no. 200102/Z/15/Z). T.D.'s work is supported by a EU-FP7 MC-ITN IN-SENS grant (no. 607616).

*The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.*

**Financial Disclosures**

O.H. has received investigator-initiated research funding from and/or participated in advisory/speaker meetings organised by Astra-Zeneca, Autifony, BMS, Eli Lilly, Heptares, Jansenn, Lundbeck, Lyden-Delta, Otsuka, Servier, Sunovion, Rand and Roche. Neither Dr Howes or his family have been employed by or have holdings/ a financial stake in any biomedical company. M.M. has consulted for Cambridge Cognition, Lundbeck and Forum Pharmaceuticals in the past 3 years. He has also received research funding from Takeda, Eli Lilly and Roche. The other authors declare no competing financial interests.

## References

- Akdeniz C, Tost H, Streit F, Haddad L, Wüst S, Schäfer A, Schneider M, Rietschel M, Kirsch P, & Meyer-Lindenberg A** (2014). Neuroimaging Evidence for a Role of Neural Social Stress Processing in Ethnic Minority–Associated Environmental Risk. *JAMA Psychiatry* **71**, 672.
- Asnaani A, Richey JA, Dimaite R, Hinton DE, & Hofmann SG** (2010). A Cross-Ethnic Comparison of Lifetime Prevalence Rates of Anxiety Disorders. *The Journal of Nervous and Mental Disease* **198**, 551–555.
- Bécares L, Nazroo J, & Stafford M** (2009). The buffering effects of ethnic density on experienced racism and health. *Health and Place* **15**, 670–678.
- Bentall RP, & Fernyhough C** (2008). Social predictors of psychotic experiences: specificity and psychological mechanisms. *Schizophrenia bulletin* **34**, 1012–20.
- Bhugra D, Hilwig M, Corridon B, Neehall J, Rudge S, Mallett R, & Leff J** (2000). A comparison of symptoms in cases with first onset of schizophrenia across four groups. *European journal of psychiatry* **14**, 241–250.
- Bosqui TJ, Hoy K, & Shannon C** (2014). A systematic review and meta-analysis of the ethnic density effect in psychotic disorders. *Social Psychiatry and Psychiatric Epidemiology* **49**, 519–529.
- Bourque F, van der Ven E, & Malla A** (2011). A meta-analysis of the risk for psychotic disorders among first- and second-generation immigrants. *Psychological medicine* **41**, 897–910.
- Breslau J, Aguilar-Gaxiola S, Kendler KS, Su M, Williams D, & Kessler RC** (2006). Specifying race-ethnic differences in risk for psychiatric disorder in a USA national sample. *Psychological medicine* **36**, 57–68.
- Bresnahan M, Begg MD, Brown A, Schaefer C, Sohler N, Insel B, Vella L, & Susser E** (2007). Race and risk of schizophrenia in a US birth cohort: Another example of health disparity? *International Journal of Epidemiology* **36**, 751–758.

- Brugha T, Jenkins R, Bebbington P, Meltzer H, Lewis G, Farrell M, Le L, & Park H** (2004). Risk factors and the prevalence of neurosis and psychosis in ethnic groups in Great Britain. *Social Psychiatry and Psychiatric Epidemiology* **39**, 939–946.
- Cantor-graee E, Ph D, & Selten J** (2005). Schizophrenia and Migration : A Meta-Analysis and Review. *American Journal of Psychiatry* **162**, 12–24.
- Chekroud AM, Everett J a C, Bridge H, & Hewstone M** (2014). A review of neuroimaging studies of race-related prejudice: does amygdala response reflect threat? *Frontiers in human neuroscience* **8**, 179.
- Cohen CI, Magai C, Yaffee R, & Walcott-Brown L** (2004). Racial Differences in Paranoid Ideation and Psychoses in an Older Urban Population. *American Journal of Psychiatry* **161**, 864–871.
- Coid JW, Kirkbride JB, Barker D, Cowden F, Stamps R, Yang M, & Jones PB** (2008). Raised Incidence Rates of All Psychoses Among Migrant Groups. *Archives of General Psychiatry* **65**, 1250.
- Combs DR, Penn DL, & Fenigstein A** (2002). Ethnic differences in subclinical paranoia: an expansion of norms of the paranoia scale. *Cultural diversity & ethnic minority psychology* **8**, 248–256.
- Cunningham W a, Johnson MK, Raye CL, Chris Gatenby J, Gore JC, & Banaji MR** (2004). Separable neural components in the processing of black and white faces. *Psychological science* **15**, 806–13.
- English Indices of Deprivation 2015:**  
[www.gov.uk/government/statistics/english-indices-of-deprivation-2015](http://www.gov.uk/government/statistics/english-indices-of-deprivation-2015) (n.d.).
- Fearon P, Kirkbride JB, Morgan C, Dazzan P, Morgan K, Lloyd T, Hutchinson G, Tarrant J, Fung WLA, Holloway J, Mallett R, Harrison G, Leff J, Jones PB, & Murray RM** (2006). Incidence of schizophrenia and other psychoses in ethnic minority groups: results from the MRC AESOP Study. *Psychological medicine* **36**, 1541–50.



- Fox AS, Oler JA, Tromp DPM, Fudge JL, & Kalin NH** (2015). Extending the amygdala in theories of threat processing. *Trends in Neurosciences* **38**, 319–329.
- Gevonden M, Myin-Germeys I, Wichers M, Booij J, van den Brink W, van Winkel R, & Selten J-P** (2016). Reactivity to social stress in ethnic minority men. *Psychiatry Research* **246**, 629–636.
- Gibbs TA, Okuda M, Oquendo MA, Lawson WB, Wang S, Thomas YF, & Blanco C** (2013). Mental health of African Americans and Caribbean blacks in the United States: Results from the national epidemiological survey on alcohol and related conditions. *American Journal of Public Health* **103**, 330–338.
- Goghari VM, Sponheim SR, & MacDonald AW** (2010). The functional neuroanatomy of symptom dimensions in schizophrenia: A qualitative and quantitative review of a persistent question. *Neuroscience and Biobehavioral Reviews* **34**, 468–486.
- Greer TM, Vendemia JMC, & Stancil M** (2012). Neural correlates of race-related social evaluations for African Americans and white Americans. *Neuropsychology* **26**, 704–12.
- Hart AJ, Whalen PJ, Shin LM, Mcinerney SC, Fischer H, & Rauch SL** (2000). Differential response in the human amygdala to racial outgroup vs ingroup face stimuli. *Neuroreport* **11**, 2351–2355.
- Himle JA, Baser RE, Joseph R, Denise R, & Jackson JS** (2009). Anxiety disorders among African Americans, blacks of Caribbean descent, and non-Hispanic whites in the United States. *Journal of Anxiety Disorders* **23**, 578–590.
- Hutchinson G, Takei N, Fahy T a., Bhugra D, Gilvarry C, Moran P, Mallett R, Sham P, Leff J, & Murray RM** (1996). Morbid risk of schizophrenia in first-degree relatives of White and African-Caribbean patients with psychosis. *British Journal of Psychiatry* **169**, 776–780.
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, Rush AJ,**

- Walters EE, Wang PS, & National Comorbidity Survey Replication** (2003). The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA* **289**, 3095–105.
- Kirkbride JB, Jones PB, Ullrich S, & Coid JW** (2014). Social deprivation, inequality, and the neighborhood-level incidence of psychotic syndromes in East London. *Schizophrenia bulletin* **40**, 169–80.
- Kirkbride JB, Morgan C, Fearon P, Dazzan P, Murray RM, & Jones PB** (2007). Neighbourhood-level effects on psychoses: re-examining the role of context. *Psychological medicine* **37**, 1413–25.
- Lieberman MD, Hariri A, Jarcho JM, Eisenberger NI, & Bookheimer SY** (2005). An fMRI investigation of race-related amygdala activity in African-American and Caucasian-American individuals. *Nature neuroscience* **8**, 720–2.
- Mindlis I, & Boffetta P** (2017). Mood disorders in first- and second-generation immigrants: systematic review and meta-analysis. *The British Journal of Psychiatry* **210**, 182–189.
- Morgan C, Charalambides M, Hutchinson G, & Murray RM** (2010). Migration, ethnicity, and psychosis: toward a sociodevelopmental model. *Schizophrenia bulletin* **36**, 655–64.
- Ødegaard Ø** (1932). *Emigration and insanity: a study of mental disease among the Norwegianborn population of Minnesota*. Levin & Munksgaard.
- Pinkham AE, Liu P, Lu H, Kriegsman M, Simpson C, & Tamminga C** (2015). Amygdala Hyperactivity at Rest in Paranoid Individuals With Schizophrenia. *American Journal of Psychiatry* **172**, 784–792.
- Reininghaus U, Kempton MJ, Valmaggia L, Craig TKJ, Garety P, Onyejiaka A, Gayer-Anderson C, So SH, Hubbard K, Beards S, Dazzan P, Pariante C, Mondelli V, Fisher HL, Mills JG, Viechtbauer W, McGuire P, van Os J, Murray RM, Wykes T, Myin-Germeys I, & Morgan C** (2016). Stress Sensitivity, Aberrant

Salience, and Threat Anticipation in Early Psychosis: An Experience Sampling

Study. *Schizophrenia Bulletin* **42**, 712–722.

**Riolo SA, Nguyen TA, Greden JF, & King CA** (2005). Prevalence of depression by race/ethnicity: Findings from the national health and nutrition examination survey III.

*American Journal of Public Health* **95**, 998–1000.

**Ronquillo J, Denson TF, Lickel B, Lu ZL, Nandy A, & Maddox KB** (2007). The effects of skin tone on race-related amygdala activity: An fMRI investigation. *Social Cognitive and Affective Neuroscience* **2**, 39–44.

**Selten J-P, Cantor-Graae E, Slaets J, & Kahn RS** (2002). Ødegaard's selection hypothesis revisited: schizophrenia in Surinamese immigrants to The Netherlands.

*The American journal of psychiatry* **159**, 669–71.

**Shaw CM, Creed F, Tomenson B, Riste L, & Cruickshank JK** (1999). Prevalence of anxiety and depressive illness and help seeking behaviour in African Caribbeans and white Europeans: two phase general population survey. *British Medical Journal* **318**, 302–306.

**Shaw RJ, Atkin K, Becares L, Albor CB, Stafford M, Kiernan KE, Nazroo JY, Wilkinson RG, & Pickett KE** (2012). Impact of ethnic density on adult mental disorders: Narrative review. *British Journal of Psychiatry* **201**, 11–19.

**Swinnen SGHA, & Selten J** (2007). Mood disorders and migration. *British Journal of Psychiatry* **190**, 6–10.

**Taylor SF, Kang J, Brege IS, Tso IF, Hosanagar A, & Johnson TD** (2012). Meta-analysis of functional neuroimaging studies of emotion perception and experience in schizophrenia. *Biological psychiatry* **71**, 136–45.

**Underwood R, Kumari V, & Peters E** (2016). Cognitive and neural models of threat appraisal in psychosis: A theoretical integration. *Psychiatry Research* **239**, 131–138.

**Veling W, Selten JP, Mackenbach JP, & Hoek HW** (2007). Symptoms at first contact for psychotic disorder: Comparison between native Dutch and ethnic

minorities. *Schizophrenia Research* **95**, 30–38.

**van der Ven E, Dalman C, Wicks S, Allebeck P, Magnusson C, van Os J, & Selten JP** (2015). Testing Ødegaard’s selective migration hypothesis: a longitudinal cohort study of risk factors for non-affective psychotic disorders among prospective emigrants. *Psychological medicine* **45**, 727–34.

**Weich S, Nazroo J, Sproston K, McManus S, Blanchard M, Erens B, Karlsen S, King M, Lloyd K, Stansfeld S, & Tyrer P** (2004). Common mental disorders and ethnicity in England: the EMPIRIC study. *Psychological Medicine* **34**, 1543–1551.

**Wheeler ME, & Fiske ST** (2005). Controlling racial prejudice: social-cognitive goals affect amygdala and stereotype activation. *Psychological science* **16**, 56–63.

**Wickham S, Taylor P, Shevlin M, & Bentall RP** (2014). The impact of social deprivation on paranoia, hallucinations, mania and depression: The role of discrimination social support, stress and trust. *PLoS ONE* **9**

**Yalonetzky G** (2015). Index of Dissimilarity. In *The Wiley Blackwell Encyclopedia of Race, Ethnicity, and Nationalism*, pp1–3. John Wiley & Sons, Ltd: Oxford, UK.

	Black Ethnic minority (n=17)	White British Ethnicity (n=19)	P
Age (years), median (IQR)	24.5 (20.8-32.2)	24.0 (22.5-28.1)	0.98 <sup>a</sup>
Sex n(%): Male	7 (41)	9 (50)	0.71 <sup>b</sup>
Female	10(59)	10 (53)	
Migration n(%): 1 <sup>st</sup> generation	7 (41)		
2 <sup>nd</sup> generation	10(59)		
Black Caribbean, n (%)	3 (18)		
Black African, n (%)	14 (82)		
% own ethnicity, median (IQR)	9.1 (6.0-24.6)	86.3 (77.2-95.4)	<0.001 <sup>a</sup>
Index of Multiple deprivation rank/1000, median (IQR)	7.2 (5.0-13.5)	22.8 (19.0-31.4)	<0.001 <sup>a</sup>
Index of segregation, median (IQR)	0.28 (0.18-0.31)	0.32 (0.28-0.42)	0.008 <sup>a</sup>
Population density: persons per hectare, median (IQR)	91.7 (59.1-126.7)	38.6 (1.9-96.1)	0.005 <sup>a</sup>

<sup>a</sup>Mann-Whitney; <sup>b</sup>chi-square test

**Table 1:** Demographic characteristics of participants included in imaging analysis

	Variable	B	SE	Beta	t	P	R <sup>2</sup>	P
Model A	Constant	3.44	6.41		.537	.601	.571	.027
	Own group ethnic density	-.172	.164	-.399	-1.05	.315		
	Index of Segregation	-12.5	12.5	-.283	-.996	.339		
	Own group ethnic density	.032	.029	.345	1.11	.288		
	Index of Multiple Deprivation	<.001	<.001	.363	1.36	.199		
Model B	Constant	7.52	1.37		5.51	<.001	.475	.002
	Own group ethnic density	-.297	.081	-.689	-3.68	.002		

**Table 2:** Linear regression of relationship between of neighbourhood variables and right amygdala reactivity to white faces. Model A includes all neighbourhood variables. Model B includes only ‘own group ethnicity’ as determined by a stepwise regression.