



King's Research Portal

DOI:
[10.1111/eip.12521](https://doi.org/10.1111/eip.12521)

Document Version
Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Vancampfort, D., De Hert, M., Myin-Germeys, I., Rosenbaum, S., Stubbs, B., Van Damme, T., & Probst, M. (2017). Validity and correlates of the International Physical Activity Questionnaire in first-episode psychosis. *Early Intervention in Psychiatry*. Advance online publication. <https://doi.org/10.1111/eip.12521>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Validity and correlates of the International Physical Activity Questionnaire in first-episode psychosis

Davy Vancampfort^{a,b,*}, Marc De Hert^b, Inez Myin-Germeys^c, Simon Rosenbaum^d, Brendon Stubbs^{e,f},
Tine Van Damme^{a,b}, Michel Probst^{a,b}

^aKU Leuven Department of Rehabilitation Sciences, Leuven, Belgium

^bUPC KU Leuven, Kortenberg-Leuven, Belgium

^cKU Leuven Department of Neurosciences, Centre for Contextual Psychiatry, Leuven, Belgium

^dSchool of Psychiatry, UNSW Australia, The Black Dog Institute, University of New South Wales,
Prince of Wales Hospital, Sydney, Australia

^ePhysiotherapy Department, South London and Maudsley NHS Foundation Trust, Denmark Hill,
London, UK

^fHealth Service and Population Research Department, Institute of Psychiatry, Psychology and
Neuroscience, King's College London, De Crespigny Park, London, UK

Running title

Validity IPAQ in first-episode psychosis

Acknowledgements

The authors wish to thank Miss Amber De Herdt and Miss Kim Sweers for their assistance in the data collection.

*Corresponding author: Tervuursevest 101, 3001 Leuven, Belgium. Tel.: +32 2 758 05 11; Fax: +32 2 759 9879. Email: davy.vancampfort@kuleuven.be

Abstract

Aim: The International Physical Activity Questionnaire (IPAQ) is a self-report tool commonly used in mental health care settings to assess physical activity. However, its validity has not yet been investigated in first episode-psychosis (FEP). The aim of this study was to examine the concurrent validity of the IPAQ compared with an objective real-life measure, the Sensewear Armband (SWA) in assessing moderate and vigorous physical activity (MVPA) in people with FEP. Secondary aim was to explore whether there are differences in correlates of the IPAQ versus SWA scores.

Methods: Nineteen outpatients with FEP (15 men; 24.4±5.1years) wore a SWA for 5 full consecutive days, subsequently completed the IPAQ, performed a maximal cardiorespiratory fitness test and were assessed with the Positive and Negative Syndrome Scale (PANSS).

Results: There was no significant correlation between time spent in MVPA according to the IPAQ and SWA. In contrast with SWA scores, there were no significant associations between IPAQ scores and cardiorespiratory fitness levels. No correlations with PANSS scores were observed in both measures.

Conclusions: The current results suggest that the IPAQ should be used with caution when assessing levels of MVPA in FEP. More accurate methods of measuring physical activity are needed in this population.

Keywords: physical activity; exercise; fitness; psychosis

INTRODUCTION

Physical activity improves symptomatic, neurocognitive, cardiorespiratory and metabolic outcomes in people with first-episode psychosis (FEP) ¹⁻⁴. Current guidelines recommend that patients with psychosis should aim to be physically active for at least 150 minutes per week at a moderate intensity ⁵. However, assisting people with FEP to initiate and maintain an active lifestyle remains a key challenge for treatment services ⁶⁻⁸. There is a paucity of published data on physical activity levels in FEP, with a recent meta-analysis only identifying two previous studies ⁹. Valid methods for assessing physical activity levels among young people with psychosis is warranted ^{10,11}. Objective methods for measuring physical activity such as accelerometers are considered to offer more precise estimates of physical activity than self-reported levels as they remove many of the issues of recall and response bias ⁶. Despite these advantages, objective assessment tools are often costly and time-consuming rendering them difficult to use routinely in mental health care settings (Vancampfort et al., 2015a). In order to be able to define clear treatment goals and to assist patients in adopting and maintaining an active lifestyle, it is also pertinent to determine the context in which physical activity occurs ¹² and pedometers or accelerometers do not provide such information (e.g. active transportation, housework, leisure time activity and sports). Questionnaires are a more feasible alternative, are more cost-effective and can provide information about the context in which patients are physically active and in which they are not ¹³. The International Physical Activity Questionnaire (IPAQ) ¹⁴ is the most widely used self-report tool, and is also commonly used in mental health care settings ¹⁵. The IPAQ was designed to assess levels of habitual physical activity in individuals ranging from young to middle-aged (i.e. 15–69 years old). In addition, there are different forms of the IPAQ based on several variations, which include the length of questionnaire (i.e. short or long form), reference period (i.e. last 7 days or usual week) and mode of administration (i.e. self-report or interviewer-based). It has previously been demonstrated in people with schizophrenia that the IPAQ exhibits measurement properties that are comparable to those reported in the general population and therefore can be considered as an appropriate surveillance tool to assess levels of physical activity in these patients ^{16,17}.

To the best of our knowledge, although the IPAQ has been used previously to assess physical activity levels in people with FEP ^{1,2}, validity data pertaining to the use of the IPAQ in young people with FEP is lacking. In order to evaluate its concurrent validity, the IPAQ can be compared with physical activity output from objective measures. Therefore, the aim of this study was to examine the concurrent

validity of time spent in moderate and vigorous physical activity as recorded with the IPAQ, compared with an accelerometer in outpatients with FEP. The secondary aim was to assess whether volume of moderate and vigorous physical activity as assessed with the IPAQ, was differently associated with mental and physical health variables compared with objectively assessed time spent in moderate and vigorous physical activity.

METHODS

Participants and procedure

Over a 9-month period, outpatients of the UPC KU Leuven campus Kortenberg in Belgium, experiencing a FEP¹⁸ were invited to participate. Participants needed to be stable on psychotropic medication for at least 4 weeks. Somatic exclusion criteria included evidence of severe cardiovascular, neuromuscular and endocrine disorders which prevented participants being physically active as per usual. All participants wore an SWA for 5 consecutive days. At the end of day 5, participants completed the International Physical Activity Questionnaire¹⁴. To maximise adherence to the study protocol, on the third day of monitoring participants received a follow-up telephone call to resolve any issues or concerns about the study. The study procedure was approved by the Scientific and Ethical Committee of the UPC KU Leuven, campus Kortenberg, Belgium and conducted in accordance with the principles of the Declaration of Helsinki. All participants gave informed written consent. There was no compensation for participation in the study.

Objective physical activity assessment: Sensewear Armband (SWA)

The SWA is worn over the right arm triceps muscle and assesses minute to minute movement through multiple sensors, namely a two-axis accelerometer and sensors measuring heat flux, galvanic skin and near body-temperature. Data are combined with gender, age, body weight and height, to estimate active energy expenditure using algorithms developed by the manufacturer (SenseWear Professional software, version 7.0). Data were collected for 5 full days (3 week days and 2 weekend days). Daily average time spent in moderate (MPA) (3-6METs) and vigorous physical activity (VPA) (≥ 6 METs) were calculated from all minutes during waking hours. We only included physical activities performed in bouts of at least 10min. In accordance with a previous validity study in people with chronic schizophrenia¹⁷, we did not assess time spent in light physical activity (< 3 MET). Although time spent in light physical activity can be considered as a measure-of-proxy for time spent walking, this can be misleading or erroneous, as for example household chores might also be executed at low intensity. In accordance with previous research¹⁹, data were accepted when the average wear time was at least 1368 minutes per day (95% of a 24-hour bout).

Subjective physical activity assessment: International Physical Activity Questionnaire (IPAQ) - short version

A structured format (short version) that requires participants to recall activities for each of the last seven preceding days was used (self-report) ¹⁴. We included the reported time spent in moderate and vigorous physical activity performed in bouts of at least 10min (min/day). The walking item was not included. In this study, the questionnaire was not interviewer-administered.

Cardiorespiratory fitness test

Cardiorespiratory fitness tests were performed according to internationally accepted standards ²⁰ on a cycle ergometer (Siemens-Elema 380B; Ergometrics 800S, Ergometrics, Bitz, Germany) in an air-conditioned laboratory where the room temperature was regulated at 18-22°C. Participants were asked to cycle at a constant rate of 60 revolutions per minute. The initial workload of 20W was increased by 20W every minute. Blood pressure was measured at rest before the test, with the patient sitting on the bicycle, and every 2 minutes during the test. Heart rate and a 12-lead electrocardiogram (Max Personal Exercise Testing[®], Marquette, WI, USA) were registered continuously. In- and expired gasses were analysed breath-by-breath by means of the Oxycon Pro (Jaeger, Mijnhardt, The Netherlands). The gas analysers and the flow meter were calibrated before each test according to the manufacturer's instructions. VO_2 max values were defined as the 30 seconds average at the highest workload achieved. All participants were asked and encouraged to perform a symptom-limited graded exercise test until exhaustion. In order to define a maximal effect we followed the criteria described by the European Association for Cardiovascular Prevention and Rehabilitation ²¹. A maximal effort was assumed if the cardiorespiratory fitness test was terminated by the patient due to exhaustion, dyspnea, pain or tiredness in the legs and if (a) a peak respiratory exchange ratio (RER) ≥ 1.10 and/or (b) a rating of perceived exertion ≥ 16 on the Borg Scale ²².

Positive and Negative Syndrome Scale (PANSS)

In this study we used the positive (7 items) and negative (7 items) subscales of the PANSS ²³. Each item is scored on a Likert type severity scale ranging from 1 (absent) to 7 (extreme). The subscale scores range from 7 to 49.

Anthropometric measurements

Body weight was measured in light clothing to the nearest 0.1kg using a SECA beam balance scale, and height to the nearest 0.1cm using a wall-mounted stadiometer. Waist circumference (WC) was measured to nearest 1cm at the level of the umbilicus and at the end of expiration with the participant upright and his/her hands by the side.

Statistical analyses

Data were assessed for normality using the Shapiro-Wilk test and except for vigorous physical activity assessments found to be normally distributed. Descriptive statistics are reported as mean and standard deviation (SD), and for uniformity reasons also for vigorous physical activity assessments. Associations between the levels of moderate physical activity using the IPAQ and the SWA were assessed using Pearson's correlation coefficients and for vigorous physical activity using Spearman Rho correlation coefficients (as not normally distributed). We also explored associations of IPAQ and SWA scores with demographical and clinical variables using respectively Pearson's correlation coefficients and Spearman Rho correlation coefficients. The significance level was set at 0.05. Statistical analysis was carried out using IBM SPSS statistics software (version 22.0).

RESULTS

Participants

Twenty-four consecutive patients were invited to participate. One patient was excluded due to a locomotor co-morbidity preventing safe participation and one declined (i.e. not interested). Three patients responded on the questionnaire that they were not able to recall their physical activity levels of the past 7 days (which is a valid option in the IPAQ) and were therefore excluded. Nineteen outpatients with FEP completed the study. There were 15 men and 4 women. Four outpatients (21%) smoked (14 ± 5 cigarettes per day). The mean time spent in moderate physical activity was 214.7 ± 156.6 min/day using the IPAQ and 158.7 ± 81.7 min/day when assessed with the SWA. The mean time spent in vigorous physical activity was 6.3 ± 12.6 min/day using the IPAQ and 4.1 ± 5.7 min/day as assessed with the SWA. Seventeen patients were on antipsychotic monotherapy: aripiprazole (n=5), amisulpride (n=3), ziprasidone (n=3), quetiapine (n=3), paliperidone (n=2). There were two patients who were antipsychotic-naïve, of which one was also psychotropic medication-naïve. Six patients took antidepressants (duloxetine (n=2), mirtazapine (n=2), escitalopram (n=2)) and 2 were on lithium carbonate. All participants wore the SWA for at least 1368 minutes per day (95% of a 24-hour bout) and could be included in the analysis. Mean and SD for other demographical and clinical variables are presented in Table 1.

Associations with and comparisons between the IPAQ and SWA assessments

The correlation between mean time (min/day) spent in moderate physical activity recorded with the SWA and the IPAQ was 0.29 ($P=0.23$) while the correlation spent in vigorous physical activity assessed with the SWA and the IPAQ was -0.11 ($P=0.66$). Compared with the SWA data for physical activity accumulated in bouts of at least 10min, the IPAQ was found to overestimate moderate and vigorous physical activity by 35% and 54% respectively.

Associations of the IPAQ and SWA scores with demographical and clinical variables

As can be seen in Table 1, there were no significant correlations for IPAQ and SWA scores with age, body mass index, waist and PANSS scores. In contrast, better cardiorespiratory fitness levels were associated with higher physical activity levels as assessed with SWA. No associations were found between cardiorespiratory fitness levels and IPAQ scores.

[Insert Table 1 about here]

DISCUSSION

General findings

To the best of the authors' knowledge, this is the first study to examine the concurrent validity of the IPAQ in people with FEP. In sum, we found no significant correlations between time spent in moderate and vigorous physical activity as assessed with the IPAQ and the SWA. Moreover, 3 participants (13.6%) were unable to recall how much physical activity they had engaged in over the previous week and were excluded from the analysis. Thus, our preliminary data suggests that the IPAQ is not a suitable method for monitoring moderate to vigorous physical activity among people with FEP in clinical practice.

There is an imperative to actively include the monitoring of physical activity in the management of FEP. However, based on our data the suitability and accuracy of the IPAQ is questionable for this purpose. As the original objective of the IPAQ was to provide a self-report measure of physical activity that was suitable for assessing general population levels of physical activity across multiple countries¹⁴, the assessment of physical activity levels in clinical settings was not an explicit original goal. In the current study, outpatients with FEP overestimated their physical activity levels by between 35% to 50%, which is however in line with previous research in young men from the general population with poor physical fitness²⁴. A potential reason for this overestimation might be that people with FEP have difficulty with accurately recalling their physical activity levels. Cognitive deficits have been noted among people in the at risk mental state^{25, 26} and those experiencing FEP²⁷ and sedentary behavior and low physical activity have been associated with worse cognition among those with established schizophrenia²⁸. Second, people with FEP are significantly less physically active and have a lower cardiorespiratory fitness versus matched controls^{29, 30} and may have difficulties in accurately recalling the intensity of various physical activities and therefore may perceive light intensity physical activities (≤ 3 MET) already as moderate. The IPAQ defines moderate intensity physical activity broadly as an activity causing somewhat harder breathing¹⁴. In physically inactive and unfit patients, light intensity physical activity might cause people to start already breathing somewhat harder. Third, people with FEP may already have elevated levels of cardio-metabolic abnormalities³¹⁻³³, which is associated with cognitive impairments³⁴ and which can make recall less accurate and therefore influence perceived exertion in actual physical activities.

Overall, our data suggest that IPAQ has questionable validity and should be used with caution when assessing moderate to vigorous intensity physical activity in clinical settings, among outpatients with FEP.

Limitations and future research

While research into physical activity measurement is associated with broad methodological limitations³⁵, the current data should be interpreted in light of some specific limitations. Firstly, while the IPAQ assesses physical activity within a 7-day framework, the Sensewear was only worn in 5 of these 7 days. Second, we did not perform an a-priori power calculation and the current data are based on a small sample of mainly male outpatients with FEP. This limits the generalizability of the current findings. Nevertheless, our results suggest that more work is needed to address the identified problems of the IPAQ when used in outpatients with FEP. There is a need for a clinical physical activity instrument that validly captures physical activity in this population. Therefore, one of the most important challenges in physical activity research in people with FEP is to develop a low cost, easy to use, reliable and valid physical activity questionnaire that captures structural moderate and vigorous physical activity. Such a reliable and valid questionnaire is necessary for standardization across studies and clinical practices and would permit meaningful comparisons of values, for example between disease stages. Another option is to assess the feasibility and validity of the experience sampling method which represents a valuable way of assessing physical activity in real world settings and across time. The experience sampling method requires participants to complete self-assessment questions when prompted by an electronic device (e.g. a mobile phone) at different times of the day, in real world settings. This methodology is flourishing in research examining mental illness³⁶. Because nowadays most people with psychosis carry a mobile phone most of the time and the vast majority is in favor of using mobile phones to enhance contact with services and support self-management³⁷, they have the potential to enhance recall of physical activity by frequent prompting and limiting the time lag between the behavior and data collection. This may reduce information bias and increase compliance. Finally, further research is required to explore whether compared with other questionnaires and objective assessment tools the IPAQ is sensitive to monitor intervention-related changes in physical activity.

REFERENCES

1. Firth J, Carney R, Elliott R, et al. Exercise as an intervention for first-episode psychosis: a feasibility study. *Early intervention in psychiatry*. 2016.
2. Curtis J, Watkins A, Rosenbaum S, et al. Evaluating an individualized lifestyle and life skills intervention to prevent antipsychotic-induced weight gain in first-episode psychosis. *Early intervention in Psychiatry*. 2016; 10: 267-76.
3. Abdel-Baki A, Brazzini-Poisson V, Marois F, Letendre E, Karelis AD. Effects of aerobic interval training on metabolic complications and cardiorespiratory fitness in young adults with psychotic disorders: a pilot study. *Schizophrenia Research*. 2013; 149: 112-5.
4. Lee EH, Hui CL, Chang WC, et al. Impact of physical activity on functioning of patients with first-episode psychosis—A 6months prospective longitudinal study. *Schizophrenia Research*. 2013; 150: 538-41.
5. Vancampfort D, Hert MD, Skjerven LH, et al. International Organization of Physical Therapy in Mental Health consensus on physical activity within multidisciplinary rehabilitation programmes for minimising cardio-metabolic risk in patients with schizophrenia. *Disability and Rehabilitation*. 2012; 34: 1-12.
6. Rosenbaum S, Lederman O, Stubbs B, Vancampfort D, Stanton R, Ward PB. How can we increase physical activity and exercise among youth experiencing first-episode psychosis? A systematic review of intervention variables. *Early Intervention in Psychiatry*. 2016; 10(5): 435-40.
7. Vancampfort D, De Hert M, Broderick J, et al. Is autonomous motivation the key to maintaining an active lifestyle in first-episode psychosis? *Early Intervention in Psychiatry*. 2016; doi: 10.1111/eip.12373.
8. Firth J, Carney R, French P, Elliott R, Cotter J, Yung AR. Long-term maintenance and effects of exercise in early psychosis. *Early Intervention in Psychiatry*. 2016; doi: 10.1111/eip.12365.
9. Stubbs B, Firth J, Berry A, et al. How much physical activity do people with schizophrenia engage in? A systematic review, comparative meta-analysis and meta-regression. *Schizophrenia Research*. 2016; 176(2-3): 431-40.
10. Vancampfort D, Rosenbaum S, Probst M, et al. What are the top 10 physical activity research questions in schizophrenia? *Disability and Rehabilitation*. 2016; 38: 2235-43.
11. Carney R, Bradshaw T, Yung AR. Monitoring of physical health in services for young people at ultra-high risk of psychosis. *Early intervention in Psychiatry*. 2015; doi: 10.1111/eip.12288.

12. Vancampfort D, De Hert M, De Herdt A, et al. Associations between physical activity and the built environment in patients with schizophrenia: a multi-centre study. *General Hospital Psychiatry*. 2013; 35: 653-8.
13. Vanhees L, Lefevre J, Philippaerts R, et al. How to assess physical activity? How to assess physical fitness? *European Journal of Cardiovascular Prevention & Rehabilitation*. 2005; 12: 102-14.
14. Craig C, Marshall A, Sjostrom M, et al. International physical activity questionnaire: 12-country reliability and validity. *Medicine & Science in Sports & Exercise*. 2003; 35: 1381-95.
15. Soundy A, Roskell C, Stubbs B, Vancampfort D. Selection, use and psychometric properties of physical activity measures to assess individuals with severe mental illness: a narrative synthesis. *Archives of Psychiatric Nursing*. 2014; 28: 135-51.
16. Faulkner G, Cohn T, Remington G. Validation of a physical activity assessment tool for individuals with schizophrenia. *Schizophrenia Research*. 2006; 82: 225-31.
17. Duncan MJ, Arbour-Nicitopoulos K, Subramanieapillai M, Remington G, Faulkner G. Revisiting the International Physical Activity Questionnaire (IPAQ): Assessing physical activity among individuals with schizophrenia. *Schizophrenia Research*. 2017; 179: 2-7.
18. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*. American Psychiatric Association, Washington, DC; 2013.
19. Vancampfort D, Wyckaert S, Sienaert P, et al. Concurrent validity of the international physical activity questionnaire in outpatients with bipolar disorder: Comparison with the Sensewear armband. *Psychiatry Research*. 2016; 237:122-6.
20. Gibbons RJ, Balady GJ, Bricker JT, et al. ACC/AHA 2002 guideline update for exercise testing: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). *Journal of the American College of Cardiology*. 2002; 40: 1531-40.
21. Mezzani A, Agostoni P, Cohen-Solal A, et al. Standards for the use of cardiopulmonary exercise testing for the functional evaluation of cardiac patients: a report from the Exercise Physiology Section of the European Association for Cardiovascular Prevention and Rehabilitation. *European Journal of Cardiovascular Prevention & Rehabilitation*. 2009; 16: 249-67.
22. Borg G. *Borg's perceived exertion and pain scales*. Human Kinetics Publishers; 1998.

23. Kay SR, Flszbein A, Opfer LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*. 1987; 13: 261.
24. Fogelholm M, Malmberg J, Suni J, et al. International physical activity questionnaire: validity against fitness. *Medicine and Science in Sports and Exercise*. 2006; 38: 753-60.
25. Hou C-L, Xiang Y-T, Wang Z-L, et al. Cognitive functioning in individuals at ultra-high risk for psychosis, first-degree relatives of patients with psychosis and patients with first-episode schizophrenia. *Schizophrenia Research*. 2016; 174: 71-6.
26. De Herdt A, Wampers M, Vancampfort D, et al. Neurocognition in clinical high risk young adults who did or did not convert to a first schizophrenic psychosis: a meta-analysis. *Schizophrenia Research*. 2013; 149: 48-55.
27. Caldiroli A, Buoli M, Serati M, Cahn W, Altamura AC. General and social cognition in remitted first-episode schizophrenia patients: a comparative study. *European Archives in Psychiatry and Clinical Neurosciences*. 2016; 266: 639-47.
28. Stubbs B, Ku P-W, Chung M-S, Chen L-J. Relationship between objectively measured sedentary behavior and cognitive performance in patients with schizophrenia vs controls. *Schizophrenia Bulletin*. 2017; 43(3): 566-74.
29. Nyboe L, Vestergaard C, Moeller M, Lund H, Videbech P. Metabolic syndrome and aerobic fitness in patients with first-episode schizophrenia, including a 1-year follow-up. *Schizophrenia Research*. 2015; 168: 381-7.
30. Nyboe L, Moeller MK, Vestergaard CH, Lund H, Videbech P. Physical activity and anomalous bodily experiences in patients with first-episode schizophrenia. *Nordic Journal of Psychiatry*. 2016: 1-7.
31. Mitchell AJ, Vancampfort D, De Herdt A, Yu W, De Hert M. Is the prevalence of metabolic syndrome and metabolic abnormalities increased in early schizophrenia? A comparative meta-analysis of first episode, untreated and treated patients. *Schizophrenia Bulletin*. 2013;39(2):295-305.
32. Vancampfort D, Wampers M, Mitchell A, Correll C, Herdt A, Probst M, et al. A meta-analysis of cardio-metabolic abnormalities in drug naive, first-episode and multi-episode patients with schizophrenia versus general population controls. *World Psychiatry*. 2013;12(3):240-50.
33. Correll CU, Robinson DG, Schooler NR, et al. Cardiometabolic risk in patients with first-episode schizophrenia spectrum disorders: Baseline results from the raise-etp study. *JAMA Psychiatry*. 2014.

34. Bora E, Akdede B, Alptekin K. The relationship between cognitive impairment in schizophrenia and metabolic syndrome: a systematic review and meta-analysis. *Psychological Medicine*. 2016: 1-11.
35. Kelly P, Fitzsimons C, Baker G. Should we reframe how we think about physical activity and sedentary behaviour measurement? Validity and reliability reconsidered. *International Journal of Behavioral Nutrition and Physical Activity*. 2016; 13: 32.
36. Myin-Germeys I, Oorschot M, Collip D, Lataster J, Delespaul P, van Os J. Experience sampling research in psychopathology: opening the black box of daily life. *Psychological Medicine*. 2009; 39: 1533.
37. Firth J, Cotter J, Torous J, Bucci S, Firth JA, Yung AR. Mobile phone ownership and endorsement of "mHealth" among people with psychosis: a meta-analysis of cross-sectional studies. *Schizophrenia Bulletin*. 2016; 42(2): 448-55.

TABLE 1. Pearson's and Spearman Rho correlation coefficients° of demographical and clinical variables in people with first-episode psychosis (n=19)

Variables	Mean±SD	r with IPAQ	P IPAQ	r with SWA	P SWA	r with IPAQ	P IPAQ	r with SWA	P SWA
		moderate	moderate	moderate	moderate	vigorous	vigorous	vigorous	vigorous
Age (years)	24.4±5.1	-0.04	0.88	-0.07	0.78	0.14	0.56	0.02	0.93
Body mass index	24.8±3.8	0.05	0.84	-0.34	0.15	0.16	0.50	-0.33	0.17
Waist (cm)	84.4±10.6	-0.10	0.78	-0.17	0.47	-0.12	0.63	-0.08	0.73
VO2 max (ml/min/kg)	30.3±7.4	0.27	0.25	0.68	0.001*	-0.22	0.36	0.49	0.032*
PANSS positive score	7.9±1.1	-0.008	0.97	0.02	0.94	-0.31	0.20	-0.07	0.79
PANSS negative score	9.6±3.1	0.06	0.80	-0.05	0.84	0.11	0.66	-0.04	0.86
Antipsychotic dose (mg/d)	273.1±120.6	0.11	0.66	0.31	0.22	-0.23	0.37	0.07	0.78

*Significance is set at P<0.05. °Pearson's correlation coefficients are calculated for the moderate physical activity levels, Spearman Rho correlation coefficients for the vigorous physical activity levels. IPAQ= International Physical Activity Questionnaire; SWA: Body Sensewear armband, VO2 max= maximum oxygen uptake.