



King's Research Portal

DOI:

[10.1016/j.jacc.2017.10.093](https://doi.org/10.1016/j.jacc.2017.10.093)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

O'Gallagher, K., Khan, F., Omar, S. A., Kalra, S., Danson, E., Cabaco, A. R., Martin, K., Melikian, N., Shah, A. M., & Webb, A. J. (2018). Inorganic Nitrite Selectively Dilates Epicardial Coronary Arteries. *Journal of the American College of Cardiology*, 71(3), 363-364. <https://doi.org/10.1016/j.jacc.2017.10.093>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Letter: Inorganic Nitrite Selectively Dilates Epicardial Coronary Arteries

Kevin O’Gallagher MBBS^{1,2,3}, Faisal Khan MBBS^{1,2,3}, Sami A. Omar MBBS PhD^{1,3},
Sundeep Kalra MA MBBS^{1,2}, Edward Danson BM BCh, DPhil^{1,2}, Ana R Cabaco MSc^{1,2},
Katherine Martin MSc^{1,2}, Narbeh Melikian MD^{1,2}, *Ajay M. Shah MD^{1,2}, *Andrew J. Webb
MBBS PhD^{1,3}

Author Affiliations:

1. King’s College London British Heart Foundation Centre, School of Cardiovascular Medicine and Sciences,
2. King’s College Hospital NHS Foundation Trust, London, UK
3. Guy’s and St Thomas’ NHS Foundation Trust, London, UK

*, Joint senior authors.

Corresponding author:

Dr Andrew J Webb,

Senior Lecturer/Honorary Consultant in Cardiovascular Clinical Pharmacology,

King’s College London British Heart Foundation Centre,

Cardiovascular Division,

Department of Clinical Pharmacology,

St.Thomas’ Hospital, London, SE1 7EH, UK.

Tel: 02071884602

Email: andrew.1.webb@kcl.ac.uk

Sources of Funding, Relationship with industry and financial disclosure: The authors’ work was supported by the British Heart Foundation and the Department of Health via a

National Institute for Health Research (NIHR) Biomedical Research Centre award to Guy's & St Thomas' NHS Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust. AJW is a share holder of HeartBeet Ltd, Suffolk, UK who manufacture and supply beetroot juice (active nitrate-containing and placebo nitrate-depleted); though not used in this study.

Inorganic nitrite (NO_2^-), as sodium nitrite, or derived from dietary nitrate found in green leafy vegetables and beetroot, has potential as a tolerance-free therapy in Heart Failure with Preserved Ejection Fraction (HFpEF)(1) including pulmonary hypertension-associated HFpEF (PH-HFpEF) (2). Nitrite affects small and large arteries and veins. Previous studies demonstrated hypoxia (versus normoxia) enhanced nitrite-induced dilatation by 30-50% in small resistance arterioles, but not veins (since venous blood is already desaturated) (3). We recently demonstrated that nitrite selectively dilates peripheral conduit arteries in a normoxia-dependent/hypoxia-inhibited manner, and selectively lowers aortic blood pressure (afterload)(4). We confirmed this latter effect with 6 months' dietary nitrate (beetroot juice) in 126 patients(5). Nitrite has similar selectivity as nitroglycerin (GTN) for conduit, versus small resistance arterioles(4). We therefore investigated whether nitrite also selectively vasodilates epicardial coronary (conduit) arteries.

Following institutional and ethical approvals and informed consent, we recruited patients undergoing diagnostic coronary angiography, including those with an angiographically-normal coronary artery/without previous coronary intervention.

A 0.014-inch coronary Doppler wire (Flow Wire, Phillips Volcano, USA) was positioned within the proximal artery and average peak velocity (APV) measured following five-minute infusions of 0.9% saline (baseline), low-dose sodium nitrite ($2.6\mu\text{mol}/\text{min}$), high-dose sodium nitrite ($26\mu\text{mol}/\text{min}$), and GTN ($1\mu\text{g}/\text{min}$). Angiographic images were obtained simultaneously. Quantitative Coronary Analysis (QCA) was performed off-line with measurements of vessel diameter (d) in the 5mm segment distal to the tip of the Doppler wire. Coronary blood flow (CBF, cm^3/s) was calculated by: $\text{CBF} = (\text{APV}/2) \times \text{cross-}$

sectional area. Coronary resistance ($\text{mmHg}/\text{cm}^3/\text{s}$) was calculated as mean arterial pressure divided by coronary blood flow.

Nine patients (mean age 56 years, 5 male, 78% hypertensive) received both doses of nitrite, and GTN, and had angiographic and Doppler flow traces of sufficient quality to be included in the analysis. The higher dose of nitrite ($26\mu\text{mol}/\text{min}$) and GTN both significantly dilated the coronary artery compared to baseline: diameter-change (mean [95% CI]): $+8.9\%$ [$+2.0,+15.8$], $p=0.01$ and $+10.8\%$ [$+1.0,+20.5$] $p=0.03$, respectively (**Figure 1A**); with no significance difference between nitrite and GTN ($p=0.45$). Lower dose nitrite ($2.6\mu\text{mol}/\text{min}$) had no significant effect ($+3.9\%$ [$-2.8,+10.6$], $p=0.3$).

GTN increased coronary flow relative to baseline (median [interquartile range]): $+23.9\%$ [$+6.3,+72.7$], $p=0.02$. Nitrite ($2.6\mu\text{mol}/\text{min}$ and $26\mu\text{mol}/\text{min}$) lacked significant effect on coronary flow: $+12.0\%$ [$-10.2,+24.1$] $p>0.99$ and $+16.9\%$ [$+3.2,+64.4$] $p>0.99$ respectively. This was despite relatively high estimated intracoronary nitrite concentrations 73.7 [$57.8, 154.9$] μM (median [IQR]) for nitrite ($2.6\mu\text{mol}/\text{min}$) and 705 [$555, 1179$] μM (median [IQR]) for nitrite ($26\mu\text{mol}/\text{min}$).

GTN also lowered coronary resistance relative to baseline (mean [95% CI]): -29.3% [$-52.8,-5.9$] $p=0.02$, and versus nitrite ($26\mu\text{mol}/\text{min}$): -10.4% [$-18.3,-2.4$] $p=0.01$. By contrast, nitrite ($2.6\mu\text{mol}/\text{min}$ and $26\mu\text{mol}/\text{min}$) had no significant effect on coronary resistance: -11.1% [$-36.7,+14.5$] $p=0.52$, and -19.0% [$-43.4,+5.4$] $p=0.13$, respectively (**Figure 1B**).

There were no significant overall differences in central systolic, diastolic, or mean blood pressure with any of the interventions versus baseline. Heart rate changed -3.1 bpm [-0.2,-6.1], $p=0.02$, with nitrite (2.6 $\mu\text{mol}/\text{min}$), but not with nitrite (26 $\mu\text{mol}/\text{min}$) or GTN.

In summary, this is the first data to show that inorganic nitrite dilates epicardial coronary arteries; moreover, this was selective (and dose-dependent). Whilst the results should be interpreted with caution, given the small sample size, the indirect measures of coronary flow, and the trend towards a change in coronary flow/resistance, they support our findings in the peripheral circulation(4) of nitrite's physiological action as a normoxia-dependent selective conduit artery dilator over resistance arteries, even in the metabolically-active myocardium. These findings also have important implications for our understanding of the potential beneficial cardiovascular properties of nitrite derived from dietary nitrate sources such as beetroot and green leafy vegetables: nitrite acts principally on large, rather than small vessels in the heart.

Figure List


Figure 1. Percentage change in: **A** coronary artery diameter, and **B** coronary resistance. Nitrite (2.6) and Nitrite (26) denote nitrite (2.6 $\mu\text{mol}/\text{min}$) and nitrite (26 $\mu\text{mol}/\text{min}$), respectively. Data shown as Mean \pm SEM. Significance shown as * $p<0.05$ vs baseline, † $p<0.05$ vs nitrite (2.6 $\mu\text{mol}/\text{min}$), ‡ $p<0.05$ vs nitrite (26 $\mu\text{mol}/\text{min}$)

References

1. Borlaug BA, Koepp KE, Melenovsky V. Sodium Nitrite Improves Exercise Hemodynamics and Ventricular Performance in Heart Failure With Preserved Ejection Fraction. *J Am Coll Cardiol* 2015;66:1672-82.
2. Simon MA, Vanderpool RR, Nouraie M et al. Acute hemodynamic effects of inhaled sodium nitrite in pulmonary hypertension associated with heart failure with preserved ejection fraction. *JCI Insight* 2016;1:e89620.
3. Maher AR, Milsom AB, Gunaruwan P et al. Hypoxic modulation of exogenous nitrite-induced vasodilation in humans. *Circulation* 2008;117:670-7.
4. Omar SA, Fok H, Tilgner KD et al. Paradoxical normoxia-dependent selective actions of inorganic nitrite in human muscular conduit arteries and related selective actions on central blood pressures. *Circulation* 2015;131:381-9; discussion 389.
5. Mills CE GV, Faconti L, et al. Reducing arterial stiffness independently of blood pressure? The randomized, factorial ‘VaSera’ trial in people with or at risk of Type 2 Diabetes. *J Am Coll Cardiol* 2017;70:1683-1684.

A

Coronary Diameter



B

Coronary Resistance

