



King's Research Portal

DOI:

[10.1177/2055207617743354](https://doi.org/10.1177/2055207617743354)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Milward, J., Drummond, C., Fincham-Campbell, S., & Deluca, P. (2018). What makes online substance-use interventions engaging? A systematic review and narrative synthesis. *Digital Health*, 4. Advance online publication. <https://doi.org/10.1177/2055207617743354>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

What makes online substance-use interventions engaging? A systematic review and narrative synthesis

Joanna Milward, Colin Drummond, Stephanie Fincham-Campbell and Paolo Deluca

Digital Health
Volume 4: 1–25
© The Author(s) 2018
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/2055207617743354
journals.sagepub.com/home/dhj


Abstract

Background: Online substance-use interventions are effective in producing reductions in harmful-use. However, low user engagement rates with online interventions reduces overall effectiveness of interventions. Identifying optimal strategies with which to engage users with online substance-use interventions may improve usage rates and subsequent effectiveness.

Objectives: (1) To identify the most prevalent engagement promoting strategies utilised to increase use of online substance-use interventions. (2) To determine whether the identified engagement promoting strategies increased said use of online substance-use interventions.

Review methods: The reviewed followed Cochrane methodology. Databases were searched for online substance-use interventions and engagement promoting strategies limited by study type (randomised controlled trial). Due to heterogeneity between engagement promoting strategies and engagement outcomes, meta-analytic techniques were not possible. Narrative synthesis methods were used.

Results: Fifteen studies were included. Five different engagement promoting strategies were identified: (1) tailoring; (2) delivery strategies; (3) incentives; (4) reminders; (5) social support. The most frequently reported engagement promoting strategies was tailoring (47% of studies), followed by reminders and social support (40% of studies) and delivery strategies (33% of studies). The narrative synthesis demonstrated that tailoring, multimedia delivery of content and reminders are potential techniques for promoting engagement. The evidence for social support was inconclusive and negative for incentives.

Conclusions: This review was the first to examine engagement promoting strategies in solely online substance-use interventions. Three strategies were identified that may be integral in promoting engagement with online substance-use interventions. However, the small number of eligible extracted studies, inconsistent reporting of engagement outcomes and diversity of engagement features prevent firmer conclusions. More high-quality trials examining engagement are required.

Keywords

Systematic review, internet, telemedicine, patient adherence, engagement

Submission date: 11 April 2017; Acceptance date: 27 October 2017

Introduction

Online interventions have been developed for a range of substance-use issues including alcohol reduction, tobacco and cannabis smoking cessation.^{1–3} Typically, they are delivered or supported via the internet and

Addictions Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

Corresponding author:

Joanna Milward, Addictions Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, Addiction Sciences Building, 4 Windsor Walk, London, SE5 8BB, UK.
Email: Joanna.milward@kcl.ac.uk

include web-based interventions such as internet-operated therapeutic software and online counselling and therapy. Meta-analyses have demonstrated effectiveness of online and computer-based substance-use interventions (SUIs) on behaviour change;^{4–7} however, engagement with the interventions remains a challenge.^{8,9}

Engagement broadly refers to how the user interacts with the technology, typically measured by web-metric usage data. Engagement related behaviours include accessing the intervention website for the first time, using the intervention website and its features, as well as revisiting the website.¹⁰ Issues such as low login rates and limited use of intervention features are consistently reported in the literature with only a minority of participants returning to the online intervention.^{11–13} Studies suggest a dose–response relationship between the level of a user’s engagement and the effectiveness of an online health intervention.^{14–17} Whilst this association may also be due to reverse causality, such as when a user experiences better outcomes they are more likely to engage, it is plausible that engagement increases effectiveness and online intervention components are important strategies to improve rates of engagement.¹⁰ Additional individual and environmental level factors have also been hypothesised to influence engagement;^{18–20} however, this review is focusing solely on technology-based intervention strategies.

Previous reviews have examined engagement strategies predominantly for internet-based health behaviour change interventions. An early systematic review by Crutzen et al.¹⁰ examined strategies that facilitated exposure to internet-delivered health behaviour change interventions for adolescents. Eight out of 17 of the included studies were targeting substance-use, utilising strategies of targeted communication, customisation of information, tailoring, professional and peer support, interactivity, reminders and incentives. Two studies examining SUIs examined the effectiveness of engagement promoting strategies (EPSs) compared with a control, with neither study reporting significant differences for interactive or tailored features. By the term ‘tailored’, we refer to a process whereby intervention components are individualised to specific characteristics or needs of users, such as tailoring text messages to gender or drinking behaviours.²¹ When including all types of health behaviour change interventions the authors reported that multi-component strategies were the most effective. Brouwer et al.⁹ also examined exposure to internet-delivered health promotion interventions with qualitative descriptive analyses. Twenty-seven out of 64 studies in the Brouwer et al.⁹ study were targeting substance-use including EPSs of interactivity, peer and counsellor support, email/phone contact and frequent updates. The authors

reported that the most effective strategies peer and counsellor support and email or phone contact.

Schubart et al.²² examined internet behavioural interventions for chronic health conditions in adults, using a positive deviance approach, where engagement strategies were compared between five studies with the highest, and five with the lowest attrition rates. However, none of the three studies included in the Schubart et al.²² study including SUIs were included in this comparison. Overall, the authors reported that tailoring, social support and updated material contributed to reducing non-engagement. Finally, Alkhaldi et al.²³ conducted a review on the use of prompts to improve engagement with digital interventions and reported borderline positive results for prompts compared with no strategy.

Previous reviews have examined EPSs across a broad range of healthcare issues;^{9,10,22,23} however, no review has focused only on online SUIs. With the exception of Alkhaldi et al.,²³ who focused on prompts for both online and offline interventions, no systematic review has been conducted on EPSs for online interventions since 2011. Furthermore, with the fast-paced development of new technologies, no review has included the use of smartphone or tablet based interventions. Finally, there is a lack of reviews considering only high quality randomised controlled trials (RCTs). As healthcare interventions moves towards mhealth, this review addresses the gap in the literature by examining the most up-to-date EPSs, considering only the studies of rigorous RCT design, to inform the development of future interventions.

The aims of this systematic review were: (1) to identify the most prevalent EPSs utilised to increase use of online SUIs; (2) to determine whether the identified EPSs increased engagement with online SUIs.

Method

Design

Systematic review of RCTs, following Cochrane methodology and Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.²⁴ The review was registered with PROSPERO (CRD42016038874).

Definitions

Online health interventions. This review targeted online health interventions delivered or supported via the internet.²⁵ This included but was not limited to smartphone apps, intervention-based websites, and online-delivered computer and tablet-based interventions. The definition of online SUIs also includes online

interventions which can be used offline (e.g. apps). Online health interventions are differentiated from digital interventions, which are not dependent upon delivery or support from the internet, such as an analogue pedometer or a CDROM based computer program.²⁶ Online SUIs were chosen because they represent the most up-to-date technology which is likely to supersede analogue (offline) interventions in the future.

Engagement. Engagement was defined by a user's interaction with the intervention. This may be visiting a particular aspect of the online intervention or completing a task such as filling out an online diary. Engagement also included how long or how often the participant used the online intervention. Previous research examining engagement has reported a range of engagement outcomes including: number of logins to the intervention; time-spent on the intervention; specific web-sessions opened and tasks completed.^{17,27–29}

EPSs. EPSs were defined as any method used to promote engagement with online SUIs. This includes prompts, tailoring of the intervention to its users; social support, online community and gamification features. Gamification is defined as the use of gaming components in non-gaming settings.³⁰

Inclusion criteria

The eligibility criteria were selected based on participants, interventions, comparisons, outcomes and study designs. A start date of 1990 was chosen as it corresponds to the time period when the internet became available.³¹

Participants. All individuals accessing an online SUI, including, but not limited to, individuals accessing online interventions targeting alcohol reduction and smoking cessation. No age limit was imposed.

Types of intervention. EPSs delivered as a component of an online SUIs designed to increase use of said intervention. These were defined from a scoping review of the literature as prompts, tailoring, social support, online community and gamification features.

Intervention delivery format. Online SUIs as per the definition above including, but not limited to: mobile phones; tablets; web-based interventions; wearable technology with online capability; push notifications including text-messaging; social networks; online social support and social media.

Comparisons. (1) Online SUIs offering no engagement promoting strategies such as tailoring versus no

tailoring; (2) online SUIs offering different levels of dosage of EPS such as low versus high dose of tailoring; (3) online SUIs offering alternative EPSs such as prompts versus tailoring.

Outcome. The main outcome was engagement with the online SUIs (measured as primary and secondary outcomes in included trials). This is typically reported as means of: number of logins; time spent on intervention (e.g. minutes); number of pages viewed; number of sessions completed; number of web-sessions opened; number of features used and tasks achieved.

Exclusion

The following were excluded: (1) trials examining CDROM and computer-based interventions which do not function in an online capacity; (2) trials where attrition from the trial (defined as drop-out attrition or loss to follow-up) could not be disentangled from attrition from the online intervention (defined as non-usage attrition); (3) studies which compared EPSs between different online SUIs; (4) studies which compared EPSs between an online intervention and offline health intervention. For (3) such studies were excluded as the effects of EPSs cannot be disentangled from the active ingredients of the behaviour change intervention.

Data sources and search methods

The search strategy combined online SUIs with engagement outcomes and EPSs limited by study type (RCT). The strategy was based on previously published systematic reviews on online health interventions and engagement with digital interventions^{4,10,32} and, in the case where systematic reviews had not been conducted on a specific topic, such as gamification, the search terms were based on the existing literature.³³ The Medline thesaurus Medical Subject Headings (MESH) terms were identified, supplemented by keyword searches and relevant filters. The data extraction tool was tested by one reviewer (JM) to determine whether 9 studies^{17,28,29,34–38} identified from a hand-search of the literature met the inclusion criteria. The Medline search terms are listed in Appendix 1.

The following databases were searched: Medline, PsychInfo, Embase, CENTRAL, CINAHL, Google Scholar (first 100 citations) and Database of Abstracts of Reviews of Effects (DARE). Unpublished literature was searched from OpenGrey and Index to Theses and Science Citation Index. Reference lists of included studies were hand searched and papers citing key papers screened.

Data collection and analysis

All studies identified were downloaded into EndNote X7 and duplicates removed. Two members of the review team independently screened article titles and abstracts (JM and SFC) in May and June 2016. Full texts were retrieved for relevant studies and eligibility assessed. If any discrepancies arose these were discussed by the two reviewers until a consensus was reached. Data from eligible papers was extracted into a template adapted from the Cochrane Collaboration's tool for assessing risk of bias.³⁹ The extraction form was piloted using a representative sample of the studies by one reviewer (JM).

Data synthesis

Reporting was based on PRISMA guidelines.²⁴ At protocol stage, meta-analytic techniques had been planned; however, they were determined to be inappropriate for comparison of the studies in the current review. This was due to high heterogeneity both in terms of outcomes as well as for type of EPS evaluated. The studies included evaluated a broad range of EPSs, including reminders, incentives and tailoring, and pooling these effects would not have been appropriate. Equally, there was high variability between outcome measures, which again were not appropriate for pooling. Consequently a modified version of narrative synthesis^{40,41} approach was used. Narrative synthesis is recommended by Cochrane when traditional meta-analytic approaches are not appropriate.⁴² As meta-analytic approaches were not appropriate, selection of a single primary outcome was not necessary (if studies reported more than one primary outcome), therefore, all engagement outcomes were reported.

Narrative synthesis adopts a textual approach and aims to provide both a summary of the knowledge-base and a rigorous evaluation, providing a robust interpretative synthesis of the effectiveness of the intervention in question. There were three stages to the modified approach to narrative synthesis: (1) developing a preliminary synthesis; (2) exploration of relationships in the data; (3) assessing the robustness of the synthesis product. A number of different tools and techniques are available for the stages above, depending on the type of studies included. At the beginning of the synthesis process the available tools and techniques were evaluated in terms of their relevance to the data by reviewer JM (see Appendix 2).

Assessment of risk of bias and robustness in included studies

Risk of bias was determined using the Cochrane's Collaboration tool.³⁹ Each study included was

evaluated for risk of bias and was coded as low-risk, high-risk or unclear risk, taking into account random sequence generation, allocation sequence concealment, blinding, incomplete data and selective outcome reporting. Other bias was considered, including selection bias, sample size, power and baseline characteristic differences between groups. A risk of bias summary was generated (see Appendix 2).

Results

Summary of search results

A total of 511 records were found. After removing duplicates 342 records were available for abstract screening. Of these, 48 studies went forward for full-text review. Thirty-three studies were excluded, the most common reason being lack of a suitable comparison group (see Figure 1).

Characteristics of included studies

In total, 15 studies were included. All of the studies were computer, web-based interventions (as opposed to smartphone or tablet). Characteristics of studies are described in Table 1. Eleven studies targeted tobacco-use,^{29,35,43–51} three studies targeted alcohol-use^{2,53,53} and one study targeted cannabis-use.⁵⁴ Two studies used a factorial RCT design^{35,29} and the remaining studies compared conditions with varying EPS types. The content and theoretical framework of the online interventions varied, but at minimum included information and advice for cutting down. Table 1 details the characteristics of the studies.

Quality of included studies

The studies varied in quality and level of risk (see Appendix 2). An overall score of quality was calculated for comparative purposes by summing the 'low risk' scores for each bias category (see Appendix 3). The studies with the least bias were McClure et al.,³⁵ Houston et al.⁴⁵ and Stoddard et al.⁵⁶ These studies reported rigorous methodologies and were of high quality.

Engagement outcomes were measured automatically so there was no blinding of outcome data bias. For missing outcome data, there was low risk of bias overall due to objective measuring. However, one study⁴⁸ excluded 6% of participants who did not log on at all and another⁵⁰ excluded 65 people who spent less than five minutes on the website, which may have introduced an element of bias. Engagement measures were only reported in three protocols⁵⁷ but were pre-specified in the methods of 11 papers.^{29,35,43,45–48,52–54,56}




Figure 1. PRISMA flow diagram.

There was potential bias of insufficient power in all studies as power analyses for engagement were not reported. Two studies^{47,51} had sample sizes of below 100 participants. Self-selection bias was found in the majority of studies, particularly for those who recruited using online and print advertisement.^{2,43,44,46–54}

Research question 1: to identify the most prevalent EPSs utilised to increase use of online SUIs

Five individual EPSs were identified: (1) tailoring; (2) reminders (3); delivery strategies; (4) social support; (5) incentives. An additional category, ‘multi-component’, was created for studies which compared a basic version of an app with an enhanced version. Due to this study design, it was not possible to disentangle the effects of individual components of EPSs on engagement in these studies.

The most frequently reported EPS was tailoring, with seven studies (47%) using this technique.

Ten different sub-types of tailoring techniques were identified (see Table 2). The most frequent sub-type of tailoring was to level of readiness to quit^{43–45,62} (40% of tailoring studies, $n = 4$) and to self-efficacy and barriers to quitting^{29,35,44} (30% of tailoring studies, $n = 3$). This was followed by tailoring to abstinence status,^{2,63} goals and motivations for quitting^{2,29} and testimonials/success stories (20% of tailoring studies, $n = 2$ for each sub-type). All other sub-types were included in only a single study.⁴⁵

Reminders^{29,35,43,45,46,62} and social support^{54,43,45,46,56,62} were used in six studies (40%), delivery strategies^{29,34,35,49,53} in five (33%). All reminders were via email. The most popular delivery strategy techniques were using multimedia content^{34,49} and variations on the timing of delivery of content.^{29,53} Social support was either peer led,^{43,46,56} therapist led,⁵⁴ or a combination of both.^{45,62} Incentives were used in two studies,^{47,51} including rewards contingent on abstinence or engagement. Two studies used a multi-component design.^{43,63}

Table 1. Study characteristics.

Author, year and location	Targeted substance/health behaviour	Study characteristics	Description of DI	n participants	Characteristics of study population	Engagement outcome measure(s)	Reported engagement results
Tailoring							
Danaher et al. 2012 USA	Smokeless tobacco	Design: RCT (three arms) Participants: adult smokers Arms: (1) active control (basic version), (2) enhanced, (3) enhanced + EPS(s); tailoring/social support/reminders/multimedia recruitment; referral from smoking clinic DI format: web-based	Name: MyLastDip Arm 1 (basic version): included step-by-step instructions for best practices for cutting down and a resources section Arm 2 (enhanced version): provided above interactively with multimedia, personalised content; email reminders; goal setting and a web blog	N = 1716 (Enhanced = 857; basic = 859)	% male: 97% % White: 96% Age: 20.8 years, SD = 2.6 Readiness to quit: mean of 9 (SD 2.1) on an 11 point scale	(1) Logins (2) Duration of visit (min)	(1) Participants in enhanced version logged in more compared with basic** (2) Participants in enhanced version had higher duration of visit (min) compared with basic***
Elfeddali et al. 2012 Netherlands	Tobacco	Design: RCT (three arms) Participants: adult smokers aged 18–65 years Arms: (1) basic version (action planning (AP) programme), (2) enhanced version (AP+), (3) control (questionnaire only) EPS(s): tailoring (basic vs. enhanced tailoring) Recruitment: online and print DI format: web-based	Name: Stay Quit for You (SQ4U) Tailored smoking cessation web-based programme using the I-Change Model ^{16b} Arm 1: basic included (AP); (1) tailored feedback before the quit attempt, (2) tailored planning strategy assignments before and after the quit attempt Arm 2: enhanced included above, plus tailored feedback after the quit attempt Arm 3: questionnaire only	N = 1812 (after exclusions) (Basic tailoring: n = 542, enhanced tailoring: n = 622, control: n = 648)	% male: 38% % White: not reported Age: 41 years (11.80) Readiness to quit: not reported	Number of planning assignments completed (six in total)	Participants in enhanced condition (AP+) completed more planning assignments than participants in the basic condition (AP)*
Houston et al. 2015 USA	Tobacco	Design: RCT (two arms) Participants: adult smokeless tobacco users Arms: three arms of increasing complexity EPS(s): interactivity/tailoring/social support/reminders/multimedia recruitment: online and print paid advertisements, radio DI format: web-based	Name: Decide2quit Arm 1: interactive, tailored quit smoking website. Including: motivational messaging tailored to readiness to quit; interactive risk, decisional balance, and cessation barrier calculators; informational resources Arm 2: enhanced version included above, plus a tailored pushed motivational email messaging system Arm 3: enhanced version + included the above ((1) and (2)) plus a social support group and communication with a trained tobacco treatment specialist	N = 900 (control = 299, enhanced = 164, enhanced+ = 437)	% male: 37% % White: 85% Age: 50% were 35–55 years (mean and SD not reported) Readiness to quit: 80% thinking about quitting	(1) Logins	(1) Participants in enhanced+ logged in more than the control*** (2) Enhanced+ vs. enhanced was non-significant

(continued)

Table 1. Continued.

Author, year and location	Targeted substance/health behaviour	Study characteristics	Description of DI	n participants	Characteristics of study population	Engagement outcome measure(s)	Reported engagement results
McClure et al. 2013 USA	Tobacco	Design: randomised fractional factorial design with four two-level intervention factors Participants: adult smokers Factors: participants randomised to one of two levels of each four factors: (1) Message tone (prescriptive vs. motivational), (2) navigation autonomy (dictated vs. not), (3) proactive email reminders (yes vs. no), (4) personally tailored testimonials (yes vs. no) EPS(s): tailoring/reminders recruitment: identified from automated health plan records DI format: web-based	Name: Q2 Tailored (to stage of readiness to quit and demographics) internet based smoking cessation programme offering material addressing: the risks of smoking; benefits of quitting smoking; perceived barriers to treatment, perceived barriers to quitting; strategies for reducing smoking Factor 1: message tone (prescriptive/didactic vs. motivational) Factor 2: navigation autonomy (dictated order vs. not) Factor 3: proactive email reminders (yes vs. no) Factor 4: inclusion of personally tailored testimonials (yes vs. no)	N = 1865	% male: 37% % White: 82% Age: 44.2 years (SD = 14.7) Readiness to quit: in next 30 days: 44%	(1) Website visits, (2) content pages viewed, (3) minutes online, (4) content areas viewed	(1) Tailored testimonials: no effect (2) Email reminders increased visits***, page views***, min online*** not content area views (3) Prescriptive message tone increased page** and content*** area views but not visits or min online (4) Dictated navigation increased content area*** and page views**, min online***, not visits
Severson et al. 2008 USA	Smokeless tobacco	Design: RCT (two arms) Participants: adult smokeless tobacco users Arms: (1) enhanced version, (2) basic version EPS(s): interactivity/tailoring/social support/reminders/multimedia recruitment: online and print paid advertisements, radio DI format: web-based	Name: Chewfree.com Arm 1 (basic version): included best practices to cut down and cessation information Arm 2 (enhanced version): arm 1 plus guided, interactive and tailored 'planning to quit module'; 'staying to quit' module; multimedia content; two web-based support groups (peer and research staff; email reminders)	N = 2523 (Enhanced = 1260; basic = 1263)	% male: 98% % White: 98% Age: 37 years, SD = 9.6 Readiness to quit: reported a mean of 8 (SD 1.8) on an 11 point scale (ref.)	(1) Logins (2) Duration of visit (min)	(1) Participants in enhanced version logged in more compared with basic*** (2) Participants in enhanced version had higher duration of visit (min) compared with basic***

(continued)

Table 1. Continued.

Author, year and location	Targeted substance/health behaviour	Study characteristics	Description of DI	n participants	Characteristics of study population	Engagement outcome measure(s)	Reported engagement results
Strecher et al. 2008 USA	Tobacco	Design: Randomised fractional factorial design with five two-level intervention factors Participants: adult smokers Factors: participants randomised to one of two levels of each of the five factors: (1) outcome expectations, (2) efficacy expectations messages, (3) use of hypothetical success stories, (4) personalisation of the message source, (5) timing of message exposure EPS(s) tailoring/feedback strategies: Recruitment: identified from automated health plan records DI format: web-based	Name: not reported Core-intervention based on cognitive-behavioural methods of smoking cessation and relapse prevention including motivations for quitting, stimulus control and self-efficacy enhancement Factor 1: tailored outcome feedback expectations (high/low) Factor 2: tailored efficacy expectations messages (high/low) Factor 3: use of hypothetical success stories (high/low) Factor 4: personalisation of the message source (high/low) Factor 5: timing of message exposure (all at once/over a five week period with reminder)	N = 1866	% male: 40% % White: 79% Age: 46 years (SD = not reported) Motivation to quit (mean on 1–10 scale): = 8.3	Number of web-based smoking cessation sections opened by the participant	Regression analyses revealed that participants receiving high-depth tailored self-efficacy components*** and single exposure information opened significantly more web-sections***
Tensil et al. 2013 Germany	Alcohol	Design: RCT (two arms) Participants: adult drinkers (18 or over who drank at level considered harmful or consumed more than 24/12 g (male/female) of pure alcohol per day on average in the past week) Arms: (1) original version, (2) revised version EPS(s): tailoring (original vs. enhanced) Recruitment: email DI format: web-based	Name: Change Your Drinking Internet-based self-help programme with automated tailored feedback. Based on solution-focused brief intervention ⁵⁹ and cognitive behavioural therapy Arm 1: includes drinking self-assessment, tailored feedback, goal planning, drinking diary and goal feedback based on motivational interviewing techniques ⁶⁰ Arm 2: revised version of above including more intensive daily short tailored feedback on the individual's alcohol use and graphical display of alcohol use, detailed tailored feedback on alcohol use and tips on how to cope with risk situations on day 7, and advice for reflecting on reasons for reducing drinking and advice on rewarding achievement of goals	N = 595 (Original = 300, revised = 295)	% male: original = 59%, revised = 63% % White: not reported Age: original mean = 29.8 (10.3), revised = 29.0 (9.4) Measure of motivation to cut down: not reported	Diary usage (dichotomous: used at least once)	The usage of the diary was significantly higher*** in the revised version of the programme (diary used at least once, original version n = 167 (55.7%), revised n = 238 (80.7%))

(continued)

Table 1. Continued.

Author, year and location	Targeted substance/health behaviour	Study characteristics	Description of DI	n participants	Characteristics of study population	Engagement outcome measure(s)	Reported engagement results
Reminders							
Danaher et al. 2012 USA	See Danaher above						
Houston et al. 2015 USA	See Houston above						
McClure et al. 2013 USA	See McClure above						
Muñoz et al. 2009 USA	Tobacco Design: RCT (four arms) Participants: adult Spanish/English speaking internet users who smoked cigarettes Arms: (1) basic version, (2) basic + email reminders, (3) basic + email reminders + mood management lessons, (4) basic + email reminders + mood management lessons + social support EPS(s): reminders/social support/online therapist guidance Recruitment: online/word of mouth DI format: web-based	Name: or a Arm 1: static online smoking cessation guide covering: reasons to quit, cessation strategies, relapse prevention and management, pharmacological aids. Also included a cigarette counter and online journal Arm 2: Basic version + automated emails with links to sections of the GUIA keyed to quit date Arm 3: arms 1+2 and eight-lesson cognitive-behavioural mood management course Arm 4: arms 1+2+3+ online support group	N = 1000 (Arm 1 = 247; arm 2 = 251; arm 3 = 251; arm 4 = 251)	% male: 55% % White: 70% Age: 38 years, SD = 11.3 Confidence in quitting (as proxy for readiness to quit): mean = 6.8 on a 10 point scale (SD = 2.0)	As a proxy for engagement (1) cigarette counter usage, (2) online journal usage	Conditions with email reminders (2, 3 and 4) had increased cigarette counter*** and journal usage*** compared with condition 1 (data not reported). Conditions 3 and 4 had increased cigarette counter usage compared with condition 2***	
Streicher et al. 2008 USA	See Strecher above						
Severson et al. 2008 USA	See Severson above						
Delivery strategies							
Lieberman et al. 2006 USA	Alcohol Design: RCT (two arms) Participants: adult drinkers Arms: (1) text based feedback, (2) multimedia based feedback EPS(s): feedback strategies (text vs. multimedia) Recruitment: online DI format: web-based	Name: Alcohol Check up Web-based alcohol screening and personalised feedback Arm 1: feedback presented in HTML formatting Arm 2: multimedia feedback that included an animated photograph of a woman's face which personalised the programme as she guided the user through the feedback modules	N = 288 Between arms not reported	% male: text = 62.8%, multimedia = 69% % White: text = 87.2%, multimedia = 86.8% Age: text mean = 37.2 (11.8), multimedia = 36.0 (12.1) Measure of motivation to cut down: not reported	Feedback modules viewed	Participants in the personalised guide arm (multimedia) viewed significantly*** more viewed feedback modules (3.9) than participants in the text-only group (3.7)	

(continued)

Table 1. Continued.

Author, year and location	Targeted substance/health behaviour	Study characteristics	Description of DI	Characteristics of study population	Engagement outcome measure(s)	Reported engagement results
McClure et al. 2013 USA	See McClure above					
Schulz et al. 2013 Netherlands/Germany	Alcohol Participants: adult drinkers (18 or over who drank at level considered unhealthy Arms: (1) wait list control, (2) summative feedback, (3) alternating feedback EPS(s): delivery (feedback) strategies Recruitment: online DI format: web-based	Design: RCT (three arms) Participants: adult drinkers (18 or over who drank at level considered unhealthy Arms: (1) wait list control, (2) summative feedback, (3) alternating feedback EPS(s): delivery (feedback) strategies Recruitment: online DI format: web-based	Name: Alcohol-Everything Within the Limits? Three session, tailored programme targeting adult problem drinkers based on the I-Change model ⁵⁸ including personalised feedback and advice for cutting down drinking Arm 1: waiting list control Arm 2: summative feedback. All feedback provided at once after answering questions Arm 3: alternating feedback where participants received questions and personal advice alternately	N = 448 (Control = 135, summative feedback = 181, alternating feedback = 132) % male: 57% % White: not reported Age: mean = 41.7 years old (SD = 15.7) Measure of motivation to cut down: not reported	(1) Programme completion	After the first session, drop out was significantly less in the alternating condition**. However, no differences in programme completion at three and six months
Stanczyk et al. 2013 Netherlands	Tobacco Participants: smokers, 16 years or older, motivated to quit, categorised as lower or higher educated participants Arms: (1) text based messages, (2) multimedia based messages EPS(s): delivery strategies (text vs. video) Recruitment: online DI format: web-based	Design: RCT (two arms) Participants: smokers, 16 years or older, motivated to quit, categorised as lower or higher educated participants Arms: (1) text based messages, (2) multimedia based messages EPS(s): delivery strategies (text vs. video) Recruitment: online DI format: web-based	Name: Steunblij Stoppen Web-based tailored smoking cessation programme based on the I-Change Model ⁵⁸ Arm 1: (text) tailored messaging delivered by text Arm 2: (video) tailored messaging delivered by video. All content between the two arms was identical	N = 204 (Text: n = 104, video: n = 100) % male: text = 40.5, video = 22 % White: not reported Age: text mean = 46.6 (11.9), multimedia = 48.2 (12.0) Readiness to quit within one month (%): text = 19.2, video = 18.5	Duration of visit (min)	Text condition spent a mean of 7.2 min online, video condition spent a mean of 7.8 min online***
Streicher et al. 2008 USA	See Strecher above					
McClure et al. 2013 USA	See McClure above					
Social support						
Danaher et al. 2012 USA	See Danaher above					

(continued)

Table 1. Continued.

Author, year and location	Targeted substance/health behaviour	Study characteristics	Description of DI	Characteristics of study population	Engagement outcome measure(s)	Reported engagement results
Ramo et al. 2015 USA	Tobacco	Design: RCT (two levels of randomisation: three arms randomised to Facebook group by stage of change, within each arm randomised to one of three incentive conditions) Participants: young adult smokers (18–25 years old) Facebook group arms: (1) not ready to quit, (2) thinking about quitting, (3) ready to quit Incentive arms: (1) no incentive, (2) personal condition: \$50 gift card, (3) altruistic incentive: \$50 gift card to charity EPS(s): incentives Recruitment: Facebook advertisement DI format: web-based	Name: Tobacco Status Project Description: a Facebook intervention tailored to stage of change; Trans-theoretical Model strategies ⁶⁰ for cessation; group messaging (posts); CBT sessions; ask the doctor sessions Arm 1: no incentive Arm 2 (personal incentive): \$50 gift card incentive emailed to participant for commenting on all 90 posts posted to their Facebook group Arm 3 (altruistic incentive): \$50 gift card incentive donated to charity for commenting on all 90 posts posted to their Facebook group	% male: 80% % White: 80% Age: 21 years, SD = 2.1 Smoking goal: 10% reported abstinence goal, 60% reduction goal, 30% no goal n participants N = 79 Arm 1 (not ready to quit): n = 35 randomised to personal condition: 8, altruistic condition: 11, no incentive: 16 Arm 2 (thinking about quitting); n = 32 Personal condition: 9, altruistic condition: 8, no incentive: 15 Arm 3 (ready to quit): n = 12 Personal condition: 3, altruistic condition: 3, no incentive: 6	As a proxy for engagement number of 'likes' and comments measured	(1) For enhanced sample: no significant difference among incentive conditions on number of comments made to Facebook groups (2) For those who commented at least once: personal incentive condition made more comments than other two conditions*
Stoops et al. 2009 USA	Tobacco	Design: RCT (two arms) Participants: adult tobacco smokers Arms: (1) Abstinence Contingent group (AC), (2) Yoked Control group (YC), EPS(s): Incentives Recruitment: local advertisements, word of mouth DI format: web-based	Name: Not given Intervention was an internet-based CO breath recorder which provided feedback and progress tracking Arm 1 (AC): participants received monetary incentives contingent on recent smoking abstinence (CO level of ≤4 parts per million) Arm 2 (YC): participants received monetary incentives independent of smoking status. Participants were matched to a participant in the AC group and were reinforced on a schedule identical to that of their 'yoked' partner	% male: AC group: 26%; YC group: 24% % White: AC group: 94%; YC group: 94% Age: AC group: mean = 38 years, range = 21–58; YC group: mean = 40, range = 18–61 Readiness to quit: not reported N = 68 (AC = 35; YC = 33)	As a proxy for engagement number of videos posted to website was used as a measure of engagement	No significant effect of group on odds of posting videos to the website. AC group posted 68% of the total number of videos expected, YC group posted 67% of the total number of videos expected

Note: * = <.05, ** = <.01, *** = <.001. RCT: randomized controlled trial; EPS: engagement promoting strategy; DI: Digital Intervention; CBT: cognitive behavioural therapy; MI: Motivational Interviewing

Table 2. Prevalence of engagement promoting strategies.

EPS category	N studies included (%)	EPS sub-category	n (%)
Tailoring	7 (47)		
		Readiness to quit	4 (40)
		Self-efficacy + barriers to quitting	3 (30)
		Abstinence status	2 (20)
		Testimonials/success stories	2 (20)
		Goals/motivation to quit	2 (20)
		Interests	1 (10)
		Mood/negative affect	1 (10)
		Health + lifestyle factors	1 (10)
		Preparatory planning	1 (10)
		Personalised message source	1 (10)
Reminders	6 (40)		
		Email	6 (40)
Delivery strategies	5 (33)		
		Multimedia (text/image/video)	3 (60)
		Single/staged/alternating feedback	2 (40)
		Message tone (prescriptive/motivational)	1 (20)
		Navigation autonomy (dictated vs. not)	1 (20)
Social support	6 (40)		
		Peer support	5 (83)
		Therapist support	3 (50)
Incentives	2 (13)		
		Contingent on abstinence	1 (50)
		Contingent on website comments	1 (50)

Research question 2: to determine whether the identified EPSs increased engagement with online SUIs

Effectiveness of EPSs

Table 3 provides an overview of the effectiveness of the different strategies. There was a lack of standardised reporting of engagement outcomes which made pooling of effects an unsuitable technique.

Where independent effects could be reported, these are described under the relevant EPS category below. Unfortunately, due to two studies^{44,63} using a multi-component design comparing a basic with an enhanced version of a SUI, which included multiple EPSs together, examination of the effects of the individual strategies reported in these studies in Table 2 was not possible. The synthesis below includes only studies where individual effects of EPS categories could be reported.

Table 3. Effectiveness of engagement promoting strategies in experimental arms.

Tailoring Author	EPS sub-type(s)	Significant
Tensil et al. 2013	1. Abstinence status 2. Goals/motivation to quit	Yes
Strecher et al. 2008	1. Goals/motivation to quit (depth outcome expectations condition) 2. Health + lifestyle (depth outcome expectations condition) 3. Self-efficacy + barriers to quitting ^a 4. Testimonials/success stories 5. Personalised source ^a	Yes
McClure et al. 2013	1. Testimonials/success stories 2. Self-efficacy + barriers to quitting	No
Elfeddali et al. 2012 ^b	1. Self-efficacy + barriers to quitting 2. Mood/negative affect 3. Level of planning	Unclear
Danaher et al. 2013 ^c	1. Readiness to quit 2. Interests	Unclear
Houston et al. 2015 ^c	1. Readiness to quit	Unclear
Severson et al. 2008 ^c	1. Abstinence status 2. Readiness to quit	Unclear
Reminders		
McClure et al. 2013	Email	Yes
Muñoz et al. 2009	Email	Yes
Strecher et al. 2008	Email	Yes
Danaher et al. 2013 ^c	Email	Unclear
Houston et al. 2015 ^c	Email	Unclear
Severson et al. 2008 ^c	Email	Unclear
Delivery strategies		
Lieberman et al. 2006	Multimedia (images ^a vs. text)	Yes
Stanczyk et al. 2013	Multimedia (video ^a vs. text)	Yes
Strecher et al. 2008	Single exposure ^a vs. staged	Yes

(continued)

Table 3. Continued.

Tailoring Author	EPS sub-type(s)	Significant
Schulz et al. 2013 ^d	Single exposure vs. alternating feedback	Unclear
McClure et al. 2013 ^d	1. Dictated order of content vs. not 2. Message tone (prescriptive/motivational)	Unclear
Social support		
Houston et al. 2015	Peer + therapist support	No
Danaher et al. 2013 ^c	Peer support	Unclear
Muñoz et al. 2009	Peer support	Unclear
Schaub et al. 2015 ^d	Therapist support	Unclear
Severson et al. 2008 ^c	Peer + therapist support	Unclear
Stoddard et al. 2008 ^d	Peer support	Unclear
Incentives		
Ramo et al. 2015 ^d	Contingent on website comments	Unclear
Stoops et al. 2009	Contingent on abstinence	No

^aIndependent effects reported for specific sub-types.^bStatistical tests not conducted.^cThese studies used a multi-component design so eliciting individual effects was not possible.^dInconsistent findings reported between engagement measures.

Tailoring

Seven studies reported tailoring as an EPS; four of these reported data for individual effects of tailoring as an EPS.^{2,29,35,45} Two reported significant increases in engagement compared with the control conditions,^{2,29} one reported increases in engagement measures compared with control but did not report statistical tests.⁴⁵ There was high variability between the types of tailoring features used, with 10 different sub-types included overall (see Table 2). Four out of seven studies tailored to more than one factor at time, subsequently, unless specified, results of individual effects of sub-types must be considered with caution.^{2,44,45,63}

Tensil et al.² compared two self-help programmes which provided automated feedback tailored to abstinence status (drinking level). Both original and revised versions included tailored feedback on participants'

alcohol-use at baseline, general information on control strategies and a drinking diary. In the revised version participants received additional daily tailored feedback messages on abstinence status and goals/motivations to quit. The authors reported increased diary usage in the revised arm compared with those in the original arm. Individual effects of the tailoring sub-types were not reported.

Strecher et al.²⁹ used a factorial design allowing for comparison of sub-types of tailoring EPSs, randomising participants to one of two levels of five different treatment components, four of which compared a high level of tailoring with a low level of tailoring. The tailoring components were: (1) depth of outcome expectations (such as health and lifestyle factors and motivations for quitting); (2) depth of efficacy expectations (such as barriers to quitting); (3) depth of success stories where participants received a hypothetical story tailored to the participant; (4) 'personalised source' where messages were presented as either friendly or informal. Analyses revealed that participants receiving high-depth tailored self-efficacy components and high-depth personalised source components opened significantly more web-sections.

Elfeddali et al.⁴⁵ compared two versions (low and high) of a tailored web-based programme targeting smoking cessation in adults whereby the level of tailoring was manipulated. A questionnaire only control condition was also included. In the low tailored group respondents received tailored feedback prior to their quit attempt. Feedback was tailored to: self-efficacy, readiness and motivations to quit and preparatory planning. In the enhanced group (AP+) participants received additional tailored feedback post-quit attempt to self, efficacy, recovery self-efficacy, mood/negative affect and level of planning. Providing additional tailored feedback after the quit attempt resulted in more planning assignments being completed, suggesting that a higher dose of tailoring, consistent with Strecher et al.²⁹ above, is more effective. However statistical tests were not completed.

McClure et al.³⁵ used a factorial design, randomising participants to one of two levels of the four experimental factors. One of these factors was tailored; participants received three highly tailored testimonials (or not) targeting self-efficacy. Tailored testimonials did not significantly affect engagement.

Reminders

Six studies reported reminders as an EPS.^{29,35,43,45,46,62} Four studies examined the individual effects of use of reminders, which are reported here. McClure et al.,³⁵ using a high quality factorial design with a large sample size ($n = 1865$), randomised participants to one of two

levels of four experimental factors. The authors examined whether reminders increased website visits, content pages and areas viewed and minutes spent online for an internet-based smoking cessation programme. Reminders significantly increased visits, views and minutes spent online compared with participants who did not receive reminders.

Muñoz et al.⁴⁷ examined a basic version of an online smoking cessation guide compared with three other versions of increasing complexity. The basic version included an instruction guide, cigarette counter and online journal; the second version included the basic version plus reminders; the third included version two plus mood management lessons and the fourth version included version three plus social support. Conditions with reminders had significantly higher tool usage (cigarette counter and diary usage) than condition one.

Strecher et al.²⁹ also used reminders in one factor of their five factor RCT. The authors reported that this EPS increased engagement; however, this result needs to be considered with caution as the use of reminders was combined with a manipulation of delivery strategy (timing of message source) so the extent to which reminders were independently associated with engagement is not known.

Finally Houston et al.⁴⁶ delivered a quit smoking website, with three arms of increasing complexity: basic, enhanced and enhanced plus. The enhanced group included the intervention from the basic arm as well as the addition of email reminders; the enhanced plus included the same content as well as a social support feature. Participants in the enhanced plus logged in more times than in the basic version, but not compared with the enhanced version, suggesting an effect of reminders upon engagement.

Delivery strategies

Five studies reported delivery strategies as an EPS. For the purpose of this review, 'delivery strategies' refers to EPSs which manipulate the type of delivery method of the intervention, for example, text versus video. Three out of five of the studies reported increases in engagement.^{29,49,52} Multimedia delivery of information may be an important delivery strategy to promote engagement, as may be providing information all at once, as opposed to in stages. However, with the exception of Strecher et al.²⁹ the studies in this EPS category were of low quality (see Appendix 3).

Two of the studies compared text-based with multimedia-based delivery formats (Lieberman⁵² and Stanczyk et al.⁴⁹). Lieberman⁵² examined the number of feedback modules completed in a study examining text versus multimedia messages for an adult web-based alcohol screening and personalised feedback

programme. In one arm participants received all feedback in HTML text, whilst in the other, feedback was delivered in multimedia format. The multimedia arm consisted of a 'personified guide', where results were presented by a photograph of a woman who would guide participants through the feedback process. Participants in the multimedia arm viewed significantly more feedback modules than participants in the text-only group.

Stanczyk et al.⁴⁹ conducted a RCT examining the delivery strategy of feedback messages in a single session web-based smoking cessation programme. Participants received feedback either by text or by video. Participants in the video condition spent significantly more time online (minutes) than those in the text condition.

The three other studies manipulated the delivery schedule of feedback.^{29,35,54} Strecher et al.²⁹ used a factorial design randomising participants to one of two levels of five different treatment components, one of which was the delivery strategy of the intervention materials. Participants received the materials either all at once (single exposure) or over the course of five weeks. Participants in the single exposure arm opened significantly more web-sections than participants who received the content over the course of five weeks. However, this result needs to be considered with a caveat: reminders were also included in this factor and it is not possible to disentangle which EPSs were having an influencing effect.

Schulz et al.⁵⁴ conducted an RCT of a web-based, three session tailored intervention targeting problem adult drinkers. Participants were randomised to a control group or the experimental group, which was divided into two sub-groups: alternating and single exposure. In the alternating group participants received alternating questions (three in total) followed by personal advice. In the single exposure group, participants answered all the questions and were given the personal advice all at once. The feedback messages were the same for both groups. In the alternating condition, significantly fewer participants dropped out after the first session; however, there were no differences in programme completion at three and six months.

McClure et al.,³⁵ using a high quality factorial design, manipulated navigational autonomy where participants could view content in any order or a pre-specified order. The authors also manipulated message tone, which was either prescriptive (in a didactic tone) or motivational. Dictated navigation significantly improved engagement on three out of four measures (content areas viewed, content pages viewed, cumulative duration), whereas prescriptive message tone significantly increased engagement on two out of four measures (content areas viewed, content pages viewed).

Social support

Six studies reported social support as an EPS.^{44,46,47,55,57-63} Four studies examined individual effects of social support features.^{46,56} Results for the use of social support as an EPS were inconclusive. Two studies reported peer social support as an EPS.^{54,46,56} Stoddard et al.,⁵⁶ in a very high quality study with large sample size ($n=1375$), examined the effect of a social bulletin board where users could interact with each other within a self-help smoking cessation programme. Stoddard reported two engagement outcomes: visit duration (minutes) and number of visits to pages; the authors reported an increase in visit duration but not in visits to pages.

Muñoz et al.⁴⁷ also used social support as a strategy, including in one of the four arms of the trial was an online social support group. The arm with a social support feature did have higher engagement (cigarette counter usage) compared with the arm with only email reminders as an EPS. However, included in the arm with social support as an EPS was a cognitive behavioural mood management course, which may also have had a moderating effect upon engagement, therefore these results need to be considered with caution.

Another study reported online therapist guidance as an EPS. Schaub et al.,⁵⁴ in a good quality study, examined whether the addition of chat counselling to a web-based tailored self-help programme for adult cannabis users increased feature usage (module and diary completion). The results for the two engagement outcomes were conflicting: participants in the chat-counselling arm did not complete significantly more modules than those in the original arm. However, participants in the chat-counselling arm did complete significantly more diary entries than those in the original arm.

Houston et al.⁴⁶ included a peer and therapist led social support group in the enhanced plus condition of a three arm trial (basic vs. enhanced vs. enhanced plus) of a quit smoking website and reported that the addition of a peer and therapist led social support group did not increase engagement over the enhanced arm, which included the basic component plus reminders. As reminders in the enhanced arm increased engagement compared with controls, this suggests an effect of reminders not of social support.

Incentives

Two studies examined the effectiveness of incentives to improve engagement.^{48,52} These results were conflicting, with one study reporting increases in engagement only for participants who had already engaged. These studies were of low quality and reported small sample sizes (see Table 2).

Ramo et al.⁴⁸ conducted a smoking cessation intervention on Facebook and examined the effect of different incentives on the number of likes and comments posted on the Facebook pages (as a proxy for engagement). There were three incentive conditions: (1) no incentive; (2) altruistic incentives where a \$50 gift card was given to charity if participants commented on all 90 posts posted on the group's Facebook page; (3) personal incentives where participants were given a \$50 gift card for commenting on all posts. Analyses reported that there were no differences in engagement between incentive arms. However, for those who had already engaged and commented at least once, personal incentive condition made significantly more comments than the other two conditions.

Stoops et al.⁵² also examined whether incentives improved engagement with an online smoking cessation website; however, incentives were not specifically targeting engagement: participants either received incentives contingent on recent smoking abstinence, or received incentives independent of smoking status. The authors measured the number of online videos participants posted to the website, which captured participants giving CO readings. There was no significant effect of group on odds of posting videos to the website.

Multi-component designs comparing a basic version with an EPS enhanced version

Two studies compared a basic with an enhanced version of the intervention.^{44,63} The enhanced versions incorporated a range of EPSs including: interactivity; tailoring; multimedia; reminders; online therapist guidance and social support. Both studies reported significant effects between basic and enhanced versions. Due to the study designs it was not possible to examine the effect of the individual strategies, therefore the EPSs in this category were considered together. Danaher et al.⁴⁴ compared a basic and enhanced version of a web-based smokeless tobacco programme targeting 14–25 year olds. The enhanced version included four additional strategies compared with the basic self-help programme (interactivity/tailoring/multimedia/reminders). The authors reported significant increases in logins and duration of visits for the enhanced version. In a similar study, Severson et al.⁶³ examined the use of six different strategies (interactivity/tailoring/social support/reminders/multimedia) in an enhanced versus basic study for smokeless tobacco users. Both logins and visit duration (minutes) were higher in the enhanced version.

Discussion

This systematic review of the effectiveness of engagement promoting strategies in online SUIs has focused

on evidence from RCTs, with the majority of studies targeting tobacco cessation. This review is the first to explore which EPSs are effective specifically in substance-use populations and the first to use Cochrane Methodology, with previous reviews examining more generally the health behaviour change literature.

Five different EPSs were identified: tailoring; reminders; delivery strategies; social support; incentives. Overall, the strongest evidence points to the use of tailoring, email reminders and multimedia and single exposure delivery strategies to improve engagement. Social support features demonstrate inconclusive results and may be effective in promoting some engagement behaviours but not others. Incentives did not demonstrate effectiveness.

Tailoring demonstrated promising results, which is consistent with previous research.^{18,22,66} Tailoring has been hypothesised to be effective because it adapts the intervention to specific attributes of the user.¹⁹ In the alcohol field tailoring has been demonstrated to improve the intervention by making it more meaningful and relatable to the user,⁶⁷ increasing their likelihood of engaging with the intervention content. Tailoring to self-efficacy demonstrated tentatively positive results, with a higher dosage of tailoring improving engagement.^{29,45} Tailoring to goals/motivations to quitting was inconclusive as one study provided positive results,² but in combination with another tailoring sub-type (abstinence status). The other study²⁹ also included this sub-type in combination with another (health and lifestyle factors) but did not report an effect of higher dosage on engagement. Tailoring to a friendly, personalised source was effective, but was only examined in a single study,²⁹ as was tailoring to abstinence status.²

There was high variability between the types and combinations of tailoring features used in this review (see Table 2) and disentangling which subtypes of tailoring EPSs are the most effective remains a challenge. Therefore these results need to be considered with caution: Only two studies^{29,35} examined effects of individual sub-components in isolation. Elfeddali et al.⁴⁵ and Tensil² examined tailoring sub-types of self-efficacy, abstinence status, goals and motivations, mood and level of planning in combination, therefore work is needed to experimentally study the influence of individual tailoring strategies to confirm these findings. Furthermore, the descriptions of how tailoring was utilised in the studies was often limited, and specific details on how tailoring was modified in the studies, for example how interventions were tailored to 'self-efficacy', were lacking, making it difficult for the design of future interventions to replicate the effective components.

In a recent meta-analysis, the use of reminders to improve engagement with digital health interventions showed small but significant results.²³ This supports the findings from the current review. Given the wealth of research from the broader healthcare field for the effectiveness of reminders to improve engagement with healthcare services,^{68,69} there is again a need for further high quality research to examine the role of reminders in promoting engagement with online substance-use interventions.

The methods through which content is delivered demonstrate potential to increase engagement. Perski et al.¹⁸ highlight that when users have control over how they view content, and have free choice over when they interact with it, evidence points towards higher engagement rates. Equally, this review found that multimedia content had higher engagement rates than simple text based content. This maps onto the concept of usability and aesthetics, which have been demonstrated to be key factors in engagement.^{70,71} For example content which is well presented and easy to digest is more likely to be utilised than content which is difficult to navigate or read, which is supported by the findings in this review.

Social support, whether by peer or therapist, was an effective strategy to increase engagement in a previous review by Brouwer et al.⁹ However, in the current review, social support features were highly inconclusive and had differential effects on engagement depending on which engagement measure was used. This points to specific EPSs targeting different engagement behaviours. User engagement research has reported that social or 'community' features are highly desired in apps for cutting down alcohol-use.⁶⁷ With numerous studies pointing to social support being a useful tool to promote behaviour change,⁷²⁻⁷⁴ further research is warranted.

It was surprising that the two studies examining the effect of incentives on engagement did not report positive results. In the substance-use field, providing voucher or prized-based reinforcement for abstinence from drugs as well as treatment adherence is effective.^{68,75} One potential moderator may be the influence of motivation level on engagement; in the present review, Ramo et al.⁴⁸ saw an increase in engagement compared with controls only for participants who had already engaged by posting previously. If incentives work to promote engagement in users, it may only be in those motivated to reduce their use already; however, this hypothesis needs further testing.

The evidence also suggests that using a multi-component EPS approach is more effective than a single EPS. Both studies using this approach demonstrated an effect of multi-component strategies compared with no strategies. Furthermore, in the Muñoz et al.⁴⁷ study, reminders were significantly more effective

compared with a basic version, but the addition of social support was more effective than reminders alone, pointing towards an additive effect of using multiple strategies.

Various theoretical models have been proposed to explain the mechanisms behind user engagement with online health interventions. Persuasive system design has been influential in the engagement field, proposing that technology has the capacity to influence, reinforce, change and shape attitudes or behaviours through supporting users to achieve the main objective of the technology.⁷⁶ Techniques such as monitoring, feedback, tailoring, aesthetic design, novelty, credible source and social support have been proposed to influence usage with online interventions.^{77,78} This model has been further extended by Short et al.²⁰ to consider individual, environmental and technology level factors that may moderate engagement. In a recent conceptual review, Perski et al.¹⁸ hypothesise that engagement is directly influenced by the context, content and delivery of the intervention as well as characteristics of the population; however, further investigation into these associations is warranted as evidence is tentative. Engagement models have been concisely described in a scoping review by Ryan et al.⁷⁹

All the studies were computer web-based, despite the increasing availability of app-based SUIs downloadable from the app stores.⁶⁷ Coupled with a lack of evidence-based design for these programs,⁸⁰ there is a need for rigorous evaluation of engagement in app-based interventions. How people use computer-based interventions may be different from how they use app-based interventions. For example, whether a smartphone or computer is being used may influence the amount of time a user engages with an intervention. Users may also respond differentially to engagement features depending on delivery mode. This review was unable to explore any of these issues; further exploratory work would be beneficial.

An inductive approach following Cochrane methodology was used here to identify EPSs. However, due to this rigorous approach there may remain EPSs identified in other frameworks that were not identified in this review, such as novelty, aesthetics and system credibility.^{18,19,81} Unfortunately, to date, these EPSs have not been evaluated using a RCT design within the SUI field, therefore whilst they are potentially promising techniques, further research is warranted.

One issue which has yet to be defined in the engagement literature is how best to measure it, both quantitatively and qualitatively. As this review highlights, many different measures are used, with no consensus between researchers, and conflicting results, whereby an EPS will have antipodal effects on different engagement measures. Perksi et al.¹⁸ suggest that multiple

dimensions of engagement should be utilised, for example self-report, web metrics and physiological or psychophysical correlates of e-health interaction. However, as Yardley et al.⁸² highlight, we still do not fully understand what constitutes ‘effective engagement’ and how to measure it. Engagement is a complex and dynamic behaviour, and it has been argued that simply ‘increased engagement’ measured via web metric data as is typical with existing RCTs might not equate to ‘effective engagement’. For example, non- or ceased engagement could signify that the intervention has been effective whilst sustained engagement might suggest that the user is not learning to self-regulate independently and is over-relying on the intervention. Therefore defining the most appropriate measures of engagement in the context of each individual intervention is important as a ‘one size fits all’ approach may disregard nuanced differences between engagement behaviours.

Limitations

The major strengths of this review are its rigorous methodology using Cochrane Guidance. However, fewer studies were identified than expected and meta-analytic techniques could not be applied as had been planned. The size of effect of the different strategies could therefore not be measured across studies, so reporting of results here is tentative. Further high-quality trials examining the effect of EPSs need to be conducted, reporting consistent engagement outcomes, in order for measures of effect to be established.

There was high heterogeneity of engagement outcomes reported in the studies, an issue which has been highlighted in previous reviews.^{9,22,23,83} Until a consistent measure of engagement is used by researchers, reviews comparing the results of different outcomes need to be considered cautiously. In the current review, four studies reported conflicting engagement results depending on the engagement measure used, demonstrating the inconsistencies different measures of engagement can produce. Future work needs to establish whether there is an optimal ‘engagement outcome’, or combination of outcomes, for future evaluations to consider. Brouwer et al.⁹ reported that different engagement features influenced different engagement behaviours, for example, peer support was associated with a longer website stay and email contact was related to more logins. This suggests that there may not be one outcome appropriate for measurement of all types of engagement strategies.

This review considered only high quality RCTs; however, it is recognised that as engagement is a complex and multi-faceted issue,⁸⁴ studies employing other designs also provide important data to the field.

Future work may benefit from consideration of qualitative work alongside RCTs to elucidate further the inconsistencies in findings.

Piloting of search strategy and review of relevance of techniques for narrative synthesis were conducted by only one reviewer (JM), potentially introducing bias.

It is noted here that only two of the RCTs in this review cited engagement as their primary outcome. Therefore the majority of studies may not have been powered to detect effects.

The majority of the studies examined adult smoking populations so the limitations in the generalisability of these findings are acknowledged. However, as the findings are in line with evidence from other healthcare areas, the results of this review are potentially applicable across the substance-use domain. It is also possible that the differences in effect of EPSs compared with controls were due to the EPS arms taking longer to deliver than controls and therefore required more user time and engagement. This is most applicable to the studies which examined an enhanced compared with a basic version.

Conclusion

This review has identified five strategies used to promote engagement in online substance-use interventions and has highlighted that interventions that use tailoring, reminders and multimedia and single-exposure delivery are potentially effective. The evidence for social support is highly inconclusive. Incentives need to be studied more thoroughly. Promoting engagement is crucial to optimally deliver the harm reduction interventions in online programmes; however, more consistent reporting of engagement outcomes and further research are still required.

Contributorship: JM, PD and CD conceived the study and contributed to protocol development. JM wrote the first draft of the manuscript. SF was the second screener. All authors reviewed and edited the manuscript and approved the final version of the manuscript. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Declaration of Conflicting Interests: The authors declare that there is no conflict of interest.

Ethical approval: Ethics approval was not required for this project.

Funding: This work was supported by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research Programme (grant reference number RP-PG-0609-10162) and Collaboration for Leadership in Applied Health Research and Care South London at King’s College Hospital NHS Foundation Trust, the NIHR Biomedical Research Centre for Mental Health at South London and Maudsley NHS Foundation Trust and King’s College London, and the NIHR Collaborations for Leadership in Applied Health Research and Care South London at King’s College Hospital NHS Foundation Trust.

Guarantor: JM.

Peer review: This manuscript was reviewed by Ghadah Alkhalidi, King Saud University, Saudi Arabia; Olga Perski, University College London, UK and Siobhan O'Connor, Edinburgh Napier University, UK.

References

1. DanaHER BG, Boles SM, Akers L, et al. Defining participant exposure measures in web-based health behavior change programs. *J Med Internet Res* 2006; 8: 13P.
2. Tensil MD, Jonas B and Struber E. Two fully automated web-based interventions for risky alcohol use: Randomized controlled trial. *J Med Internet Res* 2013; 15: e110.
3. Rooke S, Copeland J, Norberg M, et al. Effectiveness of a self-guided web-based cannabis treatment program: Randomized controlled trial. *J Med Internet Res* 2013; 15(2): e26.
4. Donoghue K, Patton R, Phillips T, et al. The effectiveness of electronic screening and brief intervention for reducing levels of alcohol consumption: A systematic review and meta-analysis. *J Med Internet Res* 2014; 16(6): e142.
5. Tait RJ, Spijkerman R and Riper H. Internet and computer based interventions for cannabis use: A meta-analysis. *Drug Alcohol Depend* 2013; 133: 295–304.
6. Rooke S, Thorsteinsson E, Karpin A, et al. Computer-delivered interventions for alcohol and tobacco use: A meta-analysis. *Addiction* 2010; 105: 1381–1390.
7. Kaner EF, Beyer FR, Garnett C, et al. Personalised digital interventions for reducing hazardous and harmful alcohol consumption in community-dwelling populations. *The Cochrane Library* 2015; 1: CD011479.
8. Kohl LF, Crutzen R and de Vries NK. Online prevention aimed at lifestyle behaviors: A systematic review of reviews. *J Med Internet Res* 2013; 15(7): e126.
9. Brouwer W, Kroeze W, Crutzen R, et al. Which intervention characteristics are related to more exposure to internet-delivered healthy lifestyle promotion interventions? A systematic review. *J Med Internet Res* 2011; 13(1): e2.
10. Crutzen R, de Nooijer J, Brouwer W, et al. Strategies to facilitate exposure to internet-delivered health behavior change interventions aimed at adolescents or young adults: A systematic review. *Health Educ Behav* 2010; 38: 49–62.
11. Brouwer W, Oenema A, Raat H, et al. Characteristics of visitors and revisitors to an Internet-delivered computer-tailored lifestyle intervention implemented for use by the general public. *Health Educ Res* 2009; 25: 585–595.
12. Verheijden MW, Jans MP, Hildebrandt VH, et al. Rates and determinants of repeated participation in a web-based behavior change program for healthy body weight and healthy lifestyle. *J Med Internet Res* 2007; 9: e1.
13. Davies CA, Spence JC, Vandelanotte C, et al. Meta-analysis of internet-delivered interventions to increase physical activity levels. *Int J Behav Nutr Phys Act* 2012; 9: 52.
14. Hutton HE, Wilson LM, Apelberg BJ, et al. A systematic review of randomized controlled trials: Web-based interventions for smoking cessation among adolescents, college students, and adults. *Nicotine Tob Res* 2011; 13: 227–238.
15. Vandelanotte C, Spathonis KM, Eakin EG, et al. Website-delivered physical activity interventions a review of the literature. *Am J Prev Med* 2007; 33: 54–64.
16. Funk KL, Stevens VJ, Appel LJ, et al. Associations of internet website use with weight change in a long-term weight loss maintenance program. *J Med Internet Res* 2010; 12: e29.
17. Couper MP, Alexander GL, Zhang N, et al. Engagement and retention: measuring breadth and depth of participant use of an online intervention. *J Med Internet Res* 2010; 12: e52.
18. Perski O, Blandford A, West R, et al. Conceptualising engagement with digital behaviour change interventions: A systematic review using principles from critical interpretive synthesis. *Transl Behav Med* 2017; 7: 254–267.
19. Ritterband LM, Thorndike FP, Cox DJ, et al. A behavior change model for internet interventions. *Ann Behav Med* 2009; 38: 18.
20. Short CE, Rebar AL, Plotnikoff RC, et al. Designing engaging online behaviour change interventions: A proposed model of user engagement. *Eur Health Psychol* 2015; 17: 32–38.
21. Evers KE. eHealth promotion: The use of the Internet for health promotion. *Am J Health Promot* 2006; 20: 1–14.
22. Schubart JR, Stuckey HL, Ganeshamoorthy A, et al. Chronic health conditions and internet behavioral interventions: A review of factors to enhance user engagement. *Comput Inform Nurs* 2011; 29: 81–92.
23. Alkhalidi G, Hamilton FL, Lau R, et al. The effectiveness of prompts to promote engagement with digital interventions: A systematic review. *J Med Internet Res* 2016; 18: e6.
24. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *Ann Intern Med* 2009; 151: W-65–W-94.
25. Barak A, Klein B and Proudfoot JG. Defining internet-supported therapeutic interventions. *Ann Behav Med* 2009; 38: 4–17.
26. Bailey JV, Murray E, Rait G, et al. Interactive computer-based interventions for sexual health promotion. *Cochrane Database Syst Rev* 2010; 10: CD006483.
27. Lieberman DZ. Effects of a personified guide on adherence to an online program for alcohol abusers. *Cyberpsychol Behav* 2006; 9: 603–607.
28. Allam A, Kostova Z, Nakamoto K, et al. The effect of social support features and gamification on a Web-based intervention for rheumatoid arthritis patients: Randomized controlled trial. *J Med Internet Res* 2015; 17: e14.
29. Strecher V, McClure J, Alexander G, et al. The role of engagement in a tailored web-based smoking cessation program: Randomized controlled trial. *J Med Internet Res* 2008; 10: e36.
30. Deterding S, Dixon D, Khaled R, et al. From game design elements to gamefulness: Defining gamification.

- In: *Proceedings of the 15th international academic MindTrek conference: Envisioning future media environments*. 22-24th September 2015. Tampere, Finland: ACM, New York 2011, pp.9–15.
31. Hilbert M and López P. The world's technological capacity to store, communicate, and compute information. *Science* 2011; 332: 60–65.
 32. Alkhaldi G, Hamilton FL, Lau R, et al. The effectiveness of technology-based strategies to promote engagement with digital interventions: A systematic review protocol. *JMIR Res Protoc* 2015; 4: e47.
 33. Cugelman B. Gamification: What it is and why it matters to digital health behavior change developers. *JMIR Serious Games* 2013; 1: e3.
 34. Lieberman DZ. Effects of a personified guide on adherence to an online program for alcohol abusers. *Cyberpsychol Behav* 2006; 9: 603–607.
 35. McClure JB, Shortreed SM, Bogart A, et al. The effect of program design on engagement with an internet-based smoking intervention: randomized factorial trial. *J Med Internet Res* 2013; 15: e69.
 36. Glasgow RE, Christiansen SM, Kurz D, et al. Engagement in a diabetes self-management website: usage patterns and generalizability of program use. *J Med Internet Res* 2011; 13: e9.
 37. Glasgow RE, Nelson CC, Kearney KA, et al. Reach, engagement, and retention in an Internet-based weight loss program in a multi-site randomized controlled trial. *J Med Internet Res* 2007; 9: e11.
 38. Ferney SL, Marshall AL, Eakin EG, et al. Randomized trial of a neighborhood environment-focused physical activity website intervention. *Prev Med* 2009; 48: 144–150.
 39. Higgins JP and Green S. *Cochrane handbook for systematic reviews of interventions*. Chichester, UK: Wiley-Blackwell, 2008.
 40. Popay J, Roberts H, Sowden A, et al. *Guidance on the conduct of narrative synthesis in systematic reviews A product from the ESRC methods programme Version* 2006; 1: b92, ESRC.
 41. Leamy M, Bird V, Le Boutillier C, et al. Conceptual framework for personal recovery in mental health: Systematic review and narrative synthesis. *Br J Psychiatry* 2011; 199: 445–452.
 42. Ryan R. Cochrane consumers and communication review group. 'Cochrane Consumers and Communication Review Group: data synthesis and analysis'. <http://ccrg.cochrane.org> (2013, accessed 18 January 2016).
 43. Danaher BG, Severson HH, Andrews JA, et al. Randomized controlled trial of MyLastDip: A web-based smokeless tobacco cessation program for chewers ages 14–25. *Nicotine Tob Res* 2013; 15: 1502–1510.
 44. Elfeddali I, Bolman C, Candel MJ, et al. Preventing smoking relapse via Web-based computer-tailored feedback: A randomized controlled trial. *J Med Internet Res* 2012; 14: e109.
 45. Houston TK, Sadasivam RS, Allison JJ, et al. Evaluating the QUIT-PRIMO clinical practice ePortal to increase smoker engagement with online cessation interventions: A national hybrid type 2 implementation study. *Implement Sci* 2015; 10: 154.
 46. Muñoz RF, Barrera AZ, Delucchi K, et al. International Spanish/English Internet smoking cessation trial yields 20% abstinence rates at 1 year. *Nicotine Tob Res* 2009; 11: 1025–1034.
 47. Ramo DE, Thrul J, Chavez K, et al. Feasibility and quit rates of the tobacco status project: A Facebook smoking cessation intervention for young adults. *J Med Internet Res* 2015; 17: e291.
 48. Severson HH, Gordon JS, Danaher BG, et al. ChewFree.com: Evaluation of a Web-based cessation program for smokeless tobacco users. *Nicotine Tob Res* 2008; 10: 381–391.
 49. Stanczyk NE, Crutzen R, Bolman C, et al. Influence of delivery strategy on message-processing mechanisms and future adherence to a Dutch computer-tailored smoking cessation intervention. *J Med Internet Res* 2013; 15: e28.
 50. Stoddard JL, Augustson EM and Moser RP. Effect of adding a virtual community (bulletin board) to smoke-free.gov: randomized controlled trial. *J Med Internet Res* 2008; 10: e53.
 51. Stoops WW, Dallery J, Fields NM, et al. An internet-based abstinence reinforcement smoking cessation intervention in rural smokers. *Drug Alcohol Depend* 2009; 105: 56–62.
 52. Lieberman DZ. Effects of a personified guide on adherence to an online program for alcohol abusers. *Cyberpsychol Behav* 2006; 9: 603–607.
 53. Schulz DN, Candel MJ, Kremers SP, et al. Effects of a Web-based tailored intervention to reduce alcohol consumption in adults: Randomized controlled trial. *J Med Internet Res* 2013; 15: e206.
 54. Schaub MP, Wenger A, Berg O, et al. A web-based self-help intervention with and without chat counseling to reduce cannabis use in problematic cannabis users: Three-arm randomized controlled trial. *J Med Internet Res* 2015; 17: e232.
 55. Swan GE, McClure JB, Jack LM, et al. Behavioral counseling and varenicline treatment for smoking cessation. *Am J Prev Med* 2010; 38: 482–490.
 56. Stoddard JL, Augustson EM and Moser RP. Effect of adding a virtual community (bulletin board) to smoke-free.gov: Randomized controlled trial. *J Med Internet Res* 2008; 10: e53.
 57. Te Poel F, Bolman C, Reubsat A, et al. Efficacy of a single computer-tailored e-mail for smoking cessation: Results after 6 months. *Health Educ Res* 2009; 24: 930–940.
 58. Berg IK, Miller SD, Hofmeister S, et al. *Kurzzeittherapie bei Alkoholproblemen: Ein lösungsorientierter Ansatz*. Heidelberg: Carl-Auer-Systeme, 1998.
 59. Miller WR and Rollnick S. *Motivational interviewing: Helping people change*. New York: Guilford Press, 2012.
 60. Steinberg KL, Roffman RA, Carroll KM, et al. *Brief counseling for marijuana dependence: A manual for treating adults*. Rockville, MD: Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration, 2005.

61. Rockville, MD: Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration; 2005.
62. Severson H, Gordon J, Danaher B, et al. ChewFree.com: Evaluation of a web-based cessation program for smokeless tobacco users. *Nicotine Tob Res* 2008; 10: 381–391.
63. Couper MP, Alexander GL, Zhang N, et al. Engagement and retention: Measuring breadth and depth of participation use of an online intervention. *J Med Internet Res* 2010; 12: 41–55.
64. Milward J, Khadjesari Z, Fincham-Campbell S, et al. User preferences for content, features, and style for an app to reduce harmful drinking in young adults: Analysis of user feedback in app stores and focus group interviews. *JMIR Mhealth Uhealth* 2016; 4: e47.
65. Milward J, Lynskey M and Strang J. Solving the problem of non-attendance in substance abuse services. *Drug Alcohol Rev* 2014; 33: 625–636.
66. Gurolo-Urganci I, de Jongh T, Vodopivec-Jamsek V, et al. Mobile phone messaging reminders for attendance at healthcare appointments. *Cochrane Database Syst Rev* 2013; 12: CD007458.
67. Crane D, Garnett C, Brown J, et al. Factors influencing usability of a smartphone app to reduce excessive alcohol consumption: Think aloud and interview studies. *Front Public Health* 2017; 5: e39.
68. Milward J, Deluca P, Drummond C, et al. Usability testing of the BRANCH smartphone app designed to reduce harmful drinking in young adults. *JMIR Mhealth Uhealth* 2017; 5(8): e109.
69. McCarthy O, Carswell K, Murray E, et al. What young people want from a sexual health website: Design and development of Sexunzipped. *J Med Internet Res* 2012; 14: e127.
70. Cafazzo JA, Casselman M, Hamming N, et al. Design of an mHealth app for the self-management of adolescent type 1 diabetes: A pilot study. *J Med Internet Res* 2012; 14: e70.
71. Abraham C and Michie S. A taxonomy of behavior change techniques used in interventions. *Health Psychol* 2008; 27: 379.
72. Benishek LA, Dugosh KL, Kirby KC, et al. Prize-based contingency management for the treatment of substance abusers: a meta-analysis. *Addiction* 2014; 109: 1426–1436.
73. Oinas-Kukkonen H and Harjuma M. Persuasive systems design: Key issues, process model, and system features. *Comm Assoc Inform Syst* 2009; 24: 28.
74. Hamari J, Koivisto J and Pakkanen T. Do persuasive technologies persuade? A review of empirical studies. *Persuasive technology*. Cham, Germany: Springer, 2014, pp.118–136.
75. Kelders SM, Kok RN, Ossebaard HC, et al. Persuasive system design does matter: A systematic review of adherence to web-based interventions. *J Med Internet Res* 2012; 14: e152.
76. Ryan C, Bergin M and Wells JS. *Int.J. Behav. Med.* (2017). <https://doi.org/10.1007/s12529-017-9678-8> (accessed 15 January 2015).
77. Crane D, Garnett C, Brown J, et al. Behavior change techniques in popular alcohol reduction apps: Content analysis. *J Med Internet Res* 2015; 17(5): e118.
78. Short CE, Rebar AL, Plotnikoff RC, et al. Designing engaging online behaviour change interventions: A proposed model of user engagement. *Eur Health Psychol* 2015; 17: 32–38.
79. Yardley L, Spring BJ, Riper H, et al. Understanding and promoting effective engagement with digital behavior change interventions. *Am J Prev Med* 2016; 51: 833–842.
80. Donkin L, Christensen H, Naismith SL, et al. A systematic review of the impact of adherence on the effectiveness of e-therapies. *J Med Internet Res* 2011; 13: e52.
81. O'Brien HL and Toms EG. What is user engagement? A conceptual framework for defining user engagement with technology. *J Am Soc Inform Sci Technol* 2008; 59: 938–955.
82. Noblit GW and Hare RD. *Meta-ethnography: Synthesizing qualitative studies*. London: SAGE Publications, 1988.
83. Light RJ and Pillemer DB. *Summing up: The science of reviewing research*. Cambridge, MA: Harvard University Press, 1984.
84. Begley CM. Using triangulation in nursing research. *J Adv Nurs* 1996; 24: 122–128.

Appendix 1. Medline search terms for engagement promoting strategies for online substance-use interventions

Terms for online interventions

(phone adj application*) or (phone adj app) or smartphone.mp,hw.
 (virtual adj reality).mp,hw.
 (Second life or secondlife).mp,hw.
 (online adj (bulletin adj board*) or bulletinboard* or messageboard* or (message adj board*)).mp,hw.
 (online or on-line).mp,hw.
 exp internet/ or ((internet adj based) or internet-based).mp,hw.
 ((web adj based) or web-based).mp,hw.
 ((world adj wide adj web) or (world-wide-web) or WWW or (world-wide adj web) or (worldwide adj web) or website*).mp,hw.
 ((electronic adj mail) or e-mail* or email*).mp,hw.
 (((mobile or cellular or cell or smart) adj (phone* or telephone*)) or smartphone).mp,hw.
 (internet based or internet-based) adj10 (computer* or laptop*).mp,hw.
 (web adj based or web-based) adj10 (computer* or laptop*).mp,hw.
 (World wide web or world-wide-web or www or world-wide web or worldwide web or

website*).mp,hw.
 (E-health or ehealth or electronic health).mp,hw.
 (m-health or mobile health or mobilehealth or mhealth or tablet).mp,hw.
 (cyberpsychology or cybertherap* or etherap* or ecounsel*).mp,hw.
 (digital intervention* or mobile device*).mp,hw.
 Exp Social media/
 (Facebook OR LinkedIn OR Twitter OR Badoo OR Orkut OR Qzone OR Xing OR Tencent OR Weibo OR Mixi OR Sina Weibo OR Hyves OR Skyrock OR Odnoklassniki OR Wer-kennt-wen OR V Kontakte OR Tuenti OR MySpace).mp,hw.
 (interactive adj ((health adj communicat*) or televis* or video* or technolog* or multimedia)).mp,hw.
 (blog* or web-log* or weblog*).mp,hw.
 ((chat adj room*) or chatroom*).mp,hw.

AND**Engagement Outcomes**

Engagement.mp,hw.
 Attrition.mp,hw.
 Adherence.mp,hw.
 Exp Patient Dropouts/
 Exp patient compliance/
 (process adj metric*).mp,hw.
 website-us*.mp.
 website us*.mp.
 (Web us* or web-us*).mp.
 (website adj5 usage).mp,hw.
 (web adj5 usage).mp,hw.
 (online adj5 us*).mp,hw.
 Usage.mp,hw.
 Login*.mp,hw.
 log-ins.mp,hw.
 (page* adj5 view*).mp,hw.
 (module* adj complet*).mp,hw.
 (session* adj complet*).mp,hw.
 (visit* adj5 website*).mp,hw.
 ((Intervention adj participation) or (website adj participation) or (online adj participation) or (web* adj section* adj open*) or (web adj page* adj visit*) or (time adj spent adj online)).mp,hw.

AND**Engagement promoting strategies**

(prompt* or reminder* or (push adj notification*)).mp,hw.
 ((chat adj room*) or chatroom*).mp,hw.
 ((bulletin adj board*) or bulletinboard* or message-board* or (message adj board*)). mp,hw.
 ((social adj support) or (peer adj support) or (online adj social adj network*) or customi\$ation or (social adj connectivity) or (social adj5 network) or (social adj media) or (online adj5 communit*) or (chat adj counsel*)).mp,hw.
 (social adj bookmark*) or (social adj technolog*) or (social adj networking adj site*).mp,hw.
 tailor* or personali* or relevan* or individuali* or feedback or (personali\$ed adj feedback).mp,hw.
 Gamification or (gam* adj strateg*) or (gam* adj design).mp,hw.
 Incentive* or reinforcement* or reward*.mp,hw.
 (goal adj setting).mp,hw.
 monitor*.mp,hw.
 (leader adj board) or (leaderboard) or (achievement adj badge*) or (progress adj report) or (progress adj chart).mp,hw.
 (persuasive adj design adj system*).mp,hw.
 Usability.mp,hw.
 Novelty.mp,hw.
 Aesthetic*.mp,hw.
Medline RCT Filter - Cochrane
 randomized controlled trial.pt.
 controlled clinical trial.pt.
 randomized.ab.
 placebo.ab.
 drug therapy.fs.
 randomly.ab.
 trial.ab.
 groups.ab.
 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8
 exp animals/not humans.sh.
 9 not 10

Appendix 2. Tools and techniques for narrative synthesis

Developing a preliminary synthesis	Description of tool	Included in study	Detail of inclusion
Textual description	Descriptive paragraph of study/ intervention	Yes	Detailed description of each EPSs will be presented
Groupings and clusterings	Organising studies into manageable groups	Yes	Studies will be organised by EPS type as a means of synthesising the EPS results
Tabulation	Visual presentation of qualitative and quantitative data	Yes	Study results and characteristics will be tabulated
Transforming data into a common rubric	Data transformed into common rubric, for example, odds ratio for a meta-analysis	No	This is not possible as outcome and EPS types are too heterogeneous to include in a meta-analysis
Vote counting as a descriptive tool	Calculating frequencies of different results across different studies	Yes	Statistically significant results will be counted for an overview of effect. Quality of studies will be considered. This will be completed as a tool for exploring relationships (see below)
Translating data: thematic analysis	Translating the data into common themes across studies	No	More appropriate for qualitative data
Translating data: content analysis	Translating data into frequencies based on coding rules	No	More appropriate for qualitative data
Graphs, frequency distributions, funnel plots, forest plots and L'Abbe plots	Graphically present relationships within and between studies	No	Quantitative data not being used
Moderator variables and subgroup analyses	Examining characteristics between and within studies to explain variability in primary results	Yes	Variations between the EPS features will be examined. Populations, motivations to quit/reduce will be discussed. A table showing the EPS components of the evaluated interventions will be included
Idea webbing and concept mapping	Create visual models to conceptualise and explore connections across studies	No	Connections across studies will be explored with textual descriptions
Translation as an approach to exploring relationships ⁸⁵	Using qualitative research techniques to synthesise findings from multiple studies	No	More appropriate for qualitative data
Qualitative case description ⁸⁶	Use of descriptive data to explain differences in statistical findings	Yes	Will be used in conjunction with textual descriptions to explore the data
Investigator and methodological triangulation ⁸⁷	Analysing data in relation to the context in which it was produced, for example, the disciplinary perspectives and expertise of the researchers	No	More appropriate for qualitative data

Appendix 3. Quality rating of included studies

Study	Adequate sequence generation	Adequate concealment	Adequate blinding of participants and personnel	Adequate blinding of engagement outcome	Incomplete engagement outcome data addressed	Free of selective reporting	Free of other bias	Quality score
Elfeddali et al. 2012	Yes	Yes	Unclear	Yes	Unclear	Unclear	No	4/7
Danaher et al. 2012	Yes	Yes	Unclear	Yes	Unclear	Unclear	No	4/7
Houston et al. 2015	Yes	Yes	Yes	Yes	Unclear	Yes	No	6/7
Lieberman et al. 2006	Unclear	Unclear	Unclear	Yes	Unclear	Unclear	No	2/7
McClure et al. 2013	Yes	Yes	Yes	Yes	Yes	Unclear	No	5/7
Munoz et al. 2009	Yes	Yes	No	Yes	Yes	Unclear	No	4/7
Ramo et al. 2015	Unclear	Unclear	Unclear	Yes	Yes	Unclear	No	2/3
Schaub et al. 2015	Yes	Yes	No	Yes	Yes	Unclear	No	4/7
Schulz et al. 2013	Unclear	Unclear	No	Yes	Yes	Unclear	No	2/7
Severson et al. 2008	Unclear	Yes	Unclear	Yes	Yes	Unclear	No	2/7
Stanczyk et al. 2013	Yes	Yes	Unclear	Yes	No	Unclear	No	3/7
Strecher et al. 2008	Unclear	Yes	Yes	Yes	Yes	Yes	No	5/7
Stoddard et al. 2008	Yes	Yes	Yes	Yes	Yes	Yes	No	6/7
Stoops et al. 2009	Yes	Yes	No	No	Unclear	Unclear	No	3/7
Tensil et al. 2013	Yes	Yes	Yes	Yes	Yes	Unclear	No	5/7